

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 June 2001 (07.06.2001)

PCT

(10) International Publication Number
WO 01/40217 A1

(51) International Patent Classification⁷: **C07D 401/14**,
A61K 31/47, A61P 35/00, C07D 401/04, 491/10, 471/04,
491/04, 413/14

LYSSIKATOS, Joseph, Peter [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US).

(21) International Application Number: PCT/IB00/01636

(74) Agents: **SPIEGEL, Allen, J. et al.**; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).

(22) International Filing Date:
10 November 2000 (10.11.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/168,217 30 November 1999 (30.11.1999) US

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **PFIZER PRODUCTS INC.** [US/US]; Eastern Point Road, Groton, CT 06340 (US).

(72) Inventors; and


Published:

(75) Inventors/Applicants (*for US only*): **BARTH, Wayne, Ernest** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **LUZZIO, Michael, Joseph** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US).

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL BENZOIMIDAZOLE DERIVATIVES USEFUL AS ANTIPROLIFERATIVE AGENTS



(1)

(57) Abstract: The invention relates to compounds of formula (1) and to pharmaceutically acceptable salts, prodrugs and solvates thereof, wherein R¹, R⁷, R⁸, R⁹, R¹⁰, and R¹¹ are as defined herein. The invention also relates to methods of treating abnormal cell growth, such as cancer, in mammals by administering the compounds of formula 1 and to pharmaceutical compositions for treating such disorders which contain the compounds of formula (1). The invention also relates to methods of preparing the compounds of formula (1).

WO 01/40217 A1

5 NOVEL BENZOIMIDAZOLE DERIVATIVES USEFUL AS ANTIPROLIFERATIVE AGENTS

Background of the Invention

This invention relates to novel benzimidazole derivatives that are useful in the treatment of abnormal cell growth, such as cancer, in mammals. This invention also relates to a method of using such compounds in the treatment of abnormal cell growth in mammals, especially
10 humans, and to pharmaceutical compositions containing such compounds.

It is known that a cell may become cancerous by virtue of the transformation of a portion of its DNA into an oncogene (i.e., a gene which, on activation, leads to the formation of malignant tumor cells). Many oncogenes encode proteins that are aberrant tyrosine kinases capable of causing cell transformation. Alternatively, the overexpression of a normal proto-
15 oncogenic tyrosine kinase may also result in proliferative disorders, sometimes resulting in a malignant phenotype.

Receptor tyrosine kinases are enzymes which span the cell membrane and possess an extracellular binding domain for growth factors such as epidermal growth factor, a transmembrane domain, and an intracellular portion which functions as a kinase to
20 phosphorylate specific tyrosine residues in proteins and hence to influence cell proliferation. Other receptor tyrosine kinases include c-erbB-2, c-met, tie-2, PDGFr, FGFr, and VEGFR. It is known that such kinases are frequently aberrantly expressed in common human cancers such as breast cancer, gastrointestinal cancer such as colon, rectal or stomach cancer, leukemia, and ovarian, bronchial or pancreatic cancer. It has also been shown that epidermal growth factor
25 receptor (EGFR), which possesses tyrosine kinase activity, is mutated and/or overexpressed in many human cancers such as brain, lung, squamous cell, bladder, gastric, breast, head and neck, oesophageal, gynecological and thyroid tumors.


Accordingly, it has been recognized that inhibitors of receptor tyrosine kinases are useful as selective inhibitors of the growth of mammalian cancer cells. For example, erbstatin, a
30 tyrosine kinase inhibitor, selectively attenuates the growth in athymic nude mice of a transplanted human mammary carcinoma that expresses epidermal growth factor receptor tyrosine kinase (EGFR) but is without effect on the growth of another carcinoma that does not express the EGF receptor. Thus, the compounds of the present invention, which are selective inhibitors of certain receptor tyrosine kinases, in particular PDGFr, are useful in the treatment of
35 abnormal cell growth, in particular cancer, in mammals.

Various other compounds, such as styrene derivatives, have also been shown to possess tyrosine kinase inhibitory properties. More recently, five European patent publications, namely EP 0 566 226 A1 (published October 20, 1993), EP 0 602 851 A1 (published June 22, 1994), EP 0 635 507 A1 (published January 25, 1995), EP 0 635 498 A1 (published January 25,
40 1995), and EP 0 520 722 A1 (published December 30, 1992), refer to certain bicyclic derivatives,

5 in particular quinazoline derivatives, as possessing anti-cancer properties that result from their tyrosine kinase inhibitory properties. Also, World Patent Application WO 92/20642 (published November 26, 1992), refers to certain bis-mono and bicyclic aryl and heteroaryl compounds as tyrosine kinase inhibitors that are useful in inhibiting abnormal cell proliferation. World Patent Applications WO96/16960 (published June 6, 1996), WO 96/09294 (published March 6, 1996),
 10 WO 97/30034 (published August 21, 1997), WO 98/02434 (published January 22, 1998), WO 98/02437 (published January 22, 1998), and WO 98/02438 (published January 22, 1998), also refer to substituted bicyclic heteroaromatic derivatives as tyrosine kinase inhibitors that are useful for the same purpose.

Summary of the Invention

15 The present invention relates to compounds of the formula 1



and to pharmaceutically acceptable salts, prodrugs and solvates thereof, wherein:

X is CH or N;

R¹ is selected from $-(CR^4R^5)_tC(O)OR^3$, $-(CR^4R^5)_tC(O)NR^3R^4$, $-(CR^4R^5)_tOR^3$,
 20 $-(CR^4R^5)_tC(O)(C_3-C_{10} \text{ cycloalkyl})$, $-(CR^4R^5)_tC(O)(C_6-C_{10} \text{ aryl})$, $-(CR^4R^5)_tC(O)(4 \text{ to } 10 \text{ membered heterocyclic})$, $-(CR^4R^5)_t(C_3-C_{10} \text{ cycloalkyl})$, $-(CR^4R^5)_t(C_6-C_{10} \text{ aryl})$, and $-(CR^4R^5)_t(4 \text{ to } 10 \text{ membered heterocyclic})$, wherein each t is independently an integer from 0 to 5; said cycloalkyl, aryl and heterocyclic R¹ moieties are optionally fused to a benzene ring, a C₅-C₈ cycloalkyl group, or a 4 to 10 membered heterocyclic group; the $-(CR^4R^5)_t$ moieties of the
 25 foregoing R¹ groups optionally include a carbon-carbon double or triple bond where t is an integer between 2 and 5; the foregoing R¹ groups are each optionally substituted by 1 or 2 groups independently selected from $-NR^3R^4$, $-OR^3$, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, and C₂-C₁₀ alkynyl, wherein said alkyl, alkenyl and alkynyl groups are substituted by 1 or 2 groups independently selected from $-NR^3R^4$ and $-OR^3$; and the foregoing R¹ groups are optionally
 30 substituted by 1 to 3 R² groups;

each R² is independently selected from H, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₁₀ cycloalkyl, oxo, halo, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, $-OR^3$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)OR^6$, $-OC(O)R^3$, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-NR^4C(O)R^3$, $-C(O)NR^3R^4$, $-NR^5C(O)NR^3R^4$, $-NR^3R^4$, $-S(O)_j(CR^4R^5)_m(C_6-C_{10} \text{ aryl})$, $-S(O)_j(C_1-C_6 \text{ alkyl})$, wherein j is an integer from 0 to 2, $-(CR^4R^5)_m(C_6-C_{10} \text{ aryl})$, $-O(CR^4R^5)_m(C_6-C_{10} \text{ aryl})$,
 35

5 -NR⁴(CR⁴R⁵)_m(C₆-C₁₀ aryl), -O(CR⁴R⁵)_m(4 to 10 membered heterocycle), -NR⁴(CR⁴R⁵)_m(4 to 10 membered heterocycle), -(CR⁴R⁵)_m(4 to 10 membered heterocyclic), and -(CR⁴R⁵)_m(C₃-C₁₀ cycloalkyl) wherein each m is independently an integer from 0 to 4; said alkyl, alkenyl and alkynyl groups optionally contain 1 or 2 hetero moieties selected from O, -S(O)_j- wherein j is an integer from 0 to 2, and -N(R³)- with the proviso that two O atoms, two S atoms, or an O and S atom are not attached directly to each other, and the proviso that an O atom, a S atom or a N atom are not attached directly to a triple bond or a non-aromatic double bond; said cycloalkyl, aryl and heterocyclic R² groups are optionally fused to a C₆-C₁₀ aryl group, a C₅-C₈ cycloalkyl group, or a 4 to 10 membered heterocyclic group; and said alkyl, cycloalkyl, aryl and heterocyclic R² groups are optionally substituted by 1 to 5 substituents independently
 10 selected from oxo (=O), halo, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -NR⁴SO₂R⁶, -SO₂NR³R⁴, -C(O)R³, -C(O)OR³, -OC(O)R³, -NR⁴C(O)OR⁶, -NR⁴C(O)R³, -C(O)NR³R⁴, -NR³R⁴, -OR³, C₁-C₁₀ alkyl, -(CR⁴R⁵)_m(C₆-C₁₀ aryl), and -(CR⁴R⁵)_m(4 to 10 membered heterocyclic), wherein each m is independently an integer ranging from 0 to 4;

each R³ is independently selected from H, C₁-C₁₀ alkyl, -(CR⁴R⁵)_m(C₆-C₁₀ aryl), and
 20 -(CR⁴R⁵)_m(4 to 10 membered heterocyclic), wherein each m is independently an integer from 0 to 4; said alkyl group optionally includes 1 or 2 hetero moieties selected from O, -S(O)_j- wherein j is an integer ranging from 0 to 2, and -N(R⁴)- with the proviso that two O atoms, two S atoms, or an O and S atom are not attached directly to each other; said cycloalkyl, aryl and heterocyclic R³ groups are optionally fused to a C₆-C₁₀ aryl group, a C₅-C₈ cycloalkyl group, or
 25 a 4 to 10 membered heterocyclic group; and the foregoing R³ substituents, except H, are optionally substituted by 1 to 5 substituents independently selected from oxo, halo, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -C(O)R⁴, -C(O)OR⁴, -OC(O)R⁴, -NR⁴C(O)R⁵, -C(O)NR⁴R⁵, -NR⁴R⁵, hydroxy, C₁-C₆ alkyl, and C₁-C₆ alkoxy;

each R⁴ and R⁵ is independently H or C₁-C₆ alkyl;
 30 or, where R⁴ and R⁵ are attached to the same carbon or nitrogen atom, R⁴ and R⁵ together with said carbon or nitrogen may be taken together to form a 4 to 10 membered ring which may be carbocyclic or heterocyclic;

each R⁶ is selected from the substituents provided in the definition of R³ except R⁶ is not H;

35 each R⁷, R⁸, R⁹, R¹⁰ and R¹¹ is independently selected from the group of substituents provided in the definition of R².

In a specific embodiment of the present invention R¹ is C₆-C₁₀ aryl or 4 to 10 membered heterocyclic wherein the foregoing R¹ groups are each substituted by 1 or 2 groups independently selected from -NR³R⁴, -OR³ and C₁-C₃ alkyl, wherein said alkyl groups

5 are substituted by 1 or 2 groups independently selected from $-NR^3R^4$ and $-OR^3$; and the foregoing R^1 groups are optionally substituted by 1 to 3 R^2 groups;

each R^2 is independently selected from H, C_1 - C_{10} alkyl, C_3 - C_{10} cycloalkyl, oxo(=O), $-OR^3$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)R^3$, $-C(O)NR^3R^4$, $-NR^3R^4$, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-(CR^4R^5)_m$ (4 to 10 membered heterocyclic), and $(CR^4R^5)_m$ (C_3 - C_{10} cycloalkyl); and said alkyl
 10 groups optionally contain 1 or 2 hetero moieties selected from O, $-S(O)_j$ - wherein j is an integer from 0 to 2, and $-N(R^3)-$ with the proviso that two O atoms, two S atoms, or an O and S atom are not attached directly to each other; and said alkyl and cycloalkyl R^2 groups are optionally substituted by 1 to 5 substituents independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)OR^6$,
 15 $-NR^4C(O)R^3$, $-C(O)NR^3R^4$, $-NR^3R^4$, $-OR^3$, and C_1 - C_{10} alkyl, wherein each m is independently an integer ranging from 0 to 4.

In another specific embodiment of the present invention, R^1 is piperidinyl, piperziny, or phenyl, wherein said R^1 groups are substituted by $-NR^3R^4$, oxo (=O), $-OR^3$ and C_1 - C_3 alkyl, wherein said alkyl groups are substituted by 1 or 2 groups independently selected from
 20 $-NR^3R^4$ and $-OR^3$; and the foregoing R^1 groups are optionally substituted by 1 to 3 R^2 groups. In a more specific embodiment, said R^1 groups are substituted by $-NR^3R^4$, oxo (=O), OR^3 , or C_1 - C_3 alkyl, wherein said alkyl group are substituted by $-NR^3R^4$.

In another specific embodiment of the present invention, R^1 is phenyl substituted by pyrrolidin-1-yl which pyrrolidin-1-yl is optionally substituted by 1 to 3 substituents
 25 independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)OR^6$, $-NR^4C(O)R^3$, $-C(O)NR^3R^4$, $-NR^3R^4$, $-OR^3$, and C_1 - C_{10} alkyl; and R^{11} is $-OR^3$. More specifically, R^1 is 4-pyrrolidin-1-ylmethyl-phenyl optionally substituted by 1 to 3 substituents independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)OR^6$, $-NR^4C(O)R^3$,
 30 $-C(O)NR^3R^4$, $-NR^3R^4$, $-OR^3$, and C_1 - C_{10} alkyl; and R^{11} is $-OR^3$. More specifically, R^{11} is connected at the 5 position of the benzimidazole moiety of the compound of formula 1 and is $-OR^3$. More specifically, R^{11} is connected at the 5 position of the benzimidazole moiety of the compound of formula 1 and is 2-methoxyethoxy.

In another specific embodiment of the present invention, R^1 is pyrrolidin-1-yl or
 35 piperidin-1-yl, said R^1 being optionally substituted by 1 to 3 substituents independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)OR^6$, $-NR^4C(O)R^3$, $-C(O)NR^3R^4$, $-NR^3R^4$, $-OR^3$, and C_1 - C_{10} alkyl. More specifically, R^1 is pyrrolidin-1-yl or piperidin-1-yl substituted by $-NR^3R^4$ and optionally substituted by 1 or 2 substituents independently selected from oxo, cyano, trifluoromethyl,
 40 trifluoromethoxy, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)OR^6$, $-NR^4C(O)R^3$,

- 5 -C(O)NR³R⁴, -NR³R⁴, -OR³, and C₁-C₁₀ alkyl; and R¹¹ is -OR³. More specifically, R¹¹ is connected at the 5 position of the benzimidazole moiety of the compound of formula 1 and is -OR³ and R⁹ and R¹⁰ are both H. More specifically, R¹¹ is connected at the 5 position of the benzimidazole moiety of the compound of formula 1 and is 2-methoxyethoxy and R⁹ and R¹⁰ are both H.
- 10 Preferred compounds include those selected from the group consisting of:
 [1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
 1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazol-5-ol;
 1-[2-[5-(Pyridin-2-ylmethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
 {1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethyl-amine;
 15 {4-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine;
 {4-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine;
 Cyclopropyl-{4-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine;
 tert-Butyl-{4-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine;
 4-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzylamine;
 20 1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
 {1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethylamine;
 1-[2-(5-Trifluoromethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
 {4-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine;
 Cyclopropyl-{4-[2-(5-ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine;
 25 tert-Butyl-{4-[2-(5-ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine;
 {4-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine;
 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one;
 1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one;
 1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
 30 tert-Butyl-{1-[2-(5-ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-amine;
 {1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine;
 2-(5-Methoxy-benzimidazol-1-yl)-8-(1-oxa-6-aza-spiro[2.5]oct-6-yl)-quinoline;
 4-Dimethylaminomethyl-1-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
 35 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-4-methylaminomethyl-piperidin-4-ol;
 4-Aminomethyl-1-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-pyrrolidin-3-ylamine;
 1-(2-Benzimidazol-1-yl-quinolin-8-yl)-piperidin-4-ylamine;
 40 1-(2-Imidazo[4,5-b]pyridin-3-yl-quinolin-8-yl)-piperidin-4-ylamine;

- 5 1-{2-[5-(4-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
 1-{2-[5-Cyclopropylmethoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
 1-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
 1-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 10 1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
 1-{2-[5-(Pyridin-4-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
 1-{2-[5-Benzylloxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
 1-{2-[5-(Pyridin-3-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 15 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
 1-{2-[5-Methoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidine-4-carboxylic acid ethyl ester;
 1-{2-[5-Methoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidine-4-carboxylic acid;
 4-Dimethylaminomethyl-1-{2-[5-methoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-
- 20 4-ol;
 N-{1-{2-[5-Methoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl}-acetamide;
 N-{1-{2-[5-Cyclopropylmethoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl}-acetamide;
 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol;
- 25 {1-{2-[5-Methoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl}-urea;
 4-Aminomethyl-1-{2-[5-(pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol;
 Cyclopropyl-(1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-amine;
- 30 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-dimethyl-amine;
 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methyl-amine;
- 35 (1-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-dimethyl-amine;
 {1-{2-[5-Cyclopropylmethoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methyl-amine;
 {1-{2-[5-Cyclopropylmethoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-dimethyl-amine;

- 5 2-Amino-N-(1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-acetamide;
 -(S)-2-Amino-N-(1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-propionamide;
 -(R)-2-Amino-N-(1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-propionamide;
- 10 2-Amino-N-(1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-isobutyramide;
 1-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamino)-2-methyl-propan-2-ol;
- 15 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-pyridin-2-ylmethyl-amine;
 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-pyridin-3-ylmethyl-amine;
 4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
- 20 [2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenoxy)-ethyl]-dimethyl-amine;
 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline;
 [2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethyl]-dimethyl-amine;
- 25 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyridin-2-ylmethyl-piperazin-1-yl)-quinoline;
 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyridin-3-ylmethyl-piperazin-1-yl)-quinoline;
 2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-2-methyl-propan-1-one;
- 30 (S)-2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-one;
 (S)-2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-one;
- 35 2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone;
 (1-Amino-cyclopropyl)-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-methanone;
 2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethylamine;
- 40

- 5 (R)-2-Amino-3-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-ol;
3-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-3-aza-bicyclo[3.1.0]hex-6-ylamine;
(S)-1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-pyrrolidin-3-ylamine;
- 10 (R)-1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-pyrrolidin-3-ylamine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-pyridin-3-yl-quinoline;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(6-methoxy-pyridin-3-yl)-quinoline;
4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzoic acid methyl
15 ester;
1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-4-methyl-piperidin-4-ylamine;
1-[2-(6,7-Dihydro-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalen-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
- 20 2-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethanol;
4-Cyclopropylaminomethyl-1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol;
1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazole-5-sulfonic acid
dimethylamide;
- 25 1-[2-(6-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-[2-(5,6-Dimethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
2-Dimethylamino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone;
1-[2-(5-Benzoyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-4-methyl-piperidin-4-ol;
- 30 (4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-dimethylamine;
(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-methylamine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-morpholin-4-ylmethyl-phenyl)-quinoline;
- 35 2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamino)-ethanol;
4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;

- 5 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-cis-
10 pyrrolidine-3,4-diol;
R,R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol);
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol;
15 R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol);
S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol);
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-azetidin-3-ol;
20 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-quinoline;
4-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperazine-1-carboxylic acid tert-butyl ester;
[1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-yl]-carbamic acid tert-butyl ester;
25 1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-ylamine;
(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methanol;
30 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-methyl-amine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-piperidin-1-yl]-quinoline;
(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-
35 dimethyl-amine;
1-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-pyrrolidin-3-ol;
C-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methylaniline;

- 5 1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-4-methyl-piperidin-4-ol;
1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol;
S,S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol);
- 10 4-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
4-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
1-[2-(5-Phenyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-[2-(5-Pyridin-4-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-{2-[5-(3-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 15 1-[2-(5-Pyridin-3-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-{2-[5-(6-Methoxy-pyridin-3-yl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-{2-[5-(4-Aminomethyl-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 20 4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl}-benzoic acid methyl ester;
4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl}-phenol;
2-(5-Methoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinoline-8-carboxylic acid methyl ester;
2-(5-Methoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid (2-dimethylamino-ethyl)-amide;
- 25 2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester;
[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-pyrrolidin-1-yl-methanone;
- 30 [2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-morpholin-4-yl-methanone;
[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-1-yl-methanone;

- 5 (3-Amino-pyrrolidin-1-yl)-[2-(5-cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-methanone;
- 8-Allyloxy-2-(5-methoxy-benzimidazol-1-yl)-quinoline;
- {2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethyl}-methyl-amine;
- {2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethyl}-dimethyl-amine;
- 10 2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethylamine;
- 1-[[2-[5-(3-Morpholinoethoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride;
- 1-[[2-[5-(3-Morpholinoethoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride;
- 15 5-{2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-[1,3,4]oxadiazol-2-ylamine;
- Ethyl 1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate;
- 1-[8-(4-Aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylic acid;
- N-(4-Morpholino)ethyl-1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-
- 20 carboxamide;
- 4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1*H*-benzimidazol-5-yl}-benzaldehyde;
- 1-{2-[5-(4-Methylaminomethyl-phenyl)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 1-{2-[5-(4-Dimethylaminomethyl-phenyl)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-
- 25 4-ylamine;
- 1-(2-{5-[2-(2-Methyl-imidazol-1-yl)-ethoxy]-benzimidazol-1-yl}-quinolin-8-yl)-piperidin-4-ylamine and
- 1-{2-[5-(2-[1,2,4]Triazol-1-yl)-ethoxy]-benzimidazol-1-yl}-quinolin-8-yl}-piperidin-4-ylamine,
- 30 and the pharmaceutically acceptable salts, prodrugs and solvates of the foregoing compounds.
- In accordance with the present invention, preferred compounds include those selected from the group consisting of:
- 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;

- 5 1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-ol;
1-{2-[5-(Pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
{1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethyl-amine;
{4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine;
{4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine;
- 10 1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
{1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethylamine;
{4-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine;
{4-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine;
{1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine;
- 15 4-Dimethylaminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-4-methylaminomethyl-piperidin-4-ol;
4-Aminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
- 20 1-{2-[5-(4-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 25 1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-{2-[5-(Pyridin-4-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-[2-(5-Benzoyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-{2-[5-(Pyridin-3-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 30 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
4-Dimethylaminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
4-Aminomethyl-1-[2-[5-(pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ol;
- 35 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl}-dimethyl-amine;
1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl}-methyl-amine;
(1-[2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-dimethyl-amine;
- 40

- 5 {1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine;
{1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethyl-amine;
10 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-pyridin-2-ylmethyl-amine;
(1-[2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-pyridin-3-ylmethyl-amine;
4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
15 [2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenoxy)-ethyl]-dimethyl-amine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline;
[2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethyl]-dimethyl-amine;
2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-2-methyl-propan-1-one;
20 (S)-2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-one;
(S)-2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-one;
25 2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone;
2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethylamine;
30 3-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-3-aza-bicyclo[3.1.0]hex-6-ylamine;
1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-4-methyl-piperidin-4-ylamine;
2-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethanol;
1-[2-(5,6-Dimethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
35 1-[2-(6,7-Dihydro-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalen-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-dimethyl-amine;
(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-methyl-amine;

- 5 2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamino)-ethanol;
4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
- 10 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-cis-pyrrolidine-3,4-diol;
- 15 R,R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol);
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol;
- 20 R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol);
S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol);
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-azetidin-3-ol
- 25 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-quinoline;
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-ylamine;
(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methanol;
- 30 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-methyl-amine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-piperidin-1-yl]-quinoline;
- 35 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-dimethyl-amine;
C-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methylamine;

- 5 S,S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol);
- 4-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
- 1-[2-(5-Phenyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
- 1-[2-(5-Pyridin-4-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
- 10 1-[2-[5-(3-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
- 1-[2-(5-Pyridin-3-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
- 1-[2-[5-(6-Methoxy-pyridin-3-yl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
- 1-[2-[5-(4-Aminomethyl-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
- 15 4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl}-benzoic acid methyl ester;
- 4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl}-phenol;
- 1-[[2-[5-(3-Morpholinoethoxy)-1H-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride;
- 20 Ethyl 1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate;
- N-(4-Morpholino)ethyl-1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxamide;
- 1-[2-[5-(4-Methylaminomethyl-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
- 25 1-[2-[5-(4-Dimethylaminomethyl-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
- 1-(2-{5-[2-(2-Methyl-imidazol-1-yl)-ethoxy]-benzoimidazol-1-yl}-quinolin-8-yl)-piperidin-4-ylamine and
- 1-[2-[5-(2-[1,2,4]Triazol-1-yl-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine,
- 30

and the pharmaceutically acceptable salts, prodrugs and solvates of the foregoing compounds.

- The invention also relates to a pharmaceutical composition for the treatment of a hyperproliferative disorder in a mammal which comprises a therapeutically effective amount of a compound of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a
- 35 pharmaceutically acceptable carrier. In one embodiment, said pharmaceutical composition is for the treatment of cancer such as brain, lung, squamous cell, bladder, gastric, pancreatic, breast, head, neck, renal, kidney, ovarian, prostate, colorectal, oesophageal, testicular, gynecological or thyroid cancer. In another embodiment, said pharmaceutical composition is for the treatment of

5 a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (e.g., psoriasis), restenosis, or prostate (e.g., benign prostatic hypertrophy (BPH)).

The invention also relates to a pharmaceutical composition for the treatment of pancreatitis or kidney disease (including proliferative glomerulonephritis and diabetes-induced renal disease) in a mammal which comprises a therapeutically effective amount of a compound
10 of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a pharmaceutically acceptable carrier.

The invention also relates to a pharmaceutical composition for the prevention of blastocyte implantation in a mammal that comprises a therapeutically effective amount of a compound of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a
15 pharmaceutically acceptable carrier.

The invention also relates to a pharmaceutical composition for treating a disease related to vasculogenesis or angiogenesis in a mammal which comprises a therapeutically effective amount of a compound of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a pharmaceutically acceptable carrier. In one embodiment, said pharmaceutical
20 composition is for treating a disease selected from the group consisting of tumor angiogenesis, chronic inflammatory disease such as rheumatoid arthritis, atherosclerosis, skin diseases such as psoriasis, excema, and scleroderma, diabetes, diabetic retinopathy, retinopathy of prematurity, age-related macular degeneration, hemangioma, glioma, melanoma, Kaposi's sarcoma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer.

The invention also relates to a method of treating a hyperproliferative disorder in a mammal that comprises administering to said mammal a therapeutically effective amount of the compound of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof. In one embodiment, said method relates to the treatment of cancer such as brain, squamous cell, bladder, gastric, pancreatic, breast, head, neck, oesophageal, prostate, colorectal, lung, renal,
30 kidney, ovarian, testicular, gynecological or thyroid cancer. In another embodiment, said method relates to the treatment of a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (e.g., psoriasis), restenosis or prostate (e.g., benign prostatic hypertrophy (BPH)).

The invention also relates to a method for the treatment of a hyperproliferative disorder
35 in a mammal which comprises administering to said mammal a therapeutically effective amount of a compound of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, in combination with an anti-tumor agent selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones,
40 angiogenesis inhibitors, and anti-androgens.

5 The invention also relates to a method of treating pancreatitis or kidney disease in a mammal which comprises administering to said mammal a therapeutically effective amount of a compound of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof.

 The invention also relates to a method of preventing blastocyte implantation in a mammal which comprises administering to said mammal a therapeutically effective amount of a
10 compound of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof.

 The invention also relates to a method of treating diseases related to vasculogenesis or angiogenesis in a mammal which comprises administering to said mammal an effective amount of a compound of formula 1, or a pharmaceutically acceptable salt, prodrug or hydrate thereof. In one embodiment, said method is for treating a disease selected from the group consisting of
15 tumor angiogenesis, chronic inflammatory disease such as rheumatoid arthritis, atherosclerosis, skin diseases such as psoriasis, excema, and scleroderma, diabetes, diabetic retinopathy, retinopathy of prematurity, age-related macular degeneration, hemangioma, glioma, melanoma, Kaposi's sarcoma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer.

 Patients that can be treated with a compounds of formula 1, and the pharmaceutically
20 acceptable salts, prodrugs and hydrates of said compounds, according to the methods of this invention include, for example, patients that have been diagnosed as having psoriasis, restenosis, atherosclerosis, BPH, lung cancer, bone cancer, CMML, pancreatic cancer, skin cancer, cancer of the head and neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, colon cancer, breast
25 cancer, testicular, gynecologic tumors (e.g., uterine sarcomas, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina or carcinoma of the vulva), Hodgkin's disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system (e.g., cancer of the thyroid, parathyroid or adrenal glands), sarcomas of soft tissues, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute
30 leukemia, solid tumors of childhood, lymphocytic lymphomas, cancer of the bladder, cancer of the kidney or ureter (e.g., renal cell carcinoma, carcinoma of the renal pelvis), or neoplasms of the central nervous system (e.g., primary CNS lymphoma, spinal axis tumors, brain stem gliomas or pituitary adenomas).

 This invention also relates to a pharmaceutical composition for inhibiting abnormal
35 cell growth in a mammal which comprises an amount of a compound of formula 1, or a pharmaceutically acceptable salt or solvate or prodrug thereof, in combination with an amount of a chemotherapeutic, wherein the amounts of the compound, salt, solvate, or prodrug, and of the chemotherapeutic are together effective in inhibiting abnormal cell growth. Many chemotherapeutics are presently known in the art. In one embodiment, the chemotherapeutic
40 is selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites,

- 5 intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, e.g. anti-androgens.

 This invention further relates to a method for inhibiting abnormal cell growth in a mammal or treating a hyperproliferative disorder which method comprises administering to the mammal an amount of a compound of formula 1, or a pharmaceutically acceptable salt or
10 solvate or prodrug thereof, in combination with radiation therapy, wherein the amount of the compound, salt, solvate or prodrug is in combination with the radiation therapy effective in inhibiting abnormal cell growth or treating the hyperproliferative disorder in the mammal. Techniques for administering radiation therapy are known in the art, and these techniques can be used in the combination therapy described herein. The administration of the compound of
15 the invention in this combination therapy can be determined as described herein.

 It is believed that the compounds of formula 1 can render abnormal cells more sensitive to treatment with radiation for purposes of killing and/or inhibiting the growth of such cells. Accordingly, this invention further relates to a method for sensitizing abnormal cells in a mammal to treatment with radiation which comprises administering to the mammal an amount
20 of a compound of formula 1 or pharmaceutically acceptable salt, prodrug or solvate thereof, which amount is effective in sensitizing abnormal cells to treatment with radiation. The amount of the compound, salt, or solvate in this method can be determined according to the means for ascertaining effective amounts of such compounds described herein.

 This invention also relates to a method of and to a pharmaceutical composition for
25 inhibiting abnormal cell growth in a mammal which comprises an amount of a compound of formula 1, a pharmaceutically acceptable salt or solvate thereof, a prodrug thereof, or an isotopically-labelled derivative thereof, and an amount of one or more substances selected from anti-angiogenesis agents, signal transduction inhibitors, and antiproliferative agents.

 Anti-angiogenesis agents, such as MMP-2 (matrix-metalloproteinase 2) inhibitors,
30 MMP-9 (matrix-metalloproteinase 9) inhibitors, and COX-II (cyclooxygenase II) inhibitors, can be used in conjunction with a compound of formula 1 and pharmaceutical compositions described herein. Examples of useful COX-II inhibitors include CELEBREX™ (alecoxib), valdecoxib, and rofecoxib. Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 (published October 24, 1996), WO 96/27583 (published March 7,
35 1996), European Patent Application No. 97304971.1 (filed July 8, 1997), European Patent Application No. 99308617.2 (filed October 29, 1999), WO 98/07697 (published February 26, 1998), WO 98/03516 (published January 29, 1998), WO 98/34918 (published August 13, 1998), WO 98/34915 (published August 13, 1998), WO 98/33768 (published August 6, 1998), WO 98/30566 (published July 16, 1998), European Patent Publication 606,046 (published July 13,
40 1994), European Patent Publication 931,788 (published July 28, 1999), WO 90/05719 (published

- 5 May 331, 1990), WO 99/52910 (published October 21, 1999), WO 99/52889 (published October 21, 1999), WO 99/29667 (published June 17, 1999), PCT International Application No. PCT/IB98/01113 (filed July 21, 1998), European Patent Application No. 99302232.1 (filed March 25, 1999), Great Britain patent application number 9912961.1 (filed June 3, 1999), United States Provisional Application No. 60/148,464 (filed August 12, 1999), United States Patent 5,863,949
 10 (issued January 26, 1999), United States Patent 5,861,510 (issued January 19, 1999), and European Patent Publication 780,386 (published June 25, 1997), all of which are incorporated herein in their entireties by reference. Preferred MMP-2 and MMP-9 inhibitors are those that have little or no activity inhibiting MMP-1. More preferred, are those that selectively inhibit MMP-2 and/or MMP-9 relative to the other matrix-metalloproteinases (*i.e.* MMP-1, MMP-3, MMP-4,
 15 MMP-5, MMP-6, MMP-7, MMP-8, MMP-10, MMP-11, MMP-12, and MMP-13).

Some specific examples of MMP inhibitors useful in the present invention are AG-3340, RO 32-3555, RS 13-0830, and the compounds recited in the following list:

- 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclopentyl)-amino]-propionic acid;
 20 3-exo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide;
 (2R, 3R) 1-[4-(2-chloro-4-fluoro-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide;
 4-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxylic acid
 25 hydroxyamide;
 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclobutyl)-amino]-propionic acid;
 4-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide;
 30 (R) 3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-3-carboxylic acid hydroxyamide;
 (2R, 3R) 1-[4-(4-fluoro-2-methyl-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide;
 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-1-methyl-ethyl)-
 35 amino]-propionic acid;
 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(4-hydroxycarbamoyl-tetrahydro-pyran-4-yl)-amino]-propionic acid;
 3-exo-3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide;

5 3-endo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; and
 (R) 3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-furan-3-carboxylic acid hydroxyamide;

 and pharmaceutically acceptable salts and solvates of said compounds.

10 Other anti-angiogenesis agents, including other COX-II inhibitors and other MMP inhibitors, can also be used in the present invention.

 A compound of formula 1, can also be used with signal transduction inhibitors, such as agents that can inhibit EGFR (epidermal growth factor receptor) responses, such as EGFR antibodies, EGF antibodies, and molecules that are EGFR inhibitors; VEGF (vascular
15 endothelial growth factor) inhibitors, such as VEGF receptors and molecules that can inhibit VEGF; and erbB2 receptor inhibitors, such as organic molecules or antibodies that bind to the erbB2 receptor, for example, HERCEPTIN™ (Genentech, Inc. of South San Francisco, California, USA).

 EGFR inhibitors are described in, for example in WO 95/19970 (published July 27,
20 1995), WO 98/14451 (published April 9, 1998), WO 98/02434 (published January 22, 1998), and United States Patent 5,747,498 (issued May 5, 1998), and such substances can be used in the present invention as described herein. EGFR-inhibiting agents include, but are not limited to, the monoclonal antibodies C225, anti-EGFR 22Mab (ImClone Systems Incorporated of New York, New York, USA), and **ABX-EGF (Abgenix antibody)** the compounds ZD-1839 (AstraZeneca),
25 BIBX-1382 (Boehringer Ingelheim), MDX-447 (Medarex Inc. of Annandale, New Jersey, USA), and OLX-103 (Merck & Co. of Whitehouse Station, New Jersey, USA), VRCTC-310 (Ventech Research) and EGF fusion toxin (Seragen Inc. of Hopkinton, Massachusetts). These and other EGFR-inhibiting agents can be used in the present invention.

 VEGF inhibitors, for example SU-5416 and SU-6668 (Sugen Inc. of South San
30 Francisco, California, USA), can also be combined with the compound of the present invention. VEGF inhibitors are described in, for example in WO 99/24440 (published May 20, 1999), PCT International Application PCT/IB99/00797 (filed May 3, 1999), in WO 95/21613 (published August 17, 1995), WO 99/61422 (published December 2, 1999), United States Patent 5,834,504 (issued November 10, 1998), WO 98/50356 (published November 12, 1998), United
35 States Patent 5,883,113 (issued March 16, 1999), United States Patent 5,886,020 (issued March 23, 1999), United States Patent 5,792,783 (issued August 11, 1998), WO 99/10349 (published March 4, 1999), WO 97/32856 (published September 12, 1997), WO 97/22596 (published June 26, 1997), WO 98/54093 (published December 3, 1998), WO 98/02438 (published January 22, 1998), WO 99/16755 (published April 8, 1999), and WO 98/02437 (published January 22, 1998),
40 all of which are incorporated herein in their entireties by reference. Other examples of some

5 specific VEGF inhibitors useful in the present invention are IM862 (Cytran Inc. of Kirkland, Washington, USA); IMC-1C11 Imclone antibody, anti-VEGF monoclonal antibody of Genentech, Inc. of South San Francisco, California; and angiozyme, a synthetic ribozyme from Ribozyme (Boulder, Colorado) and Chiron (Emeryville, California). These and other VEGF inhibitors can be used in the present invention as described herein.

10 ErbB2 receptor inhibitors, such as GW-282974 (Glaxo Wellcome plc), and the monoclonal antibodies AR-209 (Aronex Pharmaceuticals Inc. of The Woodlands, Texas, USA) and 2B-1 (Chiron), can furthermore be combined with the compound of the invention, for example those indicated in WO 98/02434 (published January 22, 1998), WO 99/35146 (published July 15, 1999), WO 99/35132 (published July 15, 1999), WO 98/02437 (published
15 January 22, 1998), WO 97/13760 (published April 17, 1997), WO 95/19970 (published July 27, 1995), United States Patent 5,587,458 (issued December 24, 1996), and United States Patent 5,877,305 (issued March 2, 1999), which are all hereby incorporated herein in their entireties by reference. ErbB2 receptor inhibitors useful in the present invention are also described in United States Provisional Application No. 60/117,341, filed January 27, 1999, and in United States
20 Provisional Application No. 60/117,346, filed January 27, 1999, both of which are incorporated in their entireties herein by reference. The erbB2 receptor inhibitor compounds and substance described in the aforementioned PCT applications, U.S. patents, and U.S. provisional applications, as well as other compounds and substances that inhibit the erbB2 receptor, can be used with the compound of the present invention in accordance with the present invention.

25 The compound of the invention can also be used with other agents useful in treating abnormal cell growth or cancer, including, but not limited to, agents capable of enhancing antitumor immune responses, such as CTLA4 (cytotoxic lymphocyte antigen 4) antibodies, and other agents capable of blocking CTLA4; and anti-proliferative agents such as farnesyl protein transferase inhibitors, and $\alpha\text{v}\beta 3$ inhibitors, such as the $\alpha\text{v}\beta 3$ antibody Vitaxin, and $\alpha\text{v}\beta 5$
30 inhibitors and the like. Specific CTLA4 antibodies that can be used in the present invention include those described in United States Provisional Application 60/113,647 (filed December 23, 1998) which is incorporated by reference in its entirety, however other CTLA4 antibodies can be used in the present invention.

The compounds of formula 1 and their pharmaceutically acceptable salts, prodrugs
35 and solvates can each independently also furthermore be used in a palliative neo-adjuvant/adjuvant therapy in alleviating the symptoms associated with the diseases recited herein as well as the symptoms associated with abnormal cell growth. Such therapy can be a monotherapy or can be in a combination with chemotherapy and/or immunotherapy.

The invention also relates to a method of preparing a compound of the formula 1

5 The terms "abnormal cell growth" and "hyperproliferative disorder" are used interchangeably in this application.

 "Abnormal cell growth", as used herein, unless otherwise indicated, refers to cell growth that is independent of normal regulatory mechanisms (e.g., loss of contact inhibition). This includes the abnormal growth of: (1) tumor cells (tumors) that proliferate by expressing a mutated tyrosine kinase or overexpression of a receptor tyrosine kinase; (2) benign and malignant cells of other proliferative diseases in which aberrant tyrosine kinase activation occurs; (4) any tumors that proliferate by receptor tyrosine kinases; (5) any tumors that proliferate by aberrant serine/threonine kinase activation; and (6) benign and malignant cells of other proliferative diseases in which aberrant serine/threonine kinase activation occurs..

15 The term "treating", as used herein, unless otherwise indicated, means reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment", as used herein, unless otherwise indicated, refers to the act of treating as "treating" is defined immediately above.

20 The term "Me" means methyl, "Et" means ethyl, and "Ac" means acetyl.

 The term "halo", as used herein, unless otherwise indicated, means fluoro, chloro, bromo or iodo. Preferred halo groups are fluoro, chloro and bromo.

 The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, branched, or cyclic moieties (including fused and bridged bicyclic and spirocyclic moieties), or a combination of the foregoing moieties. For an alkyl group to have cyclic moieties, the group must have at least three carbon atoms.

 The term "cycloalkyl", as used herein, unless otherwise indicated, includes cyclic alkyl moieties wherein alkyl is as defined above. The use of the term "cycloalkyl" shall not be construed as limiting the term "alkyl" to non-cyclic moieties.

30 The term "alkenyl", as used herein, unless otherwise indicated, includes alkyl moieties having at least one carbon-carbon double bond wherein alkyl is as defined above and including E and Z isomers of said alkenyl moiety.

 The term "alkynyl", as used herein, unless otherwise indicated, includes alkyl moieties having at least one carbon-carbon triple bond wherein alkyl is as defined above.

35 The term "alkoxy", as used herein, unless otherwise indicated, includes O-alkyl groups wherein alkyl is as defined above.

 The term "aryl", as used herein, unless otherwise indicated, includes an organic radical derived from an aromatic hydrocarbon by removal of one hydrogen, such as phenyl or naphthyl.

 The term "4 to 10 membered heterocyclic", as used herein, unless otherwise indicated, includes aromatic and non-aromatic heterocyclic groups containing one to four heteroatoms

5 each selected from O, S and N, wherein each heterocyclic group has from 4-10 atoms in its ring system, and with the proviso that the ring of said group does not contain two adjacent O or S atoms. Non-aromatic heterocyclic groups include groups having only 4 atoms in their ring system, but aromatic heterocyclic groups must have at least 5 atoms in their ring system. The heterocyclic groups include benzo-fused ring systems. An example of a 4 membered
10 heterocyclic group is azetidiny (derived from azetidine). An example of a 5 membered heterocyclic group is thiazolyl and an example of a 10 membered heterocyclic group is quinolinyl. Examples of non-aromatic heterocyclic groups are pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidino, morpholino, thiomorpholino, thioxanyl, piperazinyl, homopiperazinyl, azetidiny, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepiny, diazepiny, thiazepiny,
15 1,2,3,6-tetrahydropyridiny, 2-pyrroliny, 3-pyrroliny, indoliny, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazoliny, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidinylimidazoliny, imidazolidiny, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, azabicyclo[2.2.2]hexanyl, 3H-indolyl and quinoliziny. Examples of
20 aromatic heterocyclic groups are pyridiny, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyraziny, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnoliny, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, puriny, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl,
25 quinoxaliny, naphthyridiny, and furopyridiny. Spiro moieties are also included within the scope of this definition including 1-oxa-6-aza-spiro[2.5]oct-6-yl. The foregoing groups, as derived from the groups listed above, may be C-attached or N-attached where such is possible. For instance, a group derived from pyrrole may be pyrrol-1-yl (N-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole may be imidazol-1-yl (N-attached) or
30 imidazol-3-yl (C-attached). An example of a heterocyclic group wherein 2 ring carbon atoms are substituted with oxo (=O) moieties is 1,1-dioxo-thiomorpholinyl.

The phrase "pharmaceutically acceptable salt(s)", as used herein, unless otherwise indicated, includes salts of acidic or basic groups which may be present in the compounds of formula 1. The compounds of formula 1 that are basic in nature are capable of forming a wide
35 variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds of formula 1 are those that form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride,
40 edetate, edisylate, estolate, esylate, ethylsuccinate, fumarate, gluceptate, gluconate, glutamate,

5 glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, phosphatidylphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodode, and valerate salts. Since a single compound of the
10 present invention may include more than one acidic or basic moieties, the compounds of the present invention may include mono, di or tri-salts in a single compound.

Those compounds of the present invention that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline earth metal salts and, particularly, the calcium, magnesium,
15 sodium and potassium salts of the compounds of the present invention.

In the compounds of formula 1, where terms such as $(CR^4R^5)_m$ or $(CR^4R^5)_t$ are used, R^4 and R^5 may vary with each iteration of m or t above 1. For instance, where m or t is 2, the terms $(CR^4R^5)_m$ or $(CR^4R^5)_t$ may equal $-CH_2CH_2-$, or $-CH(CH_3)C(CH_2CH_3)(CH_2CH_2CH_3)-$, or any number of similar moieties falling within the scope of the definitions of R^4 and R^5 .

20 Certain compounds of formula 1 may have asymmetric centers and therefore exist in different enantiomeric forms. All optical isomers and stereoisomers of the compounds of formula 1, and mixtures thereof, are considered to be within the scope of the invention. With respect to the compounds of formula 1, the invention includes the use of a racemate, one or more enantiomeric forms, one or more diastereomeric forms, or mixtures thereof. The compounds of
25 formula 1 may also exist as tautomers. This invention relates to the use of all such tautomers and mixtures thereof.


The subject invention also includes isotopically-labelled compounds, which are identical to those recited in formula 1, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass
30 number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as 2H , 3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Compounds of the present invention, prodrugs thereof, and pharmaceutically acceptable salts of said compounds or of said prodrugs which contain the
35 aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labelled compounds of the present invention, for example those into which radioactive isotopes such as 3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., 3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with
40 heavier isotopes such as deuterium, i.e., 2H , can afford certain therapeutic advantages

5 resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labelled compounds of formula 1 of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples and Preparations below, by substituting a readily available isotopically labelled reagent for a
10 non-isotopically labelled reagent.

This invention also encompasses pharmaceutical compositions containing and methods of treating proliferative disorders, or abnormal cell growth, by administering prodrugs of compounds of the formula 1. Compounds of formula 1 having free amino, amido, hydroxy or carboxylic groups can be converted into prodrugs. Prodrugs include compounds wherein an
15 amino acid residue, or a polypeptide chain of two or more (e.g., two, three or four) amino acid residues is covalently joined through an amide or ester bond to a free amino, hydroxy or carboxylic acid group of compounds of formula 1. The amino acid residues include but are not limited to the 20 naturally occurring amino acids commonly designated by three letter symbols and also includes 4-hydroxyproline, hydroxylysine, demosine, isodemosine, 3-methylhistidine, norvalin, beta-alanine, gamma-aminobutyric acid, citrulline homocysteine, homoserine, ornithine
20 and methionine sulfone. Additional types of prodrugs are also encompassed. For instance, free carboxyl groups can be derivatized as amides or alkyl esters. Free hydroxy groups may be derivatized using groups including but not limited to hemisuccinates, phosphate esters, dimethylaminoacetates, and phosphoryloxymethyloxycarbonyls, as outlined in *Advanced Drug*
25 *Delivery Reviews*, 1996, 19, 115. Carbamate prodrugs of hydroxy and amino groups are also included, as are carbonate prodrugs, sulfonate esters and sulfate esters of hydroxy groups. Derivatization of hydroxy groups as (acyloxy)methyl and (acyloxy)ethyl ethers wherein the acyl group may be an alkyl ester, optionally substituted with groups including but not limited to ether, amine and carboxylic acid functionalities, or where the acyl group is an amino acid ester as
30 described above, are also encompassed. Prodrugs of this type are described in *J. Med. Chem.* 1996, 39, 10. Free amines can also be derivatized as amides, sulfonamides or phosphonamides. All of these prodrug moieties may incorporate groups including but not limited to ether, amine and carboxylic acid functionalities.


5

SCHEME 1




5

SCHEME 1A




SCHEME 1B




5

SCHEME 2



SCHEME 3


10



15


5

SCHEME 4




10

SCHEME 5




5


SCHEME 6



SCHEME 7



5

SCHEME 8Detailed Description Of The Invention

General synthetic methods which may be referred to for preparing the compounds of the present invention are provided in United States patent 5,990,146 (issued November 23, 1999)(Warner-Lambert Co.) and PCT published application number WO 99/16755 (published April 8, 1999)(Merck & Co.). The foregoing patent and patent application are incorporated herein by reference in their entirety.

Scheme 1 illustrates the synthesis of compounds of formula 1. In step 1, the diol of formula 2 is reacted with a trialkylsilyl chloride or trialkylsilyl trifluoromethanesulphonate (such as tert-butyldimethylsilyl) with a suitable organic base, such as imidazole or pyridine, in an organic solvent, such as dichloromethane (DCM) at temperature ranging from -78°C to 45°C , preferably at ambient temperature, for 1 to 12 hours to give compound of formula 3. In step 2, compound of formula 3 is reacted with a triflating reagent and base, such as N-phenylbis(trifluoromethanesulfonimide) and sodium hydride or trifluoromethanesulfonic anhydride and 2,6-dimethylpyridine, in an anhydrous organic solvent such as tetrahydrofuran (THF) or

- 5 DCM at a temperature ranging from -78°C to ambient temperature, preferably at ambient temperature, for 1 to 12 hours to give compound of formula 4.

In step 3, compound of formula 4 is reacted with the amine of formula 5, preferably $X = C$, with a palladium catalyst such as, tris(dibenzylideneacetone)dipalladium (0) or palladium(II) acetate and a base such as, cesium carbonate or sodium tert-butoxide, preferably cesium carbonate, and a palladium ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) or 1,2-bis(diphenylphosphino)ethane (DIPHOS), in a solvent such as 1,4-dioxane or toluene, preferably toluene, at a temperature ranging from ambient temperature to between $80-105^{\circ}\text{C}$, preferably at 105°C for 1 to 48 hours. In step 4, the resulting compound of formula 6 is reduced under palladium catalysis, using either 10% palladium on carbon or 20% palladium hydroxide on carbon, with a hydrogen source, such as hydrazine, ammonium formate, or formic acid in an organic solvent such as ethanol or methanol with or without a co-solvent solvent such as THF, at a temperature ranging from ambient temperature to reflux for 1 to 24 hours to give compound of the formula 7.

In step 5, compound of the formula 7 is reacted with formamidine acetate or formic acid in an organic solvent such as 2-methoxyethanol, 1-butanol, ethanol, or formic acid, preferably ethanol, at a temperature ranging from ambient temperature to reflux, preferably at reflux, for 1 to 48 hours to give compound of formula 8. In step 6, compound of formula 8 is reacted with a triflating reagent and base, such as N-phenyl-bis(trifluoromethanesulfonimide) and sodium hydride or triethylamine in an anhydrous organic solvent such as THF, with or without a co-solvent such as dimethylformamide (DMF), at a temperature ranging from -78°C to ambient temperature, preferably at ambient temperature, for 1 to 24 hours to give compound of formula 9.

In step 7, for R^1 groups that include an aryl or heteroaryl group (where Ar = aryl or heteroaryl), compound of formula 9 is reacted under palladium catalysis, such as with tetrakis(triphenylphosphine)palladium (0), with the appropriate organoborane (where $Z = B(OH)_2$ or $B(Alkyl)_2$), organostannane (where $Z = Sn(alkyl)_3$) or organozinc (where $Z = Zn(Halogen)$). When $Z = B(OH)_2$, a base such as potassium phosphate is used in a solvent such as 1,4-dioxane or 1,2-dimethoxyethane at a temperature ranging from ambient temperature to reflux, preferably at reflux, for 1 to 48 hours to give compound of the formula 1.

35 When $Z = B(alkyl)_2$, a base such as sodium carbonate with or without lithium chloride is used in a solvent system including ethanol and water with or without toluene at a temperature ranging from ambient temperature to reflux, preferably around 90°C , for 1 to 48 hours to give compound of the formula 1. When $Z = Sn(alkyl)_3$, with or without a base such as potassium phosphate, in a suitable organic solvent such as toluene or 1,4-dioxane and at a temperature

5 ranging from ambient temperature to reflux, preferably between 80-100°C, for 1 to 48 hours to give compound of the formula 1. When Z = Zn(Halogen), a suitable organic solvent such as THF, 1,4-dioxane or 1,2-dimethoxyethane is used at a temperature ranging from -78°C to reflux, preferably between 20-45°C, for 1 to 48 hours to give compound of the formula 1. For R¹ with an NR⁵R⁶ moiety, compound of the formula 9 is reacted with an amine HNR⁵R⁶ using a
10 palladium catalyst such as tris(dibenzylideneacetone)dipalladium (0) or palladium acetate and a base such as, cesium carbonate or sodium tert-butoxide, preferably cesium carbonate, and a palladium ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) or 1,2-bis(diphenylphosphino)ethane (DIPHOS) in a solvent such as 1,4-dioxane, toluene, and xylenes preferably toluene, at a temperature ranging from ambient temperature to reflux,
15 preferably at reflux for 1 to 48 hours to give compound of formula 1. Compounds of formula 1 may have protecting groups, such as R¹¹, R¹⁰ or R⁹ = OMe, that can be removed by standard conditions as discussed in "Protective Groups for Organic Synthesis". For example, R¹¹, R¹⁰ or R⁹ = OMe can be transformed to R¹¹, R¹⁰ or R⁹ = OH by treating with borontribromide in an organic solvent such as DCM at a temperature ranging from -78°C to 45°C, preferably at
20 ambient temperature for 1 to 24 hours.

Scheme 1A illustrates an alternative synthesis of compounds of formula 7. In step 1, the diol of formula 2 is reacted with benzyl chloride or benzyl bromide, preferably benzyl bromide, with a suitable base, such as potassium carbonate, sodium carbonate or cesium carbonate, in an organic solvent, such as DMF at temperature ranging from -78°C to 100°C,
25 preferably between 60-80°C for 3 to 24 hours to give compound of formula 3A. In step 2, compound of formula 3A is reacted with a triflating reagent and base, such as N-phenyl-bis(trifluoromethanesulfonimide) and sodium hydride or trifluoromethanesulfonic anhydride and 2,6-dimethylpyridine, in an anhydrous organic solvent such as tetrahydrofuran (THF) or DCM at a temperature ranging from -78°C to ambient temperature, preferably at ambient
30 temperature, for 1 to 12 hours to give compound of formula 4A (where V = OTf). Alternatively, compound of the formula 3A is reacted with a chlorinating reagent, such as phosphorus oxychloride, thionyl chloride or oxalyl chloride, in an organic solvent such as DCM, 1,2-dichloroethane (DCE) or chloroform at a temperature ranging from ambient temperature to reflux, preferably at reflux, to give compound of the formula 4A (where V = Cl). Alternatively,
35 compound of the formula 3A is reacted with a brominating reagent, such as phosphorus oxybromide, in an organic solvent such as DCM, 1,2-dichloroethane (DCE) or chloroform at a temperature ranging from ambient temperature to reflux, preferably at reflux, to give compound of the formula 4A (where V = Br).

5 In step 3, compound of formula **4A** is reacted with the amine of formula **5**, preferably
X = C, with a palladium catalyst such as, tris(dibenzylideneacetone)dipalladium (0) or
palladium(II) acetate and a base such as, cesium carbonate or sodium tert-butoxide,
preferably cesium carbonate, and a palladium ligand such as 2,2'-bis(diphenylphosphino)-1,1'-
binaphthyl (BINAP) or 1,2-bis(diphenylphosphino)ethane (DIPHOS), in a solvent such as 1,4-
10 dioxane or toluene, preferably toluene, at a temperature ranging from ambient temperature to
105°C, preferably between 80-105°C for 1 to 48 hours. In step 4, the resulting compound of
formula **6A** is reduced under palladium catalysis, using either 10% palladium on carbon or
20% palladium hydroxide on carbon, with a hydrogen source, such as ammonium formate,
triethylammonium formate or formic acid, preferably ammonium formate triethylammonium
15 formate, in an organic solvent such as ethanol (EtOH) or methanol (MeOH) with or without a
co-solvent solvent such as THF, at a temperature ranging from ambient temperature to reflux,
preferably between 75°C and at reflux, for 1 to 24 hours to give compound of the formula **7**.

Scheme 1B illustrates an alternative synthesis of compounds of formula **6A**. In step
1, the aminoquinoline of formula **2B** is reacted with benzyl chloride or benzyl bromide,
20 preferably benzyl bromide, with a suitable base, such as sodium hydride or potassium
hydride, in an organic solvent, such as DMF, THF or 1,2-dimethoxyethane, at temperature
ranging from -78°C to 65°C, preferably between 0-25°C, for 1 to 24 hours to give compound of
formula **3B**. In step 2, compound of formula **3B** is reacted with bromoaromatic compound of
formula **5A** with a palladium catalyst such as, tris(dibenzylideneacetone)dipalladium (0) or
25 palladium(II) acetate, preferably tris(dibenzylideneacetone)dipalladium (0), and a base such
as, cesium carbonate or sodium tert-butoxide, preferably sodium tert-butoxide, and a
palladium ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) or 1,2-
bis(diphenylphosphino)ethane (DIPHOS), preferably 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
(BINAP), in a solvent such as 1,4-dioxane or toluene, at a temperature ranging from ambient
30 temperature to 105°C, preferably between 80-105°C for 1 to 48 hours to give compound of the
formula **6A**.

Scheme 2 illustrates the synthesis of compounds of formula **11**. In step 1, compound
of the formula **10** is reacted under standard alkylation conditions by treating with an
electrophile R³Y, where Y can be mesylate, tosylate, bromo, iodo and chloro, preferably
35 bromo or iodo, and a base such as sodium hydride, potassium hydride, sodium carbonate,
potassium carbonate or cesium carbonate, preferably cesium carbonate in an organic solvent
such as DMF or THF, preferably DMF, for 1 to 48 hours at a temperature ranging from -78°C
to 85°C. Compounds of the type formula **11** can also be obtained by starting with the
appropriate amine compound of the formula **5** in step 3 of scheme 1 or step 2 of scheme 1A.

- 5 Alternatively, compounds of the type formula 11 can also be obtained by starting with the appropriate bromoaromatic compound of the formula 5A in step 2 of scheme 1B.

Scheme 2 also illustrates the synthesis of compounds of formula 13 (wherein Ar is an aryl or heteroaryl group). In step 2, compound of the formula 10 is reacted with a triflating reagent and base, such as N-phenyl-bis(trifluoromethanesulfonimide) and sodium hydride or
10 trifluoromethanesulfonic anhydride and pyridine, in an anhydrous organic solvent such as THF, with or without a co-solvent such as DMF, at a temperature ranging from -78°C to ambient temperature, preferably at ambient temperature, for 1 to 24 hours to give compound of formula 12. In step 2, compound of formula 12 is reacted under palladium catalysis, such as with tetrakis(triphenylphosphine)palladium (0), with the appropriate organoborane (where Z
15 = $\text{B}(\text{OH})_2$ or $\text{B}(\text{Alkyl})_2$), organostannane (where Z = $\text{Sn}(\text{alkyl})_3$) or organozinc (where Z = $\text{Zn}(\text{Halogen})$). When Z = $\text{B}(\text{OH})_2$, a base such as potassium phosphate is used in a solvent such as 1,4-dioxane or 1,2-dimethoxyethane at a temperature ranging from ambient temperature to reflux, preferably at reflux, for 1 to 48 hours to give compound of the formula 13. When Z = $\text{B}(\text{alkyl})_2$, a base such as sodium carbonate with or without lithium chloride is
20 used in a solvent system including ethanol and water with or without toluene at a temperature ranging from ambient temperature to reflux, preferably around 90°C , for 1 to 48 hours to give compound of the formula 13. When Z = $\text{Sn}(\text{alkyl})_3$, with or without a base such as potassium phosphate, in a suitable organic solvent such as toluene or 1,4-dioxane and at a temperature ranging from ambient temperature to reflux, preferably between 80 - 100°C , for 1 to 48 hours to
25 give compound of the formula 13. When Z = $\text{Zn}(\text{Halogen})$, a suitable organic solvent such as THF, 1,4-dioxane or 1,2-dimethoxyethane is used at a temperature ranging from -78°C to reflux, preferably between 20 - 45°C , for 1 to 48 hours to give compound of the formula 13. Compounds of formula 13 may have protecting groups, such as R^{11} , R^{10} or $\text{R}^9 = \text{OMe}$, that can be removed by standard conditions as discussed in "Protective Groups for Organic
30 Synthesis". For example, R^{11} , R^{10} or $\text{R}^9 = \text{OMe}$ can be transformed to R^{11} , R^{10} or $\text{R}^9 = \text{OH}$ by treating with borontribromide in an organic solvent such as DCM at a temperature ranging from -78°C to 45°C , preferably at ambient temperature for 1 to 24 hours. Compounds of the type formula 13 can also be obtained by starting with the appropriate amine compound of the formula 5 in step 3 of scheme 1 or step 2 of scheme 1A. Alternatively, compounds of the type
35 formula 11 can also be obtained by starting with the appropriate bromoaromatic compound of the formula 5A in step 2 of scheme 1B.

Scheme 3 illustrates the synthesis of compounds of formula 15 (where W = aryl, or heterocycle). In step 1, compound of formula 14 is reacted with an amine HNR^3R^4 and reducing agent, such as sodium cyanoborohydride or sodium triacetoxyborohydride, and

- 5 acetic acid in an organic solvent, such as methanol or ethanol, and with or without a co-solvent such as 1,2-dichloroethane, at a temperature ranging from 0°C to 80°C, preferably at ambient temperature, for 1 to 24 hours. Compounds of formula 15 may have protecting groups, such as R¹¹, R¹⁰ or R⁹ = OMe, that can be removed by standard conditions as discussed in "Protective Groups for Organic Synthesis". For example, R¹¹, R¹⁰ or R⁹ = OMe
- 10 can be transformed to R¹¹, R¹⁰ or R⁹ = OH by treating with borontribromide in an organic solvent such as DCM at a temperature ranging from -78°C to 45°C, preferably at ambient temperature for 1 to 24 hours. Compounds of the type formula 15 can also be obtained by starting with the appropriate ArZ or amine NR⁵R⁶ in step 7 of scheme 1.

- Scheme 4 illustrates the synthesis of compounds of formula 18. In step 1, compound
- 15 of formula 16 is reacted with trimethylsulfonium iodide and a base, such as sodium hydride, in an organic solvent, such as dimethylsulfoxide (DMSO) or THF, at a temperature ranging from -78°C to 65°C, preferably at ambient temperature, for 1 to 24 hours. The resulting compound of formula 17 is reacted in step 2 with an amine NR³R⁴ in a solvent such as THF, methanol, ethanol, water, DMF, DMSO or any combination thereof at a temperature ranging from 0°C to
- 20 100°C, preferably at 65°C in a sealed tube, for 1 to 48 hours to give compound of formula 18. Compounds of formula 18 may have protecting groups, such as R¹¹, R¹⁰ or R⁹ = OMe, that can be removed by standard conditions as discussed in "Protective Groups for Organic Synthesis". For example, R¹¹, R¹⁰ or R⁹ = OMe can be transformed to R¹¹, R¹⁰ or R⁹ = OH by treating with borontribromide in an organic solvent such as DCM at a temperature ranging
- 25 from -78°C to 45°C, preferably at ambient temperature for 1 to 24 hours. Compounds of the type formula 18 can also be obtained by starting with the appropriate amine NR⁵R⁶ in step 7 of scheme 1.

- Scheme 4 also illustrates the synthesis of compound of formula 19. In step 1, compound of formula 16 is reacted with an amine HNR³R⁴ and reducing agent, such as
- 30 sodium cyanoborohydride or sodium triacetoxyborohydride, and acetic acid in an organic solvent, such as methanol, ethanol and with or without a co-solvent such as 1,2-dichloroethane (DCE), at a temperature ranging from 0°C to 80°C, preferably at ambient temperature, for 1 to 24 hours. Compounds of formula 19 may have protecting groups, such as R¹¹, R¹⁰ or R⁹ = OMe, that can be removed by standard conditions as discussed in
- 35 "Protective Groups for Organic Synthesis". For example, R¹¹, R¹⁰ or R⁹ = OMe can be transformed to R¹¹, R¹⁰ or R⁹ = OH by treating with borontribromide in an organic solvent such as DCM at a temperature ranging from -78°C to 45°C, preferably at ambient temperature for 1 to 24 hours. Compound of the type formula 19 can also be obtained by starting with the appropriate amine NR⁵R⁶ in step 7 of scheme 1.

5 Scheme 5 and 6 illustrate an alternative synthetic scheme for compound of formula 1. In step 1, bromoaniline **20** is acylated with cinnamyl chloride in an organic solvent, such as DCM or THF, preferably DCM, in the presence of an organic base, such as pyridine or triethylamine, preferably pyridine, at a temperature ranging from -78°C to 40°C, preferably between 0°C and 25°C, for 1 to 24 hours to give compound of formula **21**. In step 2,
10 compound of formula **21** is reacted with a strong Lewis acid, such as aluminum trichloride, in an organic solvent, such as chlorobenzene, at a temperature ranging from 25°C to 120°C, preferably between 90°C and 120°C, for 1 to 24 hours to give compound of formula **22**. In step 3, compound of formula **22** is reacted with a triflating reagent and base, such as N-phenyl-bis(trifluoromethanesulfonimide) and sodium hydride or trifluoromethanesulfonic
15 anhydride and 2,6-dimethylpyridine, in an anhydrous organic solvent such as THF or DCM at a temperature ranging from -78°C to ambient temperature, preferably at ambient temperature, for 1 to 24 hours to give compound of formula **23**.

 In scheme 6, compound of formula **23**, in step 1, is reacted with an amine of formula **5**, preferably X = C, with a palladium catalyst such as, tris(dibenzylideneacetone)dipalladium
20 (0) or palladium(II) acetate and a base such as, cesium carbonate or sodium tert-butoxide, preferably cesium carbonate, and a palladium ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) or 1,2-bis(diphenylphosphino)ethane (DIPHOS), in a solvent such as 1,2-dioxane or toluene, preferably toluene, at a temperature ranging from ambient temperature to 105°C, preferably between 80°C-105°C for 1 to 48 hours. In step 2, the resulting compound of
25 formula **24** is reduced with iron powder and ammonium chloride in an organic solvent, such as ethanol or methanol, with or without a co-solvent solvent, such as water, at a temperature ranging from ambient temperature to reflux, preferably at reflux, for 1 to 24 hours to give compound of the formula **25**.

 In step 3, compound of the formula **25** is reacted with formamidine acetate or formic
30 acid in an organic solvent such as 2-methoxyethanol, 1-butanol, ethanol, or formic acid, preferably ethanol, at a temperature ranging from ambient temperature to reflux, preferably at reflux, for 1 to 48 hours to give compound of formula **26**. In step 4, for R¹ groups that include an aryl or heteroaryl group (where Ar = aryl or heteroaryl), compound of formula **26** is reacted under palladium catalysis, such as with tetrakis(triphenylphosphine)palladium (0), with the
35 appropriate organoborane (where Z = B(OH)₂ or B(Alkyl)₂), organostannane (where Z = Sn(alkyl)₃) or organozinc (where Z = Zn(Halogen)). When Z = B(OH)₂, a base such as potassium phosphate is used in a solvent such as 1,4-dioxane or 1,2-dimethoxyethane at a temperature ranging from ambient temperature to reflux, preferably at reflux, for 1 to 48 hours to give compound of the formula 1. When Z = B(alkyl)₂, a base such as sodium carbonate with

5 or without lithium chloride is used in a solvent system including ethanol and water with or without toluene at a temperature ranging from ambient temperature to reflux, preferably around 90°C, for 1 to 48 hours to give compound of the formula 1. When Z = Sn(alkyl)₃, with or without a base such as potassium phosphate, in a suitable organic solvent such as toluene or 1,4-dioxane and at a temperature ranging from ambient temperature to reflux, preferably
10 between 80-100°C, for 1 to 48 hours to give compound of the formula 1. When Z = Zn(Halogen), a suitable organic solvent such as THF, 1,4-dioxane or 1,2-dimethoxyethane is used at a temperature ranging from -78°C to reflux, preferably between 20-45°C, for 1 to 48 hours to give compound of the formula 1. For R¹ with an NR⁵R⁶ moiety, compound of the formula 26 is reacted with an amine HNR⁵R⁶ using a palladium catalyst such as
15 tris(dibenzylideneacetone)dipalladium (0) or palladium acetate and a base such as, cesium carbonate or sodium tert-butoxide, preferably cesium carbonate, and a palladium ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) or 1,2-bis(diphenylphosphino)ethane (DIPHOS) in a solvent such as 1,4-dioxane, toluene, and xylenes, preferably toluene when Y = Br and preferably xylenes when Y = Cl, at a temperature ranging from ambient temperature
20 to reflux, preferably at reflux for 1 to 72 hours to give compound of formula 1. Compounds of formula 1 may have protecting groups, such as R¹¹, R¹⁰ or R⁹ = OMe, that can be removed by standard conditions as discussed in "Protective Groups for Organic Synthesis". For example, R¹¹, R¹⁰ or R⁹ = OMe can be transformed to R¹¹, R¹⁰ or R⁹ = OH by treating with borontribromide in an organic solvent such as DCM at a temperature ranging from -78°C to
25 45°C, preferably at ambient temperature for 1 to 24 hours.

Alternatively, compound of formula 26 can be prepared from compounds of formula 22 in a two step sequence as outlined in scheme 7. In step 1, compound of formula 22, where X = Cl, is reacted with either phosphorous oxychloride, thionyl chloride or oxalyl chloride, preferably oxalyl chloride, with or without an organic solvent, such as chloroform or DCE,
30 preferably DCE, at a temperature ranging from ambient temperature to reflux, preferably at reflux, for 1 to 24 hours to give compound of the formula 27. Where X = Br, compound of formula 22 is reacted with phosphorous oxybromide with an organic solvent, such as chloroform or DCE, preferably chloroform, at a temperature ranging from ambient temperature to reflux, preferably at reflux, for 1 to 24 hours to give compound of the formula 27. In step 2,
35 compound of formula 27, where X = Br or Cl, is reacted with a compound of formula 28 in an organic solvent, such as DMF or 1-methyl-2-pyrrolidinone, with or without a base, such as sodium hydride or sodium bis(trimethylsilyl)amide, at a temperature ranging from ambient temperature to 150°C, preferably at 60-85°C when using a base and 150°C when not, for 1 to 24 hours to give compound of the formula 26.

5 In step 1 of scheme 8, compound of the formula **8** is reacted with an electrophile RY, where Y is a mesylate, tosylate, bromide, chloride or iodide, and a base, such as sodium hydride, potassium hydride, potassium carbonate, sodium carbonate or cesium carbonate, in a solvent, such as DMF, THF, DMSO or 1,2-dimethoxyethane, at a temperature ranging from -78°C to 65°C, to give compound of the formula **29**. Compounds of formula **29** may have
10 protecting groups, such as R¹¹, R¹⁰ or R⁹ = OMe, that can be removed by standard conditions as discussed in "Protective Groups for Organic Synthesis". For example, R¹¹, R¹⁰ or R⁹ = OMe can be transformed to R¹¹, R¹⁰ or R⁹ = OH by treating with borontribromide in an organic solvent such as DCM at a temperature ranging from -78°C to 45°C, preferably at ambient temperature for 1 to 24 hours.

15

Also outlined in scheme 1 in step 1, compound of the formula **9** is reacted with carbon monoxide, at a pressure range of atmospheric to 50 psi, preferably 50 psi, in the presence of an organic base, such as triethylamine, under palladium catalysis, such as palladium acetate, with a ligand, such as 1,3-bis(diphenylphosphino)propane, in a solvent, such as DMF, in the
20 presence of methanol to give compound of the formula **30**. In step 2, compound of the formula **30** is reacted with a preformed complex of an amine HNR³R⁴ (or its hydrochloride salt) with trimethylaluminum in a solvent such as DCM or DCE at a temperature range of 0°C to reflux to give compound of the formula **31**. Compounds of formula **31** may have protecting groups, such as R¹¹, R¹⁰ or R⁹ = OMe, that can be removed by standard conditions as
25 discussed in "Protective Groups for Organic Synthesis". For example, R¹¹, R¹⁰ or R⁹ = OMe can be transformed to R¹¹, R¹⁰ or R⁹ = OH by treating with borontribromide in an organic solvent such as DCM at a temperature ranging from -78°C to 45°C, preferably at ambient temperature for 1 to 24 hours.

The compounds of the present invention may have asymmetric carbon atoms.
30 Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known to those skilled in the art, for example, by chromatography or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixtures into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., alcohol), separating the diastereomers and converting (e.g., hydrolyzing)
35 the individual diastereomers to the corresponding pure enantiomers. All such isomers, including diastereomeric mixtures and pure enantiomers are considered as part of the invention.

The compounds of formulas **1**, **13**, **15**, **18**, **19**, **29**, **31** that are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often

5 desirable in practice to initially isolate the compound of formula **1, 13, 15, 18, 19, 29, 31** from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a
10 substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained. The desired acid salt can also be precipitated from a solution of the free base in an organic solvent by adding to the solution an appropriate mineral or organic acid.

15 Those compounds of formula **1, 13, 15, 18, 19, 29, 31** that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this
20 invention are those which form non-toxic base salts with the acidic compounds of formula **1, 13, 15, 18, 19, 29, 31**. Such non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium calcium and magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting
25 solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction and maximum yields of the desired final product.
30 Since a single compound of the present invention may include more than one acidic or basic moieties, the compounds of the present invention may include mono, di or tri-salts in a single compound.

The activity of the compounds of formula **1, 13, 15, 18, 19, 29, 31** may be determined by the following procedure.

35 General PGT Kinase ELISA Method

The following reagent and stock solutions are used:

adenosine triphosphate (ATP)	Sigma, cat. # A-2383
bovine serum albumin (BSA)	Sigma, cat. # A-3294
Dulbecco's PBS (dPBS)	Gibco-BRL, cat. # 14190-136
40 MaxiSorp plates	Nunc, cat. # 439454

5	MgCl ₂	Sigma, cat. # M-1028
	Poly-Glu-Tyr (PGT)	Sigma, cat. #. P-0275
	TMB Micowell Substrate	Kirkegaard & Perry, cat. # 50-76-05
	Tween 20	Sigma, cat. # P-1379
	HRP-PY54 antibody	OSI Pharmaceuticals, Inc.

10 Phosphorylation Buffer (PB): 50 mM HEPES, pH 7.3, 125 mM NaCl, 24 mM MgCl₂;
 Wash Buffer (WB): dPBS + 0.1% Tween 20 (polyoxyethylene sorbitan); and
 Blocking Buffer: 3% BSA, 0.05% Tween 20 in dPBS.

Assay procedure:

(a) For plate coating, fill Nunc MaxiSorp plate with 100 μ l per well of Poly-Glu-Tyr (PGT) diluted in dPBS (various concentrations). The plate is incubated overnight at 37°C. The supernatant PGT is then discarded, and the plates are washed 3X with Wash Buffer.

(b) The PDGF enzyme is then diluted in PB to an appropriate concentration, and 25 μ l of this stock solution is added per well.

(c) ATP is then diluted (from 20 mM stock) to an appropriate concentration (0.5 nM- 2 uM) with PB. The phosphorylation reaction is commenced by addition of 25 μ l ATP solution to each well of the assay plate. Incubation is continued for about 10 minutes, with shaking at room temperature.

(d) The reaction is stopped by aspirating off the reaction mixture. The plate is then washed 4X with WB.

(e) The HRP-PY54 antibody is diluted to an appropriate concentration in blocking buffer. 50
25 µl of this solution is then added per well, followed by incubation for 25-35 minutes at room
temperature. The antibody-containing solution is aspirated away, and the plate is again washed 4X
with WB.

(f) The extent of reaction is determined by measurement of light absorbance at 450 nm. First, color is developed by addition of TMB solution, 50 μ l per well, and the reaction is permitted to run until wells with positive signals achieve about 0.6-1.2 OD₄₅₀ units. Color development is then stopped by addition of 50 μ l per well of 0.09 M H₂SO₄. The background controls are wells without PGT, but with all other components included. As aforementioned, preferred signal is generally in the range of 0.6-1.2 OD units, with essentially no background.

The *in vitro* activity of the compounds of the present invention in inhibiting the PDGF β receptor may be determined by the following procedure.

Inhibition of tyrosine kinase activity may be measured using a recombinant enzyme in an assay that measures the ability of compounds to inhibit the phosphorylation of the exogenous substrate, polyGluTyr (PGT, Sigma™, 4:1). The cytoplasmic domain of the human PDGFβ receptor (amino acids 559-1106) (Ishikawa, F., et al. Nature 338: 557-562, 1989) is expressed in Sf9 insect cells as a glutathione S-transferase (GST)-fusion protein using the

5 baculovirus expression system. The protein is then purified from the lysates of these cells using glutathione agarose affinity columns.

The enzyme assay is performed in 96-well plates that are coated with the PGT substrate (0.625 μ g PGT per well). Test compounds are diluted in dimethylsulfoxide (DMSO), and then added to the PGT plates so that the final concentration of DMSO in the assay is
10 1.6% (v/v). The recombinant enzyme is diluted in phosphorylation buffer (50 mM Hepes, pH 7.3, 125 mM NaCl, 24 mM $MgCl_2$). The reaction is initiated by the addition of ATP to a final concentration of 10 μ M. After a 10 minute incubation at room temperature with shaking, the reaction is aspirated, and the plates are washed with wash buffer (PBS-containing 0.1% Tween-20). The amount of phosphorylated PGT is quantitated by incubation with a
15 horseradish peroxidase(HRP)-conjugated PY-54 antibody (Transduction Labs), developing with TMB peroxidase (TMB is 3,3',5,5'-tetramethylbenzidine), and detection on a BioRad™ Microplate reader at 450 nM. Inhibition of the kinase enzymatic activity by the test compound is detected as a reduced absorbance, and the concentration of the compound that is required to inhibit the signal by 50% (under the circumstances of the assay) is reported as the IC_{50}
20 value for the test compound.

To measure the ability of the compounds to inhibit PDGFR β tyrosine kinase activity for the full length protein that exists in a cellular context, the porcine aortic endothelial (PAE) cells transfected with the human PDGFR β (Westermarck, Bengt, et. al., PNAS 87, pp128-132, 1990) may be used. Cells are plated and allowed to attach to 96-well dishes in the same
25 media (Ham's F12) with 10% FBS (fetal bovine serum) for 6-8 hours. The cells are washed, re-fed with serum depleted media, and allowed to incubate over night. Immediately prior to dosing with compound, the cells are re-fed with the serum depleted media. Test compounds, dissolved in DMSO, are diluted into the media (final DMSO concentration 0.5% (v/v)). At the end of a 10 minutes incubation, PDGF-BB (100 ng/ml final) is added to the media for an 8
30 minute incubation. The cells are washed with Hepes buffered saline solution (HBSS) and lysed in 50 μ l of HNTG buffer (20 mM Hepes, pH 7.5, 150 mM NaCl, 0.2% Triton™ X-100, 10% glycerol, plus 0.2 mM PMSF (phenylmethylsulfonyl fluoride), 1 μ g/ml pepstatin, 1 μ g/ml leupeptin, 1 μ g/ml aprotinin, 2 mM sodium pyrophosphate, 2 mM sodium orthovanadate) and then diluted with 50 μ l of HG dilution buffer (20 mM Hepes, pH 7.5, 10% glycerol, 0.2 mM
35 PMSF (phenylmethylsulfonyl fluoride), 1 μ g/ml pepstatin, 1 μ g/ml leupeptin, 1 μ g/ml aprotinin, 2 mM sodium pyrophosphate, 2 mM sodium orthovanadate). The extent of phosphorylation of PDGFR β is measured using an ELISA assay. The 96-well Protein A coated plates are blocked with Superblock (Pierce) and coated with 0.5 μ g per well anti- PDGFR β P20 antibody (Santa Cruz, catalog number SC-339).

5 Any unbound antibody is washed off the plates prior to addition of the cell lysate. After a 2 hour room temperature incubation of the lysates (50 ul) with the PDGFR β antibody, the PDGFR β associated phosphotyrosine is quantitated by development with the HRP-conjugated PY-54 antibody and TMB, as described above. The ability of the compounds to inhibit the PDGF-BB stimulated autophosphorylation reaction by 50% under the conditions
10 used, relative to PDGF-BB-stimulated controls, is reported as the IC₅₀ value for the test compound. The compounds of the present invention, including the examples recited below, generally have IC₅₀ values using the foregoing procedure falling within the following range: 1-1000 nM.

Assay for Inhibition Activity Toward the KDR/VEGF Receptor

15 The *in vitro* activity of the compounds of the present invention in inhibiting the KDR/VEGF receptor may be determined by the following procedure.

The ability of the compounds of the present invention to inhibit tyrosine kinase activity may be measured using a recombinant enzyme in an assay that measures the ability of compounds to inhibit the phosphorylation of the exogenous substrate, polyGluTyr (PGT, Sigma™, 4:1). The kinase domain of the human KDR/VEGF receptor (amino acids 805-1350)
20 is expressed in Sf9 insect cells as a glutathione S-transferase (GST)-fusion protein using the baculovirus expression system. The protein is purified from the lysates of these cells using glutathione agarose affinity columns. The enzyme assay is performed in 96-well plates that are coated with the PGT substrate (0.625 μ g PGT per well). Test compounds are diluted in dimethylsulfoxide (DMSO), and then added to the PGT plates so that the final concentration of
25 DMSO in the assay is 1.6% (v/v). The recombinant enzyme is diluted in phosphorylation buffer (50 mM Hepes, pH 7.3, 125 mM NaCl, 24 mM MgCl₂). The reaction is initiated by the addition of ATP to a final concentration of 10 μ M. After a 30 minute incubation at room temperature with shaking, the reaction is aspirated, and the plates are washed with wash
30 buffer (PBS-containing 0.1% Tween-20). The amount of phosphorylated PGT is quantitated by incubation with a HRP-conjugated (HRP is horseradish peroxidase) PY-54 antibody (Transduction Labs), developed with TMB peroxidase (TMB is 3,3',5,5'-tetramethylbenzidine), and the reaction is quantitated on a BioRad™ Microplate reader at 450 nM. Inhibition of the kinase enzymatic activity by the test compound is detected as a reduced absorbance, and the
35 concentration of the compound that is required to inhibit the signal by 50% is reported as the IC₅₀ value for the test compound.

To measure the ability of the compounds to inhibit KDR tyrosine kinase activity for the full length protein that exists in a cellular context, the porcine aortic endothelial (PAE) cells transfected with the human KDR (Waltenberger et al., J. Biol. Chem. 269:26988, 1994) may

5 be used. Cells are plated and allowed to attach to 96-well dishes in the same media (Ham's F12) with 10% FBS (fetal bovine serum). The cells are then washed, re-fed with serum depleted media that contains 0.1% (v/v) bovine serum albumin (BSA), and allowed to incubate for 24 hours. Immediately prior to dosing with compound, the cells are re-fed with the serum depleted media (without BSA). Test compounds, dissolved in DMSO, are diluted into the
10 media (final DMSO concentration 0.5% (v/v)). At the end of a 2 hour incubation, VEGF₁₆₅ (50 ng/ml final) is added to the media for an 8 minute incubation. The cells are washed and lysed in HNTG buffer (20 mM Hepes, pH 7.5, 150 mM NaCl, 0.2% Triton™ X-100, 10% glycerol, 0.2 mM PMSF (phenylmethylsulfonyl fluoride), 1 µg/ml pepstatin, 1 µg/ml leupeptin, 1 µg/ml aprotonin, 2 mM sodium pyrophosphate, 2 mM sodium orthovanadate). The extent of
15 phosphorylation of KDR is measured using an ELISA assay. The 96-well plates are coated with 1 µg per well of goat anti-rabbit antibody. Unbound antibody is washed off the plate and remaining sites are blocked with Superblock buffer (Pierce) prior to addition of the anti-flk-1 C-20 antibody (0.5 µg per plate, Santa Cruz). Any unbound antibody is washed off the plates prior to addition of the cell lysate. After a 2 hour incubation of the lysates with the flk-1
20 antibody, the KDR associated phosphotyrosine is quantitated by development with the HRP-conjugated PY-54 antibody and TMB, as described above. The ability of the compounds to inhibit the VEGF-stimulated autophosphorylation reaction by 50%, relative to VEGF-stimulated controls is reported as the IC₅₀ value for the test compound.

Administration of the compounds of the present invention (hereinafter the "active
25 compound(s)") can be effected by any method that enables delivery of the compounds to the site of action. These methods include oral routes, intraduodenal routes, parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion), topical, and rectal administration.

The amount of the active compound administered will be dependent on the subject
30 being treated, the severity of the disorder or condition, the rate of administration, the disposition of the compound and the discretion of the prescribing physician. However, an effective dosage is in the range of about 0.001 to about 100 mg per kg body weight per day, preferably about 1 to about 35 mg/kg/day, in single or divided doses. For a 70 kg human, this would amount to about 0.05 to about 7 g/day, preferably about 0.2 to about 2.5 g/day. In some instances, dosage levels
35 below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several small doses for administration throughout the day.

The active compound may be applied as a sole therapy or may involve one or more other anti-tumour substances, for example those selected from, for example, mitotic inhibitors,

5 for example vinblastine; alkylating agents, for example cis-platin, carboplatin and cyclophosphamide; anti-metabolites, for example 5-fluorouracil, cytosine arabinoside and hydroxyurea, or, for example, one of the preferred anti-metabolites disclosed in European Patent Application No. 239362 such as N-(5-[N-(3,4-dihydro-2-methyl-4-oxoquinazolin-6-ylmethyl)-N-methylamino]-2-thenoyl)-L-glutamic acid; growth factor inhibitors; cell cycle inhibitors;
10 intercalating antibiotics, for example adriamycin and bleomycin; enzymes, for example interferon; and anti-hormones, for example anti-estrogens such as Nolvadex™ (tamoxifen) or, for example anti-androgens such as Casodex™ (4'-cyano-3-(4-fluorophenylsulphonyl)-2-hydroxy-2-methyl-3'-(trifluoromethyl)propionanilide). Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate dosing of the individual components of the treatment.

15 The pharmaceutical composition may, for example, be in a form suitable for oral administration as a tablet, capsule, pill, powder, sustained release formulations, solution, suspension, for parenteral injection as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository. The pharmaceutical composition may be in unit dosage forms suitable for single administration of
20 precise dosages. The pharmaceutical composition will include a conventional pharmaceutical carrier or excipient and a compound according to the invention as an active ingredient. In addition, it may include other medicinal or pharmaceutical agents, carriers, adjuvants, etc.

Exemplary parenteral administration forms include solutions or suspensions of active compounds in sterile aqueous solutions, for example, aqueous propylene glycol or dextrose
25 solutions. Such dosage forms can be suitably buffered, if desired.

Suitable pharmaceutical carriers include inert diluents or fillers, water and various organic solvents. The pharmaceutical compositions may, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. Thus for oral administration, tablets containing various excipients, such as citric acid may be employed together with various
30 disintegrants such as starch, alginic acid and certain complex silicates and with binding agents such as sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tableting purposes. Solid compositions of a similar type may also be employed in soft and hard filled gelatin capsules. Preferred materials, therefor, include lactose or milk sugar and high molecular weight
35 polyethylene glycols. When aqueous suspensions or elixirs are desired for oral administration the active compound therein may be combined with various sweetening or flavoring agents, coloring matters or dyes and, if desired, emulsifying agents or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin, or combinations thereof.

Methods of preparing various pharmaceutical compositions with a specific amount of
40 active compound are known, or will be apparent, to those skilled in this art. For examples, see

- 5 Remington's Pharmaceutical Sciences, Mack Publishing Company, Easter, Pa., 15th Edition (1975).

The examples and preparations provided below further illustrate and exemplify the compounds of the present invention and methods of preparing such compounds. It is to be understood that the scope of the present invention is not limited in any way by the scope of the following examples and preparations. In the following examples molecules with a single chiral center, unless otherwise noted, exist as a racemic mixture. Those molecules with two or more chiral centers, unless otherwise noted, exist as a racemic mixture of diastereomers. Single enantiomers/diastereomers may be obtained by methods known to those skilled in the art.


- 15 Where HPLC chromatography is referred to in the preparations and examples below, the general conditions used, unless otherwise indicated, are as follows. The column used is a ZORBAX™ RXC18 column (manufactured by Hewlett Packard) of 150 mm distance and 4.6 mm interior diameter. The samples are run on a Hewlett Packard-1100 system. A gradient solvent method is used running 100 percent ammonium acetate / acetic acid buffer (0.2 M) to 100 percent acetonitrile over 10 minutes. The system then proceeds on a wash cycle with 100 percent acetonitrile for 1.5 minutes and then 100 percent buffer solution for 3 minutes. The flow rate over this period is a constant 3 ml / minute. In the following examples and preparations, "Et" means ethyl, "Ac" means acetyl, "Me" means methyl, and "Bu" means butyl.

25

Experimental Examples

Example 1

- 30 Synthesis of 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.



Example 1A

Trifluoro-methanesulfonic acid 8-(tert-butyl-dimethyl-silanyloxy)-quinolin-2-yl ester.

-47-

- 5 2,8-Quinolinediol (20.0 g, 124 mMol) was suspended in 500 mL of dichloromethane (DCM) under an atmosphere of dry nitrogen (N_2). To this solution was added imidazole (20.3 g, 298 mMol) followed by tert-butyldimethylsilyl chloride (20.6 g, 137 mMol) and 4-dimethylaminopyridine (1.50 g, 12.4 mMol). The reaction mixture was stirred overnight at ambient temperature after which time it was partitioned between DCM and 1% aqueous sodium bisulfate ($NaHSO_4$). The DCM layer was saved and washed two more times with 1% aqueous $NaHSO_4$, then aqueous saturated sodium bicarbonate ($NaHCO_3$) and finally brine. The DCM layer was dried over sodium sulfate (Na_2SO_4), filtered and concentrated under vacuum to give crude product (40 g) as a white solid. The solid was dissolved in 500 mL of anhydrous tetrahydrofuran (THF) under an atmosphere of dry N_2 . To this solution was added
- 10 N-phenyl-bis(trifluoromethanesulfonimide) (48.7 g, 136 mMol) and the solution was cooled to $0^\circ C$. To this solution was slowly added (3.2 g, 136 mMol) sodium hydride (60% in oil). After the addition was complete, the reaction mixture was warmed to ambient temperature. An additional 1.00 g sodium hydride (60% in oil) was added after one hour and stirred for an additional 30 minutes. The mixture was concentrated under vacuum and taken up in DCM.
- 15 20 Water (1.0 mL) was slowly added dropwise to quench any unreacted sodium hydride and then the reaction mixture was extracted twice from 0.1N aqueous sodium hydroxide ($NaOH$) and then washed with brine. The DCM layer was dried over Na_2SO_4 , filtered and concentrated under vacuum to give 57 g of the crude triflate **1A** as a yellow oil.

25

Example 1B

([8-(tert-Butyl-dimethyl-silanyloxy)-quinolin-2-yl]-(4-methoxy-2-nitro-phenyl)-amine.

- Trifluoro-methanesulfonic acid 8-(tert-butyl-dimethyl-silanyloxy)-quinolin-2-yl ester **1A** (9.81 g, 24.1 mMol) and 4-methoxy-2-nitroaniline (4.86 g, 28.9 mMol) were dissolved in 100 mL of dioxane under an atmosphere of dry N_2 . To this solution was added (11.0 g, 33.7 mMol) cesium carbonate (Cs_2CO_3), (900 mg, 1.45 mMol) racemic-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) and tris(dibenzylideneacetone)dipalladium (0) (883 mg, 0.964 mMol) and the reaction mixture was heated to $100^\circ C$ and reacted at this temperature for 4 hours. The mixture was then cooled to ambient temperature, concentrated under vacuum, treated with
- 30 DCM, filtered and concentrated under vacuum to give a red solid. The solid was chromatographed on flash silica gel eluting with hexanes/DCM (3:1) to give 7.25 g of the title compound **1B** as a red solid.

Example 1C

- 40 N^1 -[8-(tert-Butyl-dimethyl-silanyloxy)-quinolin-2-yl]-4-methoxy-benzene-1,2-diamine.

5
([8-(tert-Butyl-dimethyl-silanyloxy)-quinolin-2-yl]-(4-methoxy-2-nitro-phenyl)-amine **1B** (21.9 g, 51.3 mMol) was dissolved in a solution of 200 mL ethanol (EtOH) and 70 mL of THF under an atmosphere of dry N₂. To this solution was added 10% palladium on carbon (2.18 g) followed by the dropwise addition of 10 mL of anhydrous hydrazine. The reaction mixture was stirred
10 at ambient temperature for 2 hours after which time it was filtered through Celite™ and the Celite™ was washed with DCM. The combined filtrates were concentrated under vacuum and the resulting residue was partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was then washed again with saturated NaHCO₃ and then brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give 18.3 g of a tan solid as the title
15 compound **1C**.

Example 1D

2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-ol.

N¹-[8-(tert-Butyl-dimethyl-silanyloxy)-quinolin-2-yl]-4-methoxy-benzene-1,2-diamine **1C** (18.3
20 g, 46.1 mMol) was dissolved in 40 mL of 2-methoxyethanol under an atmosphere of dry N₂. To this solution was added formamidine acetate (5.28 g, 50.7 mMol) and the reaction mixture was heated to 125°C and reacted at this temperature for 1.5 hours. The solvent was removed under vacuum and the resulting solid was triturated with ethyl ether (Et₂O), dried under vacuum to give 13.3 g of a pink solid as the title compound **1D**.

25

Example 1E

Trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester.

2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-ol **1D** (13.9 g, 47.8 mMol) was dissolved in 100
30 mL of anhydrous THF under an atmosphere of dry N₂. To this solution was added N-phenyl-bis(trifluoromethanesulfonimide) (20.3 g, 47.8 mMol) and then the solution was subsequently cooled to 0°C. To this solution was slowly added (1.31 g, 54.9 mMol) sodium hydride (60% in oil). After the addition was complete the reaction mixture was warmed to ambient temperature. After 30 minutes, 500 mg more of sodium hydride (60% in oil) was added
35 followed by 3.50 g of N-phenyl-bis(trifluoromethanesulfonimide) and the reaction mixture was stirred at ambient temperature for 1 hour. The solvent was then removed under vacuum and the resulting residue was taken up in DCM. To this solution was slowly added 1.0 mL of water to decompose any unreacted sodium hydride. The mixture was subsequently partitioned between DCM and 0.1 N aqueous NaOH. The DCM layer was then washed again with 0.1 N

- 5 aqueous NaOH, followed by brine and then dried over magnesium sulfate (MgSO_4), filtered and concentrated under vacuum to give 20.7 g of a pink solid as the crude title compound **1E**.

Example 1F

{1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester.

10

- Trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** (15.0 g, 35.4 mMol) and piperidin-4-yl-carbamic acid tert-butyl ester (14.2 g, 70.9 mMol) were dissolved in 200 mL of dioxane under an atmosphere of dry N_2 . To this solution was added Cs_2CO_3 (16.2 g, 49.6 mMol), racemic-BINAP (1.28 g, 2.12 mMol) and
- 15 tris(dibenzylideneacetone)dipalladium (0) (1.29 g, 1.41 mMol) and the reaction mixture was heated to 100°C and reacted at this temperature overnight. The mixture was then cooled to ambient temperature, filtered, and concentrated under vacuum to give an orange foam. The foam was chromatographed on flash silica gel eluting with a gradient from ethyl acetate (EtOAc)/DCM (1:5) to EtOAc/DCM (7:3) give 12.3 g of the title compound **1F** as a slightly
- 20 yellow solid.

Example 1G

1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.


- 25 {1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester **1F** (8.40 g, 17.7 mMol) was dissolved in 50 mL of trifluoroacetic acid (TFA) under an atmosphere of dry N_2 . The reaction mixture was stirred at ambient temperature for 15 minutes after which time it was concentrated under vacuum to give a yellow oil. The oil was partitioned between DCM and 0.1N aqueous NaOH. The DCM layer was washed again with
- 30 0.1N aqueous NaOH. The DCM layer was dried over Na_2SO_4 , filtered and concentrated to give 5.85 g of the title compound **1** as a yellow solid.

- C.I. m/z 374 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.66 (s, 1 H), 8.37 (d, J = 8.9 Hz, 1 H), 8.30 (d, J = 8.7 Hz, 1 H), 7.68 (d, J = 8.9 Hz, 1 H), 7.47 (m, 2 H), 7.35 (d, J = 2.3 Hz, 1 H), 7.25 (m, 1 H), 7.06 (dd, J = 2.5, 8.9 Hz, 1 H), 3.91 (s, 3 H), 3.88 (m, 2 H), 2.90 (m, 3 H), 2.05 (m, 2 H), 1.83 (m, 2 H), 1.50 (brs, 2 H).
- 35

Example 2

1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-ol.

-50-



5


1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** (500 mg, 1.10 mMol) was dissolved in 10 mL of DCM under an atmosphere of dry N_2 . To this solution was added boron tribromide (300 μ L, 3.30 mMol) and the mixture was stirred overnight at ambient temperature. Then an additional 200 μ L of borontribromide was added and the mixture was stirred for two hours. The reaction mixture was then poured over crushed ice and the pH of the resulting solution was adjusted to 9 with the careful addition of sodium carbonate (Na_2CO_3). The slurry was filtered and the solid was washed with water followed by Et_2O and then dried under vacuum to give the title compound **2** as a yellow solid.

C.I. m/z 360 $[M+1]$; 1H NMR (DMSO) δ 9.07 (s, 1 H), 8.76 (d, $J = 8.9$ Hz, 1 H), 8.48 (d, $J = 8.9$ Hz, 1 H), 8.10 (d, $J = 8.9$ Hz, 1 H), 7.56 (d, $J = 7.4$ Hz, 1 H), 7.45 (m, 1 H), 7.26 (d, $J = 7.4$ Hz, 1 H), 7.01 (d, $J = 2.2$ Hz, 1 H), 6.95 (dd, $J = 2.2, 8.9$ Hz, 1 H), 3.72 (m, 2 H), 2.76 (m, 3 H), 1.88 (m, 2 H), 1.65 (m, 2 H).

20

Example 3

1-[2-[5-(Pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine.



Example 3A

{1-[2-(5-Hydroxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester.

25

1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-ol **2** (460 mg, 1.30 mMol) was dissolved in 5 mL of anhydrous DMF under an atmosphere of dry N_2 . To this solution was added di-tert-butylidicarbonate (279 mg, 1.30 mMol) and the reaction mixture was stirred at ambient temperature overnight. The reaction mixture was then concentrated under vacuum and partitioned between DCM and aqueous saturated $NaHCO_3$. The DCM layer was dried over Na_2SO_4 , filtered and concentrated under vacuum to give a yellow solid. The solid was

30

- 5 chromatographed on flash silica gel eluting with EtOAc to give 273 mg of the title compound **3A** as a yellow solid.

Example 3B

(1-{2-[5-(Pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-carbamic acid
tert-butyl ester.

10

- {1-[2-(5-Hydroxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester **3A** (73 mg, 0.16 mMol) was dissolved in 1 mL of anhydrous DMF under an atmosphere of dry N₂. To this solution was added (37 mg, 0.17 mMol) potassium bis(trimethyl)silylamide (95%) followed by 2-picolyl chloride (25 µL, 0.17 mMol). The reaction mixture was stirred
15 overnight at ambient temperature after which time the reaction mixture was concentrated under vacuum and then partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow gel. The gel was chromatographed on flash silica gel eluting with a gradient from DCM to DCM/MeOH (98:2) to give 55 mg of the title compound **3B**.

20

Example 3C

1-{2-[5-(Pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.

- (1-{2-[5-(Pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-carbamic acid
25 tert-butyl ester **3B** (55 mg, 0.094 mMol) was dissolved in 1 mL of TFA under an atmosphere of dry N₂ and stirred for 15 minutes at ambient temperature. The reaction mixture was then concentrated under vacuum to give an oil which was subsequently partitioned between 0.1 N aqueous NaOH and DCM. The DCM layer was dried over Na₂SO₄, filtered and concentrated under vacuum to give 38.9 mg of a yellow film as the title compound **3**.


- 30 C.I. m/z 451 [M+1]; ¹H NMR (CDCl₃) δ 8.64 (s, 1 H), 8.60 (m, 1 H), 8.42 (dd, J = 1.5, 8.9 Hz, 1 H), 8.27 (d, J = 8.9 Hz, 1 H), 7.64-7.72 (m, 2 H), 7.57 (d, J = 7.9 Hz, 1 H), 7.45 (m, 2 H), 7.40 (d, J = 2.5 Hz, 1 H), 7.18-7.25 (m, 3 H), 5.29 (s, 2 H), 3.87 (m, 2 H), 2.92 (m, 3 H), 2.04 (m, 2 H), 1.80 (m, 2 H), 1.60 (brs, 2 H).

35

Example 4

{1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethyl-amine.

-52-



5


1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** (160 mg, 0.43 mMol) was dissolved in 2 mL of chloroform under an atmosphere of dry N₂. To this solution was added 50 µL of 37% aqueous formaldehyde and 100 µL formic acid and the reaction mixture was subsequently heated to 65°C and reacted at this temperature for 4 hours. The reaction mixture was then partitioned between DCM and 0.1 N aqueous NaOH. The DCM layer was dried over Na₂SO₄, filtered and concentrated under vacuum to give an orange solid as the title compound **4**.

C.I. m/z 402 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.42 (d, J = 8.9 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.9 Hz, 1 H), 7.45 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.25 (m, 1 H), 7.05 (dd, J = 2.5, 8.9 Hz, 1 H), 4.00 (m, 2 H), 3.89 (s, 3 H), 2.80 (m, 2 H), 2.38 (m, 1 H), 2.37 (s, 6 H), 1.99 (m, 4 H).

15

Example 5

{4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine.



20

Example 5A

4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzaldehyde.

25 Trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** (265 mg, 0.630 mMol) was dissolved in 3 mL of dioxane under an atmosphere of dry N₂. To this solution was added 4-formylbenzene boronic acid (145 mg, 0.940 mMol), potassium phosphate (267 mg, 1.26 mMol) and tetrakis(triphenylphosphine)palladium (0) (36 mg, 0.032 mMol). The reaction mixture was heated to 105°C and reacted at this temperature overnight.

30 The reaction mixture was then cooled to ambient temperature, concentrated under vacuum

- 5 and partitioned between DCM and aqueous saturated NaHCO_3 . The DCM layer was then washed with brine, dried over Na_2SO_4 , filtered and concentrated under vacuum to give a yellow solid as the title compound **5A** which was pushed on without further purification.


Example 5B

- 10 {4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine.

4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzaldehyde **5A** (120 mg, 0.32 mMol) was dissolved in 2 mL of methanol under an atmosphere of dry N_2 . To this solution was added 800 μL of a solution of 2.0 M methylamine in methanol and then acetic acid (AcOH) was added dropwise until the pH of the solution was ~5. To this solution was added (42 mg, 0.64 mMol) sodium cyanoborohydride (NaCNBH_3) and the reaction mixture was stirred at ambient temperature overnight. The reaction mixture was then concentrated under vacuum and subsequently partitioned between DCM and 0.1 N aqueous NaOH . The DCM layer was washed again with 0.1 N aqueous NaOH , then dried over MgSO_4 , filtered and concentrated under vacuum to give 220 mg of a green residue. The residue was chromatographed on flash silica gel eluting with a gradient DCM/MeOH (5/95) to DCM/MeOH (15/85) to DCM/MeOH/ NH_4OH (15/84.5/0.5) to give 50 mg of a white solid of the title compound **5**.
C.I. m/z 395 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.55 (s, 1 H), 8.29 (d, J = 8.9 Hz, 1 H), 7.92 (d, J = 8.9 Hz, 1 H), 7.79 (d, J = 8.1 Hz, 1 H), 7.74 (m, 1 H), 7.63 (m, 3 H), 7.56 (m, 1 H), 7.44 (d, J = 8.1 Hz, 2 H), 7.23 (d, J = 2.3 Hz, 1 H), 6.73 (dd, J = 2.3, 8.9, 1 H), 3.88 (s, 2 H), 3.83 (s, 3 H), 2.53 (s, 3 H), 2.43 (brs, 1H).

Example 6

{4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine.




- 30 The same procedure that used for Example 5 was followed with the exception that 2.0 M dimethylamine in methanol was used in the place of 2.0 M methylamine in methanol in Example 5B to give the title compound **6** as a white solid.

C.I. m/z 409 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.56 (s, 1 H), 8.29 (d, J = 8.7 Hz, 1 H), 7.98 (d, J = 9.1 Hz, 1 H), 7.79 (d, J = 8.1 Hz, 1 H), 7.76 (m, 1 H), 7.63 (m, 3 H), 7.56 (m, 1 H), 7.45 (d, J = 7.9

- 5 Hz, 2 H), 7.25 (d, J = 2.3 Hz, 1 H), 6.74 (dd, J = 2.3, 8.9, 1 H), 3.84 (s, 3 H), 3.57 (s, 2 H), 2.35 (s, 6 H).

Example 7

Cyclopropyl-{4-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine.




10

- 4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzaldehyde **5A** (50 mg, 0.13 mMol) was dissolved in 500 μ L of dichloroethane (DCE) under an atmosphere of dry N_2 . To this solution was added 80 μ L of AcOH, cyclopropylamine (50 μ L, 0.65 mMol) and (42 mg, 0.20 mMol) sodium triacetoxyborohydride ($NaHB(OAc)_3$). The reaction mixture was stirred at ambient
- 15 temperature for 2 hours after which time an additional 20 μ L of cyclopropylamine and 20 mg of $NaHB(OAc)_3$ were added and the resulting mixture was stirred overnight. The reaction mixture was then partitioned between aqueous 0.1 N NaOH and DCM. The DCM layer washed again with aqueous 0.1 N NaOH and then dried over Na_2SO_4 , filtered and concentrated under vacuum to give a 60 mg of the title compound **7** as a yellow solid.
- 20 C.I. m/z 421 [M+1]; 1H NMR ($CDCl_3$) δ 8.58 (s, 1 H), 8.33 (d, J = 8.7 Hz, 1 H), 7.96 (d, J = 9.1 Hz, 1 H), 7.83 (d, J = 7.7 Hz, 1 H), 7.77 (d, J = 7.3 Hz, 1 H), 7.64 (m, 3 H), 7.59 (m, 1 H), 7.45 (d, J = 7.9 Hz, 2 H), 7.25 (m, 1 H), 6.75 (dd, J = 2.3, 8.9, 1 H), 3.97 (s, 2 H), 3.86 (s, 3 H), 2.26 (m, 1 H), 1.95 (brs, 1 H), 0.49 (m, 4 H).

25

Example 8

tert-Butyl-{4-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine.




- 5 The same procedure that used for Example 7 was followed with the exception that tert-butylamine was used in the place of cyclopropylamine to give the title compound **8** as a yellow solid.

C.I. m/z 437 [M+1]; ¹H NMR (CDCl₃) δ 8.59 (s, 1 H), 8.33 (d, J = 8.9 Hz, 1 H), 7.95 (d, J = 8.9 Hz, 1 H), 7.82 (m, 1 H), 7.77 (m, 1 H), 7.65 (m, 3 H), 7.58 (m, 1 H), 7.49 (d, J = 8.1 Hz, 2 H),
10 7.26 (d, J = 2.3 Hz, 1 H), 6.79 (dd, J = 2.3, 8.9, 1 H), 3.86 (s, 5 H), 1.24 (s, 9 H).

Example 9

4-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzylamine.




- 15 The same procedure that used for Example 5 was followed with the exception that ammonium acetate was used in the place of 2.0 M methylamine in methanol in Example 5B to give a yellow solid. The solid was chromatographed on flash silica gel eluting with a gradient from DCM/MeOH (5/95) to DCM/MeOH (15/85) to DCM/MeOH/NH₄OH (15/94.5/0.5) to give the title compound **9** as a white solid.
- 20 C.I. m/z 381 [M+1]; ¹H NMR (CDCl₃) δ 8.59 (s, 1 H), 8.35 (d, J = 8.9 Hz, 1 H), 7.94 (d, J = 8.9 Hz, 1 H), 7.84 (m, 1 H), 7.78 (dd, J = 1.5, 7.3 Hz, 1 H), 7.66 (m, 3 H), 7.60 (m, 1 H), 7.46 (d, J = 8.1 Hz, 2 H), 7.26 (d, J = 2.5 Hz, 1 H), 6.76 (dd, J = 2.5, 8.9, 1 H), 4.00 (s, 2 H), 3.85 (s, 3 H), 1.78 (brs, 2H).

25

Example 10

1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.




- The same procedure that used for Example 1 was followed with the exception that 4-ethoxy-2-nitroaniline was used in the place of that 4-methoxy-2-nitroaniline in example 1B to give the
30 title compound **10** as a yellow solid.

- 5 C.I. m/z 388 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.36 (d, J = 8.9 Hz, 1 H), 8.26 (d, J = 8.7 Hz, 1 H), 7.64 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.32 (d, J = 2.5 Hz, 1 H), 7.25 (m, 1 H), 7.05 (dd, J = 2.5, 8.9 Hz, 1 H), 4.10 (q, J = 7.0 Hz, 2 H), 3.86 (m, 2 H), 2.90 (m, 3 H), 2.04 (m, 2 H), 1.79 (m, 2 H), 1.45 (t, J = 7.0 Hz, 3 H).

10

Example 11

{1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethyl-amine.




- The same procedure that used in Example 4 was followed with the exception 1-[2-(5-ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **10** was used in the place of 1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** to give the title compound **11** as a yellow solid.

- C.I. m/z 416 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.39 (d, J = 8.9 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.45 (m, 2 H), 7.32 (d, J = 2.5 Hz, 1 H), 7.24 (m, 1 H), 7.05 (dd, J = 2.5, 8.9 Hz, 1 H), 4.10 (q, J = 7.0 Hz, 2 H), 4.00 (m, 2 H), 2.80 (m, 2 H), 2.40 (m, 1 H), 2.38 (s, 6 H), 2.00 (m, 4 H), 1.46 (t, J = 7.0 Hz, 3 H).

Example 12

1-[2-(5-Trifluoromethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.




25

The same procedure that used in Example 1 was followed with the exception that 2-nitro-4-(trifluoromethoxy)aniline was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **12** as a yellow solid.

- 5 C.I. m/z 428 [M+1]; ^1H NMR (CDCl_3) δ 8.74 (s, 1 H), 8.50 (d, $J = 8.9$ Hz, 1 H), 8.33 (d, $J = 8.7$ Hz, 1 H), 7.74 (d, $J = 1.0$ Hz, 1 H), 7.67 (d, $J = 8.7$ Hz, 1 H), 7.47 (m, 2 H), 7.32 (m, 1 H), 7.25 (m, 1 H), 3.87 (m, 2 H), 2.90 (m, 3 H), 2.05 (m, 2 H), 1.77 (m, 2 H), 1.71 (brs, 2 H).

Example 13

- 10 (4-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl)-methyl-amine.




The same procedure that used for Example 5 was followed with the exception that 4-ethoxy-2-nitroaniline was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound 13 as a tan solid.

- 15 C.I. m/z 409 [M+1]; ^1H NMR (CDCl_3) δ 8.54 (s, 1 H), 8.26 (d, $J = 8.7$ Hz, 1 H), 7.92 (d, $J = 9.1$ Hz, 1 H), 7.77 (d, $J = 8.1$ Hz, 1 H), 7.72 (d, $J = 7.1$ Hz, 1 H), 7.62 (m, 3 H), 7.54 (m, 1 H), 7.52 (d, $J = 8.1$ Hz, 2 H), 7.21 (d, $J = 2.5$ Hz, 1 H), 6.73 (dd, $J = 2.5, 8.9$, 1 H), 4.04 (q, $J = 7.0$ Hz, 2 H), 3.91 (s, 2 H), 2.97 (brs, 1H), 2.55 (s, 3 H), 1.41 (t, $J = 7.0$ Hz, 3 H).

20

Example 14

Cyclopropyl-[4-[2-(5-ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl]-amine.



The same procedure that used for Example 7 was followed with the exception that 4-ethoxy-2-nitroaniline was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound 14 as a yellow solid.


25

C.I. m/z 435 [M+1]; ^1H NMR (CDCl_3) δ 8.57 (s, 1 H), 8.32 (d, $J = 8.7$ Hz, 1 H), 7.94 (d, $J = 8.9$ Hz, 1 H), 7.83 (m, 1 H), 7.77 (m, 1 H), 7.65 (m, 3 H), 7.58 (m, 1 H), 7.45 (d, $J = 8.1$ Hz, 2 H),

- 5 7.25 (m, 1 H), 6.76 (dd, J = 2.5, 9.1, 1 H), 4.06 (q, J = 7.0 Hz, 2 H), 3.97 (s, 2 H), 2.28 (m, 1 H), 1.93 (brs, 1 H), 1.44 (t, J = 7.0 Hz, 3 H), 0.48 (m, 4 H).

Example 15

- 10 tert-Butyl-{4-[2-(5-ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine.




The same procedure that used for Example 8 was followed with the exception that 4-ethoxy-2-nitroaniline was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **15** as a yellow solid.

- 15 C.I. m/z 451 [M+1]; ¹H NMR (CDCl₃) δ 8.55 (s, 1 H), 8.33 (d, J = 8.7 Hz, 1 H), 7.92 (d, J = 10.1 Hz, 1 H), 7.79 (dd, J = 1.3, 8.1 Hz, 1 H), 7.71 (d, J = 7.1 Hz, 1 H), 7.65 (m, 3 H), 7.55 (m, 1 H), 7.48 (d, J = 8.1 Hz, 2 H), 7.23 (d, J = 2.5 Hz, 1 H), 6.78 (dd, J = 2.5, 8.9, 1 H), 4.06 (q, J = 7.0 Hz, 2 H), 3.89 (s, 2 H), 1.43 (t, J = 7.0 Hz, 3 H), 1.28 (s, 9 H).

Example 16

- 20 {4-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine.




The same procedure that used for Example 6 was followed with the exception that 4-ethoxy-2-nitroaniline was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **16** as a yellow solid.

- 25 C.I. m/z 423 [M+1]; ¹H NMR (CDCl₃) δ 8.57 (s, 1 H), 8.31 (d, J = 8.7 Hz, 1 H), 7.96 (d, J = 8.9 Hz, 1 H), 7.80 (m, 1 H), 7.75 (m, 1 H), 7.64 (m, 3 H), 7.56 (m, 1 H), 7.42 (d, J = 7.9 Hz, 2 H),

- 5 7.20 (d, J = 2.4 Hz, 1 H), 6.73 (dd, J = 2.4, 8.9, 1 H), 4.06 (q, J = 7.0 Hz, 2 H), 3.55 (s, 2 H), 2.31 (s, 6 H), 1.41 (t, J = 7.0 Hz, 3 H).

Example 17

1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one.




10

The same procedure that used for Example 1 was followed with the exception that 4-piperdone was used in the place of piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound 17 as a white solid.

- C.I. m/z 373 [M+1]; ¹H NMR (CDCl₃) δ 8.67 (s, 1 H), 8.35 (m, 2 H), 7.72 (d, J = 8.9 Hz, 1 H),
15 7.55 (dd, J = 1.3, 8.2, 1 H), 7.50 (m, 1 H), 7.36 (d, J = 2.5 Hz, 1 H), 7.28 (m, 1 H), 7.04 (dd, J = 2.5, 8.9 Hz, 1 H), 3.90 (s, 3 H), 3.71 (m, 4 H), 2.79 (m, 4 H).

Example 18

1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one.



20


The same procedure that was used for example 17 was followed except that 4-ethoxy-2-nitroaniline was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound 18 as a white solid.

- C.I. m/z 387 [M+1]; ¹H NMR (CDCl₃) δ 8.57 (s, 1 H), 8.30 (m, 2 H), 7.65 (d, J = 8.7 Hz, 1 H),
25 7.50 (dd, J = 1.3, 8.1, 1 H), 7.44 (m, 1 H), 7.30 (d, J = 2.3 Hz, 1 H), 7.23 (m, 1 H), 6.99 (dd, J = 2.5, 8.9 Hz, 1 H), 4.07 (q, J = 7.0 Hz, 2 H), 3.65 (m, 4 H), 2.74 (m, 4 H), 1.42 (t, J = 7.0 Hz, 3 H).

5

Example 19

1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol.




1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one **18** (140 mg, 0.36 mMol) was dissolved in 1.5 mL of methanol under an atmosphere of dry N₂. To this solution was added sodium borohydride (NaBH₄) (14 mg, 0.36 mMol) and the solution was stirred at ambient temperature overnight. The reaction mixture was then concentrated under vacuum and partitioned between DCM and saturated aqueous NaHCO₃. The DCM layer was then washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give a green foam. The foam was chromatographed on flash silica gel eluting with a gradient from MeOH/DCM (1/99) to MeOH/DCM (4/96) to give the title compound **19** as a yellow foam.

C.I. m/z 389 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.39 (d, J = 8.9 Hz, 1 H), 8.30 (d, J = 8.7 Hz, 1 H), 7.68 (d, J = 8.9 Hz, 1 H), 7.47 (m, 2 H), 7.33 (d, J = 2.3 Hz, 1 H), 7.25 (m, 1 H), 7.07 (dd, J = 2.5, 8.9 Hz, 1 H), 4.13 (q, J = 7.0 Hz, 2 H), 3.97 (m, 1 H), 3.75 (m, 2 H), 3.10 (m, 2 H), 2.10 (m, 2 H), 1.96 (m, 2 H), 1.46 (t, J = 7.0 Hz, 3 H).

20

Example 20

Cyclopropyl-{1-[2-(5-ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-amine.



1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one **18** (130 mg, 0.340 mMol) was dissolved in 1.5 mL of DCE under an atmosphere of dry N₂. To this solution was added cyclopropyl amine (110 μL, 1.70 mMol) and 200 μL of AcOH and the solution was stirred for 10 minutes. To this solution was added NaHB(OAc)₃ (107 mg, 0.50 mMol) and the solution was stirred at ambient temperature for 5 hours. The reaction mixture was then partitioned between DCM and aqueous 0.1 N NaOH. The DCM layer was again washed with aqueous


- 5 0.1 N NaOH, dried over Na₂SO₄, filtered and concentrated under vacuum to give 150 mg of a green foam. The foam was chromatographed on flash silica gel eluting with a gradient from MeOH/DCM (2/99) to MeOH/DCM (4/96) to give the title compound **20** as a yellow solid.

C.I. m/z 428 [M+1]; ¹H NMR (CDCl₃) δ 8.64 (s, 1 H), 8.38 (d, J = 9.1 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.45 (m, 2 H), 7.32 (d, J = 2.5 Hz, 1 H), 7.25 (m, 1 H), 7.07 (dd, J = 2.5, 9.1 Hz, 1 H), 4.13 (q, J = 7.0 Hz, 2 H), 3.91 (m, 2 H), 2.90 (m, 3 H), 2.25 (m, 1 H), 2.16 (m, 2 H), 1.83 (m, 2 H), 1.46 (t, J = 7.0 Hz, 3 H), 0.43 (m, 2 H), 0.42 (m, 2 H).

Example 21

tert-Butyl-{1-[2-(5-ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-amine.

15




The same procedure that was used for example 20 was followed except tert-butylamine was used in the place of cyclopropylamine to give the title compound **21** as a yellow solid.

C.I. m/z 444 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.35 (d, J = 8.9 Hz, 1 H), 8.28 (d, J = 8.9 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.45 (m, 2 H), 7.32 (d, J = 2.5 Hz, 1 H), 7.24 (m, 1 H), 7.08 (dd, J = 2.5, 8.91 Hz, 1 H), 4.13 (q, J = 7.0 Hz, 2 H), 3.90 (m, 2 H), 2.91 (m, 2 H), 2.79 (m, 1 H), 2.00 (m, 2 H), 1.88 (m, 2 H), 1.47 (t, J = 7.0 Hz, 3 H), 1.17 (s, 9 H).

Example 22

25 {1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine.



1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one **17** (200 mg, 0.540 mMol) was dissolved in 2 mL of MeOH under an atmosphere of dry N₂. To this solution was added 1.34 mL of a 2.0 M solution of methylamine in methanol and AcOH was then added until the pH=5. To this solution was added 95% NaCNBH₃ (93 mg, 0.540 mMol) and the reaction


30

5 mixture was stirred at ambient temperature for 4 hours. The reaction mixture was then concentrated under vacuum and subsequently partitioned between DCM and aqueous 0.1 N NaOH. The DCM layer was then washed again with aqueous 0.1 N NaOH, dried over Na_2SO_4 , filtered and concentrated under vacuum to give a green foam. The foam was chromatographed on flash silica gel eluting with a gradient from MeOH/DCM (5/95) to MeOH/DCM (10/90) to MeOH/DCM/ NH_4OH (10/89/1) give the title compound **22** as a green solid.

C.I. m/z 388 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.65 (s, 1 H), 8.38 (d, $J = 8.9$ Hz, 1 H), 8.29 (d, $J = 8.7$ Hz, 1 H), 7.67 (d, $J = 8.7$ Hz, 1 H), 7.46 (m, 2 H), 7.33 (d, $J = 2.3$ Hz, 1 H), 7.25 (m, 1 H), 7.08 (dd, $J = 2.5, 8.9$ Hz, 1 H), 3.93 (m, 2 H), 3.89 (s, 3 H), 2.90 (m, 2 H), 2.62 (m, 1 H), 2.51 (s, 3 H), 2.14 (m, 2 H), 1.82 (m, 2 H), 1.75 (brs, 1 H).

Example 23

2-(5-Methoxy-benzoimidazol-1-yl)-8-(1-oxa-6-aza-spiro[2.5]oct-6-yl)-quinoline.




20 Trimethyl sulfonium iodide (326 mg, 1.60 mMol) was dissolved in 6 mL of anhydrous dimethylsulfoxide (DMSO) under an atmosphere of dry N_2 . To this solution was added (67.7 mg, 1.7 mMol) sodium hydride (60% in oil) and the reaction mixture was stirred for 10 minutes at ambient temperature. To this solution was added a solution of 1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one **17** (450 mg, 1.21 mMol) in 4 mL of anhydrous DMSO and the reaction mixture was subsequently stirred overnight at ambient temperature. The reaction mixture was then partitioned between EtOAc and water. The EtOAc layer was washed with 3 more times with water and then with brine. The EtOAc layer was dried over Na_2SO_4 , filtered and concentrated under vacuum to give a yellow foam as the title compound **23**.

30 C.I. m/z 387 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.64 (s, 1 H), 8.39 (d, $J = 8.9$ Hz, 1 H), 8.31 (d, $J = 8.9$ Hz, 1 H), 7.68 (d, $J = 8.7$ Hz, 1 H), 7.48 (m, 2 H), 7.34 (d, $J = 2.3$ Hz, 1 H), 7.29 (m, 1 H), 7.06 (dd, $J = 2.5, 8.9$ Hz, 1 H), 3.90 (s, 3 H), 3.68 (m, 2 H), 3.41 (m, 2 H), 2.78 (s, 2 H), 2.32 (m, 2 H), 1.75 (m, 2 H).

5

Example 24

4-Dimethylaminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol.




2-(5-Methoxy-benzoimidazol-1-yl)-8-(1-oxa-6-aza-spiro[2.5]oct-6-yl)-quinoline **23** (250 mg, 0.647 mMol) and 2 mL of 2.0 M dimethylamine in THF were suspended in 2 mL of methanol in a pressure vial. The vial was capped, heated to 65°C and reacted at this temperature for 2 hours. The reaction mixture was concentrated under vacuum to give a yellow solid. The solid was chromatographed on flash silica gel eluting with a gradient from MeOH/DCM (10/90) to MeOH/DCM/NH₄OH (10/89/1) give 160 mg of the title compound **24** as a yellow solid.

C.I. m/z 432 [M+1]; ¹H NMR (CDCl₃) δ 8.64 (s, 1 H), 8.39 (d, J = 8.9 Hz, 1 H), 8.29 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.30 (m, 1 H), 7.01 (dd, J = 2.5, 8.9 Hz, 1 H), 3.89 (s, 3 H), 3.68 (m, 2 H), 3.24 (m, 2 H), 2.42 (s, 8 H), 2.00 (m, 2 H), 1.90 (brs, 1 H), 1.79 (m, 2 H).

20

Example 25

1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-4-methylaminomethyl-piperidin-4-ol.



The same procedure that was used in example 24 was followed except that 2.0 M methylamine in methanol was used in the place of 2.0 M dimethylamine in THF to give the title compound **25**


C.I. m/z 418 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.32 (m, 2 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.33 (d, J = 2.3 Hz, 1 H), 7.29 (m, 1 H), 7.02 (dd, J = 2.5, 8.9 Hz, 1 H), 3.90 (s, 3 H), 3.70 (m, 2 H), 3.23 (m, 2 H), 2.65 (s, 2 H), 2.54 (s, 3 H), 2.04 (m, 2 H), 1.79 (m, 2 H).

30

Example 26

4-Aminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol.

-64-



5


The same procedure that was used in example 24 was followed except that ammonium hydroxide was used in the place of 2.0 M dimethylamine in THF and the solvent was THF rather than methanol to give the title compound **26**.

C.I. m/z 404 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.30 (m, 2 H), 7.66 (d, J = 8.9 Hz, 1 H), 7.47 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.29 (m, 1 H), 7.01 (dd, J = 2.3, 8.9 Hz, 1 H), 3.89 (s, 3 H), 3.71 (m, 2 H), 3.23 (m, 2 H), 2.78 (s, 2 H), 1.96 (m, 2 H), 1.79 (m, 2 H).

10

Example 27

1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-pyrrolidin-3-ylamine.



15

The same procedure that was used in example 1 was followed except that (+/-)-3-(t-butoxycarbonylamino)-pyrrolidine was used in the place of the piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound **27** as a yellow solid.

C.I. m/z 360 [M+1]; ¹H NMR (CDCl₃) δ 8.61 (s, 1 H), 8.23 (d, J = 8.7 Hz, 1 H), 7.83 (d, J = 8.9 Hz, 1 H), 7.59 (d, J = 8.7 Hz, 1 H), 7.40 (m, 1 H), 7.33 (d, J = 2.3 Hz, 1 H), 7.19 (d, J = 8.5 Hz, 1 H), 7.00 (dd, J = 2.5, 8.9 Hz, 1 H), 6.83 (d, J = 7.9 Hz, 1 H), 4.00 (m, 1 H), 3.90 (m, 1 H), 3.88 (s, 3 H), 3.75 (m, 2 H), 3.56 (m, 1 H), 2.24 (m, 1 H), 1.92 (brs, 2 H), 1.82 (m, 1 H).


20

Example 28

1-(2-Benzoimidazol-1-yl-quinolin-8-yl)-piperidin-4-ylamine.

25

-65-



5

Example 28A

8-Benzyloxy-quinolin-2-ylamine.

- 10 2-Amino-8-hydroxyquinoline (20.0 g, 122 mMol) was dissolved in 50 mL of anhydrous DMF under an atmosphere of dry N₂. The reaction mixture was cooled to 0°C after which time (4.89 g, 122 mMol) sodium hydride (60% in oil) was slowly added and the reaction mixture was stirred for 30 minutes. To this solution was added benzyl bromide (14.5 mL, 122 mMol) and the reaction mixture was subsequently warmed to ambient temperature and stirred overnight.
- 15 The reaction mixture was then filtered and the resulting solid was washed with Et₂O and then dried under vacuum to give 29.4 g of the title compound **28A** as a white solid.

Example 28B

(8-Benzyloxy-quinolin-2-yl)-(2-nitro-phenyl)-amine.

20

- 8-Benzyloxy-quinolin-2-ylamine **28A** (3.50 g, 14.0 mMol) and 1-bromo-2-nitrobenzene (3.20 g, 15.4 mMol) were dissolved in 70 mL of dioxane under an atmosphere of dry N₂. To this solution was added Cs₂CO₃ (18.3 g, 56.0 mMol), racemic-BINAP (1.00 g, 1.68 mMol) and (513 mg, 0.560 mMol) tris(dibenzylideneacetone)dipalladium (0). The reaction mixture was
- 25 heated to 100°C and reacted at this temperature overnight. The mixture was then cooled to ambient temperature, concentrated under vacuum, treated with DCM, filtered and concentrated under vacuum to give a red solid. The solid was chromatographed on flash silica gel eluting with DCM to give 5.16 g of the title compound **28B** as an orange solid.

30

Example 28C

2-(2-Amino-phenylamino)-quinolin-8-ol.

- (8-Benzyloxy-quinolin-2-yl)-(2-nitro-phenyl)-amine **28B** (5.16 g, 13.9 mMol) was suspended in 60 mL of EtOH under an atmosphere of dry N₂. To this solution was added ammonium
- 35 formate (17.5 g, 278 mMol) and 550 mg of 20% palladium hydroxide on carbon. The reaction

- 5 mixture was heated to 78°C and stirred at this temperature for 2 hours after which time it was cooled to ambient temperature and filtered through Celite™. The Celite™ was washed with ethanol, the filtrates were combined and concentrated under vacuum to give the title compound **28C** which was subsequently carried on crude to the next step.

10

Example 28D

1-(2-Benzoimidazol-1-yl-quinolin-8-yl)-piperidin-4-ylamine.

The same procedure that was used in example 1 except that compound 2-(2-amino-phenylamino)-quinolin-8-ol **28C** was used in the place of 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-ol **1D** in example 1E to give the title compound **28**.


15

C.I. m/z 344 [M+1]; ¹H NMR (CDCl₃) δ 8.71 (s, 1 H), 8.43 (m, 1 H), 8.31 (d, J = 8.9 Hz, 1 H), 7.89 (m, 1 H), 7.71 (d, J = 8.7 Hz, 1 H), 7.36-7.49 (m, 4 H), 7.24 (m, 1 H), 3.89 (m, 2 H), 2.90 (m, 3 H), 2.04 (m, 2 H), 1.82 (m, 2 H), 1.28 (brs, 2 H).

20

Example 29

1-(2-Imidazo[4,5-b]pyridin-3-yl-quinolin-8-yl)-piperidin-4-ylamine.



The same procedure that was used in example 28 that 2-chloro-3-nitropyridine was used in the place of 1-bromo-2-nitrobenzene in example 28B to give the title compound **29**.


- 25 C.I. m/z 345 [M+1]; ¹H NMR (CDCl₃) δ 9.36 (s, 1 H), 9.00 (d, J = 8.9 Hz, 1 H), 8.49 (dd, J = 1.4, 4.7 Hz, 1 H), 8.34 (d, J = 8.9 Hz, 1 H), 8.15 (dd, J = 1.4, 8.1, 1 H), 7.45 (m, 2 H), 7.33 (dd, J = 4.8, 8.1 Hz, 1 H), 7.17 (dd, J = 1.7, 7.1 Hz, 1 H), 3.93 (m, 2 H), 2.98 (m, 1 H), 2.91 (m, 2 H), 2.12 (m, 2 H), 1.98 (brs, 2 H), 1.88 (m, 2 H).

30

Example 30

1-(2-[5-(4-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine.

-67-



5

Example 30A

4'-Methoxy-3-nitro-biphenyl-4-ylamine.

- 10 To a solution of 30 mL of water and 40 mL of dioxane were added 4-methoxyphenylboronic acid (1.69 g, 11.1 mMol), 4-bromo-2-nitroaniline (2.18 g, 10.0 mMol), tetrakis(triphenylphosphine)palladium (0) (580 mg, 0.502 mMol) and Na₂CO₃ (6.00 g, 56.6 mMol) under an atmosphere of dry N₂. The reaction mixture was subsequently heated to 80°C and stirred at this temperature overnight. The reaction mixture was then partitioned
- 15 between DCM and water. The DCM layer was then washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum. The resulting residue was chromatographed on flash silica gel eluting with a gradient starting from DCM/hexanes (1:1) to DCM to give 2.38 g of the title compound **30A**.

20

Example 30B

1-{2-[5-(4-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.

- The same procedure that was used in example 1 was followed except that 4'-methoxy-3-nitro-biphenyl-4-ylamine was used in the place of 4-methoxy-2-nitroaniline in example 1B to give
- 25 the title compound **30**.


C.I. m/z 450 [M+1]; ¹H NMR (CDCl₃) δ 8.71 (s, 1 H), 8.43 (d, J = 8.7 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 8.01 (d, J = 1.5 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.62 (m, 3 H), 7.45 (m, 2 H), 7.35 (dd, J = 1.3, 5.8 Hz, 1 H), 6.99 (dd, J = 2.1, 6.9 Hz, 2 H), 3.98 (m, 2 H), 3.83 (s, 3 H), 2.88 (m, 3 H), 2.04 (m, 2 H), 1.83 (m, 2 H), 1.77 (brs, 2 H).

30

Example 31

(4-{2-[5-(4-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-methyl-amine.

5




The same procedure that was used for example 5 was followed except that compound 4'-methoxy-3-nitro-biphenyl-4-ylamine **30A** was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **31**.

- 10 C.I. m/z 471 [M+1]; ¹H NMR (CDCl₃) δ 8.60 (s, 1 H), 8.27 (d, J = 8.9 Hz, 1 H), 8.02 (d, J = 8.7 Hz, 1 H), 7.91 (d, J = 1.5 Hz, 1 H), 7.76 (m, 2 H), 7.64 (m, 3 H), 7.47-7.57 (m, 5 H), 7.31 (dd, J = 1.7, 8.5 Hz, 1 H), 6.96 (m, 2 H), 3.91 (s, 2 H), 3.82 (s, 3 H), 2.87 (brs, 1 H), 2.55 (s, 3 H).

Example 32

- 15 1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.




- 20 {1-[2-(5-Hydroxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester **3A** (200 mg, 0.435 mMol) was dissolved in 1.5 mL of anhydrous DMF under an atmosphere of dry N₂. To this solution was added Cs₂CO₃ (170 mg, 0.520 mMol) followed by cyclopropyl methane bromide (46 μL, 0.48 mMol). The reaction mixture was subsequently heated to 65°C and stirred at this temperature for 4 hours. The reaction mixture was then cooled to ambient temperature and partitioned between EtOAc and water. The EtOAc layer was washed 4 more times with water and then with brine. The EtOAc was then dried over Na₂SO₄, filtered and concentrated under vacuum and the resulting green oil was chromatographed on flash silica gel eluting with MeOH/DCM (2:98) to give the a green oil.
- 25 The oil was dissolved in 1.5 mL of TFA under an atmosphere of dry N₂. The reaction mixture was stirred at ambient temperature for 10 minutes after which time it was concentrated under vacuum and the resulting residue was partitioned between DCM and aqueous 0.1 N NaOH.

- 5 The DCM layer was then washed with basic brine (pH = 10), dried over Na₂SO₄, filtered and concentrated under vacuum to give 118 mg of the title compound as a yellow solid **32**.

C.I. m/z 414 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.37 (d, J = 8.9 Hz, 1 H), 8.27 (d, J = 8.7 Hz, 1 H), 7.65 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.30 (d, J = 2.5 Hz, 1 H), 7.24 (m, 1 H), 7.09 (dd, J = 2.5, 8.9 Hz, 1 H), 3.87 (m, 4 H), 2.87 (m, 3 H), 2.03 (m, 2 H), 1.81 (m, 2 H), 1.56 (brs, 2 H), 1.32 (m, 1 H), 0.66 (m, 2 H), 0.39 (m, 2 H).

Example 33

1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.



- 15 The same procedure that was used for example 1 was followed except that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F to give the title compound **33**.


C.I. m/z 418 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.40 (d, J = 8.7 Hz, 1 H), 8.29 (d, J = 8.7 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.45 (m, 2 H), 7.35 (d, J = 2.5 Hz, 1 H), 7.25 (m, 1 H), 7.13 (dd, J = 2.5, 8.7 Hz, 1 H), 4.22 (m, 2 H), 3.88 (m, 2 H), 3.81 (m, 2 H), 3.48 (s, 3 H), 2.89 (m, 3 H), 2.05 (m, 2 H), 1.82 (m, 2 H), 1.57 (brs, 2 H).

25

Example 33A

1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine p-toluenesulphonate salt.

-70-




5 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **33** (15.13 g, 36.2 mMol) was dissolved in 93 mL of EtOH and the reaction was heated to reflux under an atmosphere of dry N₂. To this solution was added p-toluenesulphonic acid mono hydrate (6.89 g, 36.2 mMol) which was then stirred at reflux overnight. The reaction mixture was then
 10 cooled to ambient temperature and filtered. The filtercake was washed with EtOH, dried under vacuum to give 18.46 g of the title compound **33A** as an off-white solid.

¹H NMR (CD₃OD) δ 8.93 (s, 1 H), 8.57 (d, J = 9.1 Hz, 1 H), 8.40 (d, J = 8.7 Hz, 1 H), 7.89 (d, J = 8.7 Hz, 1 H), 7.68 (d, J = 6.2 Hz, 2 H), 7.58 (m, 1 H), 7.49 (m, 1 H), 7.33 (dd, J = 1.2, 7.9 Hz, 1 H), 7.25 (d, J = 2.5 Hz, 1 H), 7.17 (m, 3 H), 4.18 (m, 2 H), 3.93 (m, 2 H), 3.77 (m, 2 H), 3.43
 15 (s, 3 H), 3.30 (m, 1 H), 2.86 (m, 2 H), 2.31 (s, 3 H), 2.16 (m, 2 H), 2.06 (m, 2 H).

Example 34

1-{2-[5-(Pyridin-3-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.



20


The same procedure that was used for example 32 was followed except 3-picolyl chloride hydrochloride was used in the place of cyclopropyl methane bromide in example 32 and the equivalents of Cs₂CO₃ was doubled to give the title compound **34**.

C.I. m/z 451[M+1]; ¹H NMR (CDCl₃) δ 8.73 (d, J = 2.3 Hz, 1 H), 8.65 (s, 1 H), 8.58 (dd, J = 1.4, 5.0 Hz, 1 H), 8.42 (d, J = 8.9 Hz, 1 H), 8.29 (d, J = 8.7 Hz, 1 H), 7.82 (m, 1 H), 7.66 (d, J = 8.7
 25

- 5 Hz, 1 H), 7.46 (m, 2 H), 7.41 (d, J = 2.5 Hz, 1 H), 7.31 (m, 1 H), 7.26 (m, 1 H), 7.13 (m, 1 H), 5.17 (s, 2 H), 3.88 (m, 2 H), 2.91 (m, 3 H), 2.03 (m, 2 H), 1.80 (m, 2 H), 1.61 (brs, 2 H).

Example 35

1-[2-(5-Benzoyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.




10

The same procedure that was used for example 32 was followed except benzyl bromide was used in the place of cyclopropyl methane bromide to give the title compound 35.

- C.I. m/z 450[M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.42 (d, J = 8.9 Hz, 1 H), 8.30 (d, J = 8.7 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.37-7.51 (m 7 H), 7.33 (d, J = 7.3 Hz, 1 H), 7.26 (m, 1 H),
 15 7.18 (dd, J = 2.5, 8.9 Hz, 1 H), 5.16 (s, 2 H), 3.88 (m, 2 H), 2.90 (m, 3 H), 2.04 (m, 2 H), 1.81 (m, 2 H), 1.40 (brs, 2 H).

Example 36

1-[2-[5-(Pyridin-4-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine.



20

The same procedure that was used for example 32 was followed except the hydrochloride salt of 4-picolyl chloride was used in the place of cyclopropyl methane bromide and the number of equivalents of Cs₂CO₃ was doubled to give the title compound 36.


- C.I. m/z 451[M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.63 (m, 2 H), 8.44 (d, J = 9.1 Hz, 1 H),
 25 8.31 (d, J = 8.9 Hz, 1 H), 7.68 (d, J = 8.7 Hz, 1 H), 7.47 (m, 2 H), 7.41 (d, J = 6.0 Hz, 2 H),

-72-

- 5 7.37 (d, J = 2.3 Hz, 1 H), 7.26 (m, 1 H), 7.16 (dd, J = 2.3, 9.1 Hz, 1 H), 5.19 (s, 2 H), 3.90 (m, 2 H), 2.90 (m, 3 H), 2.05 (m, 2 H), 1.80 (m, 2 H), 1.55 (brs, 2 H).

Example 37

1-{2-[5-(2-Dimethylamino-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.




10

The same procedure that was used for example 32 was followed except that the hydrochloride salt of 2-dimethylamino ethyl chloride was used in the place of cyclopropyl methane bromide and the number of equivalents of Cs₂CO₃ was doubled to give the title compound 37.

- C.I. m/z 431 [M+1]; ¹H NMR (CDCl₃) δ 8.61 (s, 1 H), 8.45 (d, J = 8.9 Hz, 1 H), 8.27 (d, J = 8.7 Hz, 1 H), 7.62 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.31 (m, 1 H), 7.24 (m, 1 H), 7.12 (m, 1 H), 4.12 (m, 2 H), 3.72-3.88 (m, 4 H), 2.90 (m, 3 H), 1.78-2.10 (m, 12 H).
- 15

Example 38

1-{2-[5-(3-Amino-propoxy)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.



20

Example 38A

[1-(2-{5-[3-(1,3-Dioxo-1,3-dihydro-isindol-2-yl)-propoxy]-benzimidazol-1-yl]-quinolin-8-yl})-piperidin-4-yl]-carbamic acid tert-butyl ester.

25

- 5 The same procedure that was used for example 32 was followed except N-(3-bromopropyl)-phthalimide was used in the place of cyclopropyl methane bromide and, the tert-butyl carbamate ester intermediate was not subjected to TFA mediated cleavage, to give the title compound **38A** as a yellow solid.

Example 38B

- 10 (1-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-carbamic acid tert-butyl ester.

- [1-(2-{5-[3-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-propoxy]-benzoimidazol-1-yl}-quinolin-8-yl)-piperidin-4-yl]-carbamic acid tert-butyl ester **38A** (200 mg, 0.31 mMol) was dissolved in 1 mL of EtOH under an atmosphere of dry N₂. To this solution was added 50 µL of hydrazine and the resulting solution was stirred at ambient temperature for 3 hours. The reaction mixture was then filtered, concentrated under vacuum and purified via silica gel chromatography eluting with DCM/MeOH/NH₄OH (89/10/1) to give 120 mg of the title compound **38B** as a yellow film.

20

Example 38C


1-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.

- (1-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-carbamic acid tert-butyl ester **38B** (175mg, 0.42 mMol) was dissolved in a solution of 0.5 mL of TFA and 0.5 mL of DCM under an atmosphere of dry N₂ and was stirred at ambient temperature for 1 hour. The reaction mixture was then concentrated under vacuum and subsequently partitioned between DCM and aqueous 1 N NaOH. The DCM layer was then dried over MgSO₄, filtered and concentrated to give the title compound **38** as a light green solid.
- 30 C.I. m/z 417 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.36 (d, J = 8.7 Hz, 1 H), 8.26 (d, J = 8.7 Hz, 1 H), 7.64 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.31 (d, J = 2.5 Hz, 1 H), 7.23 (m, 1 H), 7.04 (m, 1 H), 4.12 (t, J = 6.2 Hz, 2 H), 3.87 (m, 2 H), 2.89 (m, 3 H), 2.03 (m, 2 H), 1.94 (m, 2 H), 1.77 (m, 2 H), 1.54 (brs, 4 H).

35

Example 39

1-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.



5

(1-[2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-carbamic acid tert-butyl ester **38B** (120 mg, 0.288 mMol) was dissolved in a cooled (0°C) solution of 1 mL of acetonitrile (ACN) and 0.5 mL of formaldehyde (37% wt in water) under an atmosphere of dry N₂. To this solution was added NaCNBH₃ (72 mg, 1.2 mMol) and the solution was stirred for 10 30 minutes after which time 200 µL of AcOH was added. The reaction mixture was subsequently warmed to ambient temperature and stirred overnight. The reaction mixture was then concentrated under vacuum and partitioned between aqueous 0.1 N NaOH and DCM. The aqueous layer was washed 3 more times with DCM. The DCM extracts were combined, dried over Na₂SO₄, filtered and concentrated under vacuum to give a green film.


15 The film was purified using flash silica gel chromatography eluting with DCM/MeOH/NH₄OH (89.9/10/0.1) to give 70 mg of the tert-butylcarbonyl protected product. This residue was dissolved in a solution of 1 mL of TFA and 1 mL of DCM under an atmosphere of dry N₂. The reaction mixture was stirred for 1 hour after which time it was concentrated under vacuum, partitioned between DCM and aqueous 1 N NaOH. The aqueous layer was washed 20 2 more times with DCM. The DCM extracts were combined and dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow film. The film was purified using flash silica gel chromatography eluting with DCM/MeOH/NH₄OH (84.9/15/0.1) to give 30 mg of the title compound **39** as a yellow film.

C.I. m/z 445 [M+1]; ¹H NMR (CD₃OD) δ 8.87 (s, 1 H), 8.58 (d, J = 8.8 Hz, 1 H), 8.31 (dd, J = 2.0, 8.7 Hz, 1 H), 7.80 (dd J = 2.0, 8.7 Hz, 1 H), 7.48 (m, 1 H), 7.43 (m, 1 H), 7.25 (m, 1 H), 7.17 (d, J = 2.1 Hz, 1 H), 7.10 (m, 1 H), 4.02 (m, 2 H), 3.78 (m, 2 H), 2.71-2.84 (m, 3 H), 2.53 (m, 2 H), 2.27 (s, 6 H), 2.03 (m, 2 H), 1.99 (m, 4 H), 1.79 (m, 2 H).

Example 40

30 1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidine-4-carboxylic acid ethyl ester.

-75-



5


The same procedure that was used in example 1 was followed except that ethyl isonipecotate was used in the place of the piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound **40** as a yellow solid.

C.I. m/z 431 [M+1]; ¹H NMR (CDCl₃) δ 8.64 (s, 1 H), 8.40 (d, J = 8.9 Hz, 1 H), 8.28 (d, J = 8.9 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.33 (m, 1 H), 7.23 (m, 1 H), 7.08 (dd, J = 2.5, 8.9 Hz, 1 H), 4.18 (q, J = 7.5, 2 H), 3.89 (s, 3 H), 3.85 (m, 2 H), 2.90 (m, 2 H), 2.52 (m, 1 H), 2.14 (m, 4 H), 1.29 (t, J = 7.5 Hz, 3 H).

10

Example 41

1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidine-4-carboxylic acid.



15

1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidine-4-carboxylic acid ethyl ester **40** (28 mg, 0.065 mMol) was dissolved in 1 mL of ethanol (EtOH). To this solution was added 1 mL of 1 N aqueous NaOH and the mixture was subsequently heated to 60°C and reacted at this temperature overnight. The reaction mixture was then cooled to ambient temperature and partitioned between 10% aqueous citric acid and ethyl acetate. The organic layer was washed again with 10% aqueous citric acid then three more times with water. The ethyl acetate layer was dried over Na₂SO₄, filtered and concentrated to give 7 mg of the title compound **41** as a yellow solid.

20


C.I. m/z 403 [M+1]; ¹H NMR (CD₃OD) δ 8.94 (s, 1 H), 8.73 (d, J = 9.2 Hz, 1 H), 8.40 (d, J = 8.9 Hz, 1 H), 7.90 (d, J = 8.9 Hz, 1 H), 7.55 (m, 1 H), 7.47 (m, 1 H), 7.31 (m, 1 H), 7.23 (d, J = 2.3 Hz, 1 H), 7.07 (dd, J = 2.4, 9.0 Hz, 1 H), 3.87 (s, 3 H), 3.79 (m, 2 H), 2.85 (m, 2 H), 2.50 (m, 1 H), 2.08 (m, 4 H).

25

5

Example 42

4-Dimethylaminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol.

Example 42A

10

5-(2-Methoxy-ethoxy)-2-nitro-phenylamine.

4-Amino-3-nitrophenol (58.53 g, 379.7 mMol) was dissolved in 600 mL of anhydrous DMF under an atmosphere of dry N₂ and the solution was mechanically stirred. The reaction mixture was then cooled to 0°C and to this mixture was added Cs₂CO₃ (177.4 g, 455.7 mMol), sodium iodide (5.7 g, 37.9 mMol) and 2-bromoethyl methyl ether (39.3 mL, 417.7 mMol). After stirring for 15 minutes, the reaction mixture was then warmed up to ambient temperature and then stirred overnight. The reaction mixture is then poured into 6 L of water. The precipitate was collected via suction filtration. The wet precipitate was taken up in toluene and then concentrated under vacuum to remove the water. Finally, the solid was recrystallized from 2-propanol to give 57.93 g of 5-(2-Methoxy-ethoxy)-2-nitro-phenylamine **42A** as an orange solid. A second crop of 12.5 g of compound **42A** was also obtained.

Example 42B

25


4-Dimethylaminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol.

The same procedure that was used in example 24 was followed except that 5-(2-Methoxy-ethoxy)-2-nitro-phenylamine **42A** was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **42** as a yellow solid.

C.I. m/z 476 [M+1]; ¹H NMR (CD₃OD) δ 8.85 (s, 1 H), 8.55 (d, J = 9.1 Hz, 1 H), 8.29 (d, J = 8.7 Hz, 1 H), 7.76 (d, J = 8.7 Hz, 1 H), 7.43 (m, 2 H), 7.27 (d, J = 7.5 Hz, 1 H), 7.21 (d, J = 2.5 Hz, 1 H), 7.00 (dd, J = 2.5, 9.0 Hz, 1 H), 4.13 (m, 2 H), 3.77 (m, 2 H), 3.44 (s, 3 H), 3.32 (m, 2 H), 3.16 (m, 2 H), 2.42 (s, 2 H), 2.38 (s, 6 H), 1.97 (m, 2 H), 1.75 (m, 2 H).

Example 43


- 5 N-{1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-acetamide.



- 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** (100 mg, 0.268 mMol) and sodium cyanate (35 mg, 0.536 mMol) were dissolved in 1 mL of AcOH under an atmosphere of dry N₂. The reaction mixture was stirred overnight at ambient temperature.
- 10 The reaction mixture was then concentrated under vacuum and partitioned between DCM and saturated aqueous NaHCO₃. The organic layer was dried over Na₂SO₄, filtered and concentrated to a yellow foam which was chromatographed on flash silica gel eluting with MeOH/NH₄OH/DCM (2/0.2/97.8) to give the title compound **43**.
- C.I. m/z 416 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.30 (m, 2 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.43 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.23 (dd, J = 1.7, 7.1 Hz, 1 H), 7.03 (dd, J = 2.5, 9.0 Hz, 1 H), 5.64 (brd, J = 8.3 Hz, 1 H), 4.03 (m, 1 H), 3.89 (s, 3 H), 3.84 (m, 2 H), 2.93 (m, 2 H), 2.16 (m, 2 H), 2.02 (s, 3 H), 1.88 (m, 2 H).
- 15

Example 44

- 20 N-{1-[2-(5-Cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-acetamide.




- The procedure used in example 43 was followed except that 1-[2-(5-cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **32** was used in the place of 1-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** to give the title compound **44** as a yellow solid.
- 25 C.I. m/z 456 [M+1]; ¹H NMR (CDCl₃) δ 8.61 (s, 1 H), 8.33 (d, J = 8.7 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.65 (d, J = 8.7 Hz, 1 H), 7.45 (m, 2 H), 7.28 (d, J = 2.1 Hz, 1 H), 7.22 (m, 1 H), 7.07 (dd, J = 1.5, 8.7 Hz, 1 H), 5.63 (brd, J = 8.3 Hz, 1 H), 4.03 (m, 1 H), 3.88 (d, J = 7.0 Hz, 2 H),

-78-

- 5 3.85 (m, 2 H), 2.92 (m, 2 H), 2.14 (m, 2 H), 2.04 (s, 3 H), 1.88 (m, 2 H), 1.33 (m, 1 H), 0.67 (m, 2 H), 0.38 (m, 2 H).

Example 45

1-[2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ol.



10


The same procedure that was used for example 19 was followed except that 5-(2-methoxy-ethoxy)-2-nitro-phenylamine **42A** was used in the place of was used in the place of 4-methoxy-2-nitroaniline nitroaniline in example 1B to give the title compound **45** as a yellow solid.

- 15 C.I. m/z 419 [M+1]; ¹H NMR (CDCl₃) δ 8.62 (s, 1 H), 8.39 (d, J = 9.1 Hz, 1 H), 8.24 (d, J = 8.7 Hz, 1 H), 7.62 (d, J = 8.7 Hz, 1 H), 7.43 (m, 2 H), 7.31 (d, J = 2.1 Hz, 1 H), 7.23 (m, 1 H), 7.10 (m, 1 H), 4.17 (m, 2 H), 3.93 (m, 1 H), 3.77 (m, 2 H), 3.71 (m, 2 H), 3.44 (s, 3 H), 3.04 (m, 2 H), 2.47 (brd, J = 3.3 Hz, 1 H), 2.14 (m, 2 H), 1.97 (m, 2 H).

20

Example 46

{1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-urea.



- 1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** (150 mg, 0.402 mMol) and 1,1'-carbonyldiimidazole (78 mg, 0.48 mMol) were dissolved in 1 mL of anhydrous THF under an atmosphere of dry N₂. The reaction mixture was stirred for 2 hours at ambient temperature. 1.0 mL of concentrated ammonium hydroxide (NH₄OH) was then added to the reaction mixture that was then stirred overnight. The reaction mixture was then partitioned between DCM and water. The DCM layer was dried over Na₂SO₄, filtered and concentrated
- 25


-79-

5 under vacuum. The residue was chromatographed over flash silica gel eluting with MeOH/DCM (5/95) to give 15 mg of the title compound **46** as a white solid.

C.I. m/z 417 [M+1]; ¹H NMR (CD₃OD) δ 8.89 (s, 1 H), 8.59 (d, J = 8.9 Hz, 1 H), 8.36 (J = 8.9 Hz, 2 H), 7.85 (d, J = 8.7 Hz, 1 H), 7.74 (brs, 1 H), 7.53 (d, J = 8.1 Hz, 1 H), 7.45 (m, 1 H), 7.30 (d, J = 7.5 Hz, 1 H), 7.23 (m, 1 H), 7.05 (m, 1 H), 3.86 (s, 3 H), 3.73 (m, 2 H), 3.66 (m, 1 H), 2.87 (m, 2 H), 2.08 (m, 2 H), 1.83 (m, 2 H).

Example 47

4-Aminomethyl-1-{2-[5-(pyridin-2-ylmethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol.



15

Example 47A

2-Nitro-4-(pyridin-2-ylmethoxy)-phenylamine.

4-Amino-3-nitrophenol (5.00 g, 32.4 mMol), Cs₂CO₃ (22.8 g, 70 mMol), sodium iodide (476 mg, 3.20 mMol) and 2-picoyl chloride hydrochloride (11.2 g, 35 mMol) were added to 20 mL of anhydrous DMF under an atmosphere of dry N₂. The reaction mixture was stirred overnight at ambient temperature. The reaction mixture is then poured into water. The precipitate was collected via suction filtration. The precipitate was partitioned between DCM and 1.0 N aqueous NaOH. The organic layer was washed 2 more times with 1 N aqueous NaOH to remove any unreacted phenol. The organic layer was then washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give 7.7 g of a dark red solid. The solid was chromatographed on flash silica gel eluting with EtOAc/DCM (20/80) to give 3.7 g of the title compound **47A** as an orange solid.

30

Example 47B

4-Aminomethyl-1-{2-[5-(pyridin-2-ylmethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol.


The same procedure that was used in example 26 was followed except that 2-nitro-4-(pyridin-2-ylmethoxy)-phenylamine **47A** was used in the place of 4-methoxy-2-nitroaniline in example 35 1B to give the title compound **47**.

- 5 C.I. m/z 481 [M+1]; ¹H NMR (CDCl₃) δ 8.64 (s, 1 H), 8.60 (dd, J = 0.8, 5.0 Hz, 1 H), 8.31 (d, J = 9.1 Hz, 1 H), 8.27 (d, J = 8.7 Hz, 1 H), 7.68 (m, 1 H), 7.63 (d, J = 9.1 Hz, 1 H), 7.55 (d, J = 7.9 Hz, 1 H), 7.44 (m, 2 H), 7.38 (d, J = 2.1 Hz, 1 H), 7.27 (m, 1 H), 7.24 (m, 1 H), 7.12 (m, 1 H), 5.27 (s, 2 H), 3.67 (m, 2 H), 3.22 (m, 2 H), 2.78 (s, 2 H), 2.40 (brs, 3 H), 1.96 (m, 2 H), 1.81 (m, 2 H).

10

Example 48

Cyclopropyl-(1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-amine.




- 15 The same procedure that was used in example 20 was followed except that 5-(2-methoxy-ethoxy)-2-nitro-phenylamine **42A** was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **48** as a yellow solid.

- C.I. m/z 458 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.40 (d, J = 8.7 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.45 (m, 2 H), 7.35 (d, J = 2.5 Hz, 1 H), 7.25 (m, 1 H), 7.15 (m, 1 H), 4.22 (m, 2 H), 3.82 (m, 2 H), 3.80 (m, 2 H), 3.47 (s, 3 H), 2.88 (m, 3 H), 2.27 (m, 1 H), 2.18 (m, 2 H), 1.85 (m, 2 H), 0.50 (m, 2 H), 0.47 (m, 2 H).

20

Example 49

(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-dimethyl-amine.



25


The same procedure that was used in example 4 was followed except that 5-(2-methoxy-ethoxy)-2-nitro-phenylamine **42A** was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **49**.

- 5 C.I. m/z 446 [M+1]; ¹H NMR (CDCl₃) δ 8.59 (s, 1 H), 8.35 (d, J = 8.7 Hz, 1 H), 8.18 (d, J = 8.7 Hz, 1 H), 7.57 (d, J = 8.7 Hz, 1 H), 7.38 (m, 2 H), 7.28 (d, J = 2.5 Hz, 1 H), 7.15 (d, J = 2.5, 6.2 Hz, 1 H), 7.06 (m, 1 H), 4.18 (m, 2 H), 3.94 (m, 2 H), 3.75 (m, 2 H), 3.42 (s, 3 H), 2.72 (m, 2 H), 2.40 (m, 1 H), 2.34 (s, 6 H), 1.95 (m, 4 H).

10

Example 50

(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methyl-amine.



- The same procedure that was used in example 22 was followed except that 5-(2-methoxy-ethoxy)-2-nitro-phenylamine **42A** was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **50**.


15

C.I. m/z 432 [M+1]; ¹H NMR (CDCl₃) δ 8.62 (s, 1 H), 8.37 (d, J = 9.1 Hz, 1 H), 8.23 (d, J = 8.7 Hz, 1 H), 7.61 (d, J = 8.7 Hz, 1 H), 7.42 (m, 2 H), 7.31 (d, J = 2.1 Hz, 1 H), 7.20 (d, J = 3.0, 5.8 Hz, 1 H), 7.11 (dd, J = 2.5, 9.1 Hz, 1 H), 4.18 (m, 2 H), 3.88 (m, 2 H), 3.76 (m, 2 H), 3.45 (s, 3 H), 2.85 (m, 2 H), 2.59 (m, 1 H), 2.47 (s, 3 H), 2.18 (brs, 1 H), 2.11 (m, 2 H), 1.75 (m, 2 H).

20

Example 51

(1-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-dimethyl-amine.



25


The same procedure that was used in example 4 was followed except that 1-{2-[5-(3-dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **39** was used in the place of 1-{2-[5-methoxy-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **1** to give the title compound **51**.

- 5 C.I. m/z 473 [M+1]; ¹H NMR (CD₃OD) δ 8.87 (s, 1 H), 8.61 (d, J = 9.1 Hz, 1 H), 8.31 (d, J = 8.7 Hz, 1 H), 7.78 (d, J = 8.7 Hz, 1 H), 7.48 (d, J = 6.4 Hz, 1 H), 7.42 (m, 1 H), 7.21 (m, 2 H), 7.05 (m, 1 H), 4.05 (m, 2 H), 3.82 (m, 2 H), 3.19 (m, 1 H), 2.65 (m, 5 H), 2.40 (m, 1 H), 2.34 (s, 6 H), 1.95 (m, 4 H).

10

Example 52

{1-[2-(5-Cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine.

Example 52A

15

4-Cyclopropylmethoxy-2-nitro-phenylamine.

- 4-Amino-3-nitrophenol (13.3 g, 84.8 mMol), Cs₂CO₃ (33.2 g, 102 mMol) and cyclopropylmethyl bromide (9.1 mL, 93.3 mMol) were added to 30 mL of anhydrous DMF under an atmosphere of dry N₂. The reaction mixture was stirred overnight at ambient temperature. The reaction mixture is then poured into water. The precipitate was collected via suction filtration. The precipitate was partitioned between DCM and 1.0 N aqueous NaOH. The organic layer was washed 2 more times with 1 N aqueous NaOH to remove any unreacted phenol. The organic layer was then washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give 18.6 g of the title compound **52A** as an orange solid.

25

Example 52B

{1-[2-(5-Cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine.


- The same procedure that was used in example 22 was followed except that 4-cyclopropylmethoxy-2-nitro-phenylamine **52A** was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **52**.

C.I. m/z 428 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.38 (d, J = 9.1 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.65 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.30 (d, J = 2.5 Hz, 1 H), 7.24 (m, 1 H), 7.06

- 5 (m, 1 H), 3.89 (m, 4 H), 2.90 (m, 2 H), 2.71 (m, 1 H), 2.55 (s, 3 H), 2.16 (m, 2 H), 2.00 (brs, 1 H), 1.89 (m, 2 H), 1.30 (m, 1 H), 0.64 (m, 2 H), 0.38 (m, 2 H).

Example 53

{1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethyl-amine.



10

The same procedure that was used in example 4 was followed except that {1-[2-(5-cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine **52** was used in the place of 1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** to give the title compound **53**.


15

C.I. m/z 442 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.38 (d, J = 9.0 Hz, 1 H), 8.30 (d, J = 9.0 Hz, 1 H), 7.69 (d, J = 8.7 Hz, 1 H), 7.48 (m, 2 H), 7.31 (d, J = 2.3 Hz, 1 H), 7.24 (m, 1 H), 7.09 (d, J = 2.3, 9.0 Hz, 1 H), 4.03 (m, 2 H), 3.89 (d, J = 6.8 Hz, 2 H), 2.83 (m, 2 H), 2.55 (m, 1 H), 2.44 (s, 6 H), 2.06 (m, 4 H), 1.30 (m, 1 H), 0.68 (m, 2 H), 0.39 (m, 2 H).

20

Example 54

2-Amino-N-(1-[2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-acetamide.




- 25 N-(tert-butoxycarbonyl)glycine (105 mg, 0.600 mMol) was dissolved in 3 mL of DCM under an atmosphere of dry N₂. To this mixture was added 1,1'-carbonyldiimidazole (100 mg, 0.61 mMol) and the reaction mixture was stirred at ambient temperature for 30 minutes. To this solution was then added 4-dimethylaminopyridine (8.0 mg, 0.065 mMol) and 1-[2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine **33** (170 mg, 0.408

5 mMol). The reaction mixture was subsequently stirred overnight at ambient temperature. The reaction mixture was then partitioned between DCM and saturated aqueous NaHCO_3 . The organic layer was saved and washed again with saturated aqueous NaHCO_3 , dried over Na_2SO_4 , filtered and concentrated under vacuum to give a yellow foam. The residue was chromatographed on flash silica gel eluting with MeOH/DCM (3/97) to give 182 mg of a yellow
10 residue. The residue was dissolved in 0.5 ml of TFA and stirred at ambient temperature for 30 minutes under an atmosphere of dry N_2 . The reaction mixture was subsequently concentrated under vacuum and then partitioned between 0.1 N aqueous NaOH. The organic layer was saved and washed again with 0.1 N aqueous NaOH, dried over Na_2SO_4 , filtered and concentrated under vacuum to give a yellow foam. The foam was chromatographed on flash
15 silica gel eluting with MeOH/DCM (5/95) to give 75 mg of the title compound as **54** a yellow solid.

C.I. m/z 475 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.62 (s, 1 H), 8.31 (d, $J = 8.7$ Hz, 1 H), 8.27 (d, $J = 8.7$ Hz, 1 H), 7.64 (d, $J = 8.7$ Hz, 1 H), 7.44 (m, 2 H), 7.38 (brd, $J = 7.9$ Hz, 1 H), 7.32 (d, $J = 2.1$ Hz, 1 H), 7.22 (m, 1 H), 7.06 (d, $J = 2.5, 9.2$ Hz, 1 H), 4.20 (m, 2 H), 4.02 (m, 1 H), 3.86 (m, 2
20 H), 3.78 (m, 2 H), 3.46 (s, 3 H), 3.37 (s, 2 H), 2.94 (m, 2 H), 2.13 (m, 2 H), 1.90 (m, 2 H), 1.74 (brs, 2 H).

Example 55

25 -(S)-2-Amino-N-(1-{2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-propionamide.




The same procedure that was used in example 54 was followed except that N-(tert-butoxycarbonyl)-L-alanine was used in the place of N-(tert-butoxycarbonyl)glycine to give the title compound **55** as a yellow solid.

30 C.I. m/z 489 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.61 (s, 1 H), 8.27 (d, $J = 8.7$ Hz, 1 H), 8.25 (d, $J = 9.1$ Hz, 1 H), 7.63 (d, $J = 8.7$ Hz, 1 H), 7.42 (m, 3 H), 7.30 (d, $J = 2.5$ Hz, 1 H), 7.22 (dd, $J = 2.1, 8.7$ Hz, 1 H), 7.07 (d, $J = 2.5, 8.7$ Hz, 1 H), 4.17 (m, 2 H), 3.95 (m, 1 H), 3.85 (m, 2 H), 3.77 (m, 2 H), 3.50 (m, 1 H), 3.43 (s, 3 H), 2.91 (m, 2 H), 2.10 (m, 2 H), 1.85 (m, 4 H), 1.34 (d, $J = 6.7$ Hz, 3 H).

5

Example 56

-(R)-2-Amino-N-(1-(2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-yl)-propionamide.




- 10 The same procedure that was used in example 54 was followed except that N-(tert-butoxycarbonyl)-D-alanine was used in the place of N-(tert-butoxycarbonyl)glycine to give the title compound **56** as a yellow solid.

C.I. m/z 489 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.26 (m, 2 H), 7.65 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.38 (brd, J = 8.7 Hz, 1 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.22 (dd, J = 2.1, 7.0 Hz, 1 H), 7.10 (d, J = 2.5, 8.7 Hz, 1 H), 4.20 (m, 2 H), 3.99 (m, 1 H), 3.86 (m, 2 H), 3.79 (m, 2 H), 3.50 (m, 1 H), 3.46 (s, 3 H), 2.96 (m, 2 H), 2.12 (m, 2 H), 1.88 (m, 2 H), 1.52 (brs, 2 H), 1.35 (d, J = 7.1 Hz, 3 H).

Example 57

- 20 2-Amino-N-(1-(2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-yl)-isobutyramide.




- N-(tert-butoxycarbonyl)glycine (73 mg, 0.36 mMol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (69 mg, 0.36 mMol) were dissolved in 1 mL of anhydrous
 25 DMF under an atmosphere of dry N₂ and stirred at ambient temperature for 30 minutes. Triethylamine (167 μL, 1.24 mMol) and 1-(2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine **33** (146 mg, 0.35 mMol) were then added to the reaction mixture which was subsequently stirred overnight. The reaction mixture was then partitioned

5 between DCM and saturated aqueous NaHCO_3 . The organic layer was saved and washed two more times with saturated aqueous NaHCO_3 , dried over Na_2SO_4 , filtered and concentrated under vacuum to give a yellow oil. The residue was chromatographed on flash silica gel eluting with a gradient of DCM to MeOH/DCM (2/98) to give 30 mg of a yellow residue. The residue was dissolved in 0.25 ml of TFA and stirred at ambient temperature for
10 30 minutes under an atmosphere of dry N_2 . The reaction mixture was subsequently concentrated under vacuum and then partitioned between 0.1 N aqueous NaOH . The organic layer was saved and washed again with 0.1 N aqueous NaOH , dried over Na_2SO_4 , filtered and concentrated under vacuum to give the title compound **57** as a yellow residue.

C.I. m/z 503 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.70 (s, 1 H), 8.29 (d, $J = 8.7$ Hz, 1 H), 8.21 (d, $J = 9.1$
15 Hz, 1 H), 7.66 (m, 2 H), 7.44 (m, 2 H), 7.25 (d, $J = 2.5$ Hz, 1 H), 7.21 (dd, $J = 2.6, 9.1$ Hz, 1 H), 7.12 (m, 1 H), 4.19 (m, 2 H), 3.95 (m, 1 H), 3.88 (m, 2 H), 3.79 (m, 2 H), 3.46 (s, 3 H), 2.93 (m, 2 H), 2.09 (m, 2 H), 1.90 (m, 2 H), 1.49 (s, 6 H).

Example 58

20 1-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamino)-2-methyl-propan-2-ol.



25

Example 58A

(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamino)-acetic acid ethyl ester.

Bromo-acetic acid ethyl ester (80 μL , 0.710 mMol), diisopropylethylamine (180 μL , 1.00 mMol)
30 and 1-(2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine **33** (270 mg, 0.647 mMol) were dissolved in 3 mL of anhydrous DCM under an atmosphere of dry N_2 and stirred overnight at ambient temperature. The reaction mixture was then partitioned between DCM and 0.1 N aqueous NaOH . The organic layer was saved and washed sequentially with 0.1 N aqueous NaOH and brine, dried over Na_2SO_4 , filtered and

- 5 concentrated under vacuum to give a yellow residue. The residue was chromatographed on flash silica gel eluting with MeOH/DCM (2/98) to give 220 mg of the title compound **58A**.

Example 58B


10 1-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamino)-2-methyl-propan-2-ol.

(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamino)-acetic acid ethyl ester (110 mg, 0.218 mMol) **58A** was dissolved in 2 mL of anhydrous THF under an atmosphere of dry N₂. The reaction mixture was cooled to -78°C and then 260 µL of a
15 solution of 1.0 M methylmagnesium bromide in THF was added. The reaction mixture was slowly warmed up to ambient temperature and stirred overnight. The reaction mixture was then partitioned between DCM and saturated aqueous NaHCO₃. The organic layer was saved and washed again with saturated aqueous NaHCO₃, dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow residue. The residue was chromatographed on
20 flash silica gel eluting with MeOH/DCM/NH₄OH (2/97.9/0.1) to give 20 mg of the title compound **58** as a yellow residue.

C.I. m/z 490 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.35 (d, J = 9.1 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.22 (m, 1 H), 7.12 (m, 1 H), 4.21 (m, 2 H), 3.88 (m, 2 H), 3.80 (m, 2 H), 3.47(s, 3 H), 2.88 (m, 2 H), 2.67 (m, 1 H),
25 2.62 (s, 2 H), 2.13 (m, 2 H), 1.78 (m, 2 H), 1.19 (s, 6 H).

Example 59

(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-pyridin-2-ylmethyl-amine.




30 1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **33** (50 mg, 0.119 mMol) and pyridine-2-carbaldehyde (11 µL, 0.119 mMol) were dissolved in a solution of 2 mL of EtOH and 500 µL of DCE under an atmosphere of dry N₂. The reaction mixture was subsequently heated to reflux and stirred at this temperature for 1.5 hours. The reaction
35 mixture was then cooled to ambient temperature and NaBH₄ (14 mg, 0.357 mMol) was added.

- 5 The reaction mixture was stirred overnight at ambient temperature. The solvent was removed under vacuum and the resulting residue was partitioned between DCM and 0.1 N aqueous NaOH. The DCM layer was dried over Na₂SO₄, filtered and concentrated under vacuum to give 47 mg of the title compound **59** as a yellow residue.

C.I. m/z 509 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.55 (m, 1 H), 8.39 (d, J = 9.1 Hz, 1 H),
 10 8.26 (d, J = 8.7 Hz, 1 H), 7.65 (m, 2 H), 7.44 (m, 2 H), 7.36 (d, J = 8.9 Hz, 1 H), 7.32 (d, J = 2.5 Hz, 1 H), 7.22 (m, 1 H), 7.12 (m, 2 H), 4.20 (m, 2 H), 4.01 (s, 2 H), 3.89 (m, 2 H), 3.79 (m, 2 H), 3.46 (s, 3 H), 2.90 (m, 2 H), 2.77 (m, 1 H), 2.13 (m, 2 H), 1.88 (m, 3 H).

Example 60

- 15 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-pyridin-3-ylmethyl-amine.




- The same procedure that was used in example 59 was followed except that pyridine-3-carbaldehyde was used in the place of pyridine-2-carbaldehyde to give the title compound **60**
 20 as a yellow residue.

C.I. m/z 509 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (s, 1 H), 8.59 (d, J = 2.7 Hz, 1 H), 8.50 (dd, J = 2.7, 5.0, 1 H), 8.38 (d, J = 8.7 Hz, 1 H), 8.29 (d, J = 8.7 Hz, 1 H), 7.76 (m, 1 H), 7.67 (m, 1 H), 7.45 (m, 2 H), 7.34 (d, J = 2.5 Hz, 1 H), 7.27 (m, 1 H), 7.22 (m, 1 H), 7.12 (dd, J = 2.5, 8.7, 1 H), 4.22 (m, 2 H), 3.91 (s, 2 H), 3.88 (m, 2 H), 3.80 (m, 2 H), 3.47 (s, 3 H), 2.90 (m, 2 H), 2.75
 25 (m, 1 H), 2.14 (m, 2 H), 1.85 (m, 2 H), 1.75 (brs, 1 H).

Example 61

4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol.



Example 61A

N-(2-Bromo-phenyl)-3-phenyl-acrylamide.

- 10 2-Bromoaniline (48.41 g, 281.4 mMol) was dissolved in 500 mL of anhydrous DCM under an atmosphere of dry N₂. Pyridine (45.5 mL, 563 mMol) was then added and the reaction was subsequently cooled to 0°C after which time cinnamyl chloride (46.9 g, 281.4 mMol) was added. The reaction mixture was slowly warmed up to ambient temperature and then stirred overnight at ambient temperature. The reaction was diluted with 300 mL of saturated aqueous NaHCO₃ and then extracted with DCM. The DCM layer was then washed three times with 10% aqueous NaHSO₄ followed by saturated aqueous NaHCO₃ and finally brine. The DCM layer was then dried over Na₂SO₄, filtered and concentrated under vacuum to give 85.08 g of a tan solid as the title compound **61A**.

Example 61B

8-Bromo-quinolin-2-ol.

- 20 N-(2-Bromo-phenyl)-3-phenyl-acrylamide **61A** (85.0 g, 281.3 mMol) was dissolved in 500 mL of chlorobenzene under an atmosphere of dry N₂. Aluminum trichloride (187.5 g, 1.40 mMol) was then added to the solution and the reaction mixture was subsequently heated to 90°C for 3 hours after which time the temperature was raised to 120°C. After stirring for 1 hour at 120°C for 1 hour, an additional 40 g of aluminum trichloride was added and the reaction mixture was stirred for an additional hour. The reaction mixture was then cooled to ambient temperature and then slowly poured into a solution of 2 L of ice water and 1 L of DCM. The DCM layer was washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give a volume ~100 mL. The pink precipitate was collected and washed with hexanes and then dried to give 44.1 g of the title compound **61B**.

Example 61C

Trifluoro-methanesulfonic acid 8-bromo-quinolin-2-yl ester.

5

8-Bromo-quinolin-2-ol **61B** (24.4 g, 109 mMol) and 2,6-dimethylpyridine (19 mL, 163 mMol) were dissolved in 500 mL of anhydrous DCM under an atmosphere of dry N₂. The reaction mixture was then cooled to 0°C and trifluoromethanesulfonic anhydride (22.0 mL, 131 mMol) was added to the solution dropwise. After the addition was complete, the reaction was warmed to ambient temperature and stirred for 1 hour. The reaction mixture was then quenched with water and partitioned between DCM and aqueous NaHCO₃. The DCM layer was then washed four times with 10% aqueous citric acid, twice with aqueous NaHCO₃ and once with brine. The DCM layer was then dried over Na₂SO₄, filtered and concentrated under vacuum to give the title compound **61C**.

15

Example 61D

(8-Bromo-quinolin-2-yl)-[4-(2-methoxy-ethoxy)-2-nitro-phenyl]-amine.

Trifluoro-methanesulfonic acid 8-bromo-quinolin-2-yl ester **61C** (40.8 g, 114 mMol) and 5-(2-methoxy-ethoxy)-2-nitro-phenylamine **42A** (25.2 g, 119 mMol) were dissolved in 300 mL of toluene under an atmosphere of dry N₂. To this solution was added (49.3 g, 151 mMol) Cs₂CO₃, (3.91g, 6.50 mMol) racemic-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) and (1.98 g, 2.16 mMol) tris(dibenzylideneacetone)dipalladium (0) and the reaction mixture was heated to 80°C and reacted overnight at this temperature. The mixture was then cooled to ambient temperature, concentrated under vacuum, treated with DCM, filtered and concentrated under vacuum to give a red solid. The solid was triturated with EtOAc and dried under vacuum to give 22.0 g of the title compound **61D**.

25

Example 61E

30

N¹-(8-Bromo-quinolin-2-yl)-4-(2-methoxy-ethoxy)-benzene-1,2-diamine.

(8-Bromo-quinolin-2-yl)-[4-(2-methoxy-ethoxy)-2-nitro-phenyl]-amine **61D** (22.0 g, 52.6 mMol) was suspended in a solution of 500 mL of EtOH and 100 mL of water under an atmosphere of dry N₂. To this heterogeneous solution were added ammonium chloride (1.75 g, 63.1 mMol) and iron powder (23.5 g, 421 mMol). The reaction mixture was then heated to 100°C and reacted at this temperature for 6 hours. The reaction mixture was then cooled to ambient temperature, filtered and the filtrate was concentrated under vacuum. The resulting slurry was partitioned between DCM and water. The DCM layer was then washed two more times with water and once with brine. The DCM layer was then dried over Na₂SO₄, filtered and

35

- 5 concentrated under vacuum to give 20.5 g of the title compound **61E**. This material was used without further purification.

Example 61F

8-Bromo-2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinoline.

10

- N¹-(8-Bromo-quinolin-2-yl)-4-(2-methoxy-ethoxy)-benzene-1,2-diamine **61E** (20.5 g, 52.7 mMol) and formamidine acetate (6.03 g, 58.0 mMol) were dissolved in 150 mL of 2-methoxyethanol under an atmosphere of dry N₂. The reaction mixture was heated to reflux and stirred overnight. The reaction mixture was then cooled to ambient temperature that
15 caused the product to precipitate. The pink solid was collected via suction filtration, washed with 2-methoxyethanol and then dried under vacuum to give 16.1 g of the title compound **61F**. The filtrate was concentrated under vacuum and the resulting residue was chromatographed on flash silica gel eluting with a gradient of DCM to MeOH/DCM (1/99) to give an additional 4.40 g of the title compound **61F**.

20

Example 61G


4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol.

- 8-Bromo-2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinoline **61F** (585 mg, 1.47 mMol), was dissolved in 10 mL of dioxane under an atmosphere of dry N₂. To this solution were
25 added 4-hydroxyphenyl boronic acid (240 mg, 1.74 mMol), potassium phosphate (620 mg, 2.92 mMol) and tetrakis(triphenylphosphine)palladium (0) (85 mg, 0.074 mMol). The reaction mixture was heated to 105°C and reacted at this temperature for 48 hours. The reaction mixture was then cooled to ambient temperature, concentrated under vacuum and partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was then washed with brine,
30 dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow residue. The residue was chromatographed on flash silica gel eluting with MeOH/DCM/NH₄OH (2/97.9/0.1) to give 365 mg of a white foam as the title compound **61**.

- C.I. m/z 412 [M+1]; ¹H NMR (CDCl₃) δ 8.64 (s, 1 H), 8.21 (d, J = 8.7 Hz, 1 H), 7.93 (d, J = 9.1 Hz, 1 H), 7.68 (d, J = 7.9 Hz, 2 H), 7.56 (d, J = 9.1 Hz, 1 H), 7.49 (m, 3 H), 7.20 (d, J = 2.5 Hz,
35 1 H), 7.07 (d, J = 7.9 Hz, 2 H), 6.90 (m, 1 H), 4.02 (m, 2 H), 3.65 (m, 2 H), 3.37 (s, 3 H).

Example 62

[2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenoxy)-ethyl]-dimethyl-amine.




5 4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol **61** (100 mg, 0.243 mMol) was dissolved in 600 μ L of anhydrous DMF under an atmosphere of dry N_2 . Cs_2CO_3 (237 mg, 0.729 mMol) and (2-chloro-ethyl)-dimethyl-amine hydrochloride (39 mg, 267 mMol) were added to the reaction mixture. The reaction was heated to 80°C and reacted at this
 10 temperature for 3 hours. The reaction mixture was partitioned between DCM and 0.1 N aqueous NaOH. The DCM layer was washed two more times with 0.1 N aqueous NaOH, dried over Na_2SO_4 , filtered and concentrated under vacuum to give 125 mg of a brown oil. The oil was dissolved into 1 mL of DCM to which was added a 3.0 mL of a solution of 1 N hydrochloric acid (HCl) in ethyl ether. The precipitate was collected, washed with ether,
 15 dissolved in methanol and finally concentrated under vacuum to give 92 mg of the bis-hydrochloride salt of the title compound **62** as a tan solid.

C.I. m/z 483 [M+1]; 1H NMR (CD_3OD) δ 10.2 (s, 1 H), 8.73 (d, J = 9.1 Hz, 1 H), 8.21 (d, J = 9.1 Hz, 1 H), 8.17 (d, J = 8.7 Hz, 1 H), 8.07 (d, J = 8.3 Hz, 1 H), 7.88 (d, J = 6.2 Hz, 1 H), 7.78 (m, 1 H), 7.64 (d, J = 8.7 Hz, 2 H), 7.34 (d, J = 2.1 Hz, 1 H), 7.21 (d, J = 8.3 Hz, 2 H), 7.01 (m, 1
 20 H), 4.46 (m, 2 H), 4.22 (m, 2 H), 3.80 (m, 2 H), 3.69 (m, 2 H), 3.44 (s, 3 H), 3.04 (s, 6 H).

Example 63

2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline.




25 The same procedure that was used in example 1 was followed except that 8-bromo-2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinoline **61F** was used in the place of trifluoromethanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** and t-butyl-1-piperazine-carboxylate in the place of piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound **63** as a tan solid.

- 5 C.I. m/z 404 [M+1]; ¹H NMR (CDCl₃) δ 8.62 (s, 1 H), 8.41 (d, J = 9.1 Hz, 1 H), 8.30 (d, J = 8.7 Hz, 1 H), 7.67 (d, J = 9.1 Hz, 1 H), 7.48 (m, 2 H), 7.34 (d, J = 2.1 Hz, 1 H), 7.23 (m, 1 H), 7.13 (dd, J = 2.5, 9.1 Hz, 1 H), 4.22 (m, 2 H), 3.80 (m, 2 H), 3.46 (s, 3 H), 3.42 (m, 4 H), 3.25 (m, 4 H).

10

Example 64

[2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethyl]-dimethyl-amine.




- 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline **63** (100 mg, 0.248
15 mMol), 2-dimethylamino ethyl chloride hydrochloride (53 mg, 0.37 mMol) and N,N-diisopropylethylamine (130 μL, 0.743 mMol) were dissolved in 1 mL of ACN under an atmosphere of dry N₂. The reaction mixture was heated to 82°C and reacted at this temperature overnight. The reaction mixture was then concentrated under vacuum and then partitioned between DCM and 1 N aqueous NaOH. The DCM layer was then washed again
20 with 1 N aqueous NaOH, dried over Na₂SO₄, filtered and concentrated under vacuum to give a tan residue. The residue was chromatographed on flash silica gel eluting first with MeOH/DCM (5/95) to remove less polar impurities and then with MeOH/DCM/NH₄OH (8/91.9/0.1) to give the title compound **64**.

- C.I. m/z 475 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.44 (d, J = 8.7, 1 H), 8.27 (d, J = 8.7 Hz,
25 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.45 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.22 (m, 1 H), 7.12 (m, 1 H), 4.22 (m, 2 H), 3.81 (m, 2 H), 3.47 (s, 3 H), 3.46 (m, 4 H), 2.83 (m, 4 H), 2.63 (m, 2 H), 2.52 (m, 2 H) 2.30 (s, 6 H).

Example 65

- 30 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyridin-2-ylmethyl-piperazin-1-yl)-quinoline.



5


The same procedure that was used in example 64 was used except that 2-picolyl chloride hydrochloride was used in the place of 2-dimethylamino ethyl chloride hydrochloride to give the title compound **65** as a yellow solid.

C.I. m/z 495 [M+1]; ¹H NMR (CDCl₃) δ 8.62 (s, 1 H), 8.43 (d, J = 9.1, 1 H), 8.28 (d, J = 9.1 Hz, 1 H), 7.68 (m, 2 H), 7.48 (m, 3 H), 7.33 (d, J = 2.1 Hz, 1 H), 7.23 (m, 2 H), 7.10 (m, 1 H), 4.22 (m, 2 H), 3.82 (m, 4 H), 3.49 (s, 3 H), 3.47-3.49 (m, 4 H), 2.92 (m, 4 H).

10

Example 66

2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyridin-3-ylmethyl-piperazin-1-yl)-quinoline.



15


The same procedure that was used in example 64 was used except that 3-picolyl chloride hydrochloride was used in the place of 2-dimethylamino ethyl chloride hydrochloride to give the title compound **66** as a yellow solid.

C.I. m/z 495 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (m, 2 H), 8.52 (dd, J = 2.6, 5.0 Hz, 1 H), 8.43 (d, J = 9.1, 1 H), 8.27 (d, J = 8.7 Hz, 1 H), 7.75 (brd, J = 7.5 Hz, 1 H), 7.66 (d, J = 9.1 Hz, 1 H), 7.46 (m, 2 H), 7.33 (d, J = 2.1 Hz, 1 H), 7.27 (m, 1 H), 7.22 (m, 1 H), 7.10 (m, 1 H), 4.23 (m, 2 H), 3.82 (m, 2 H), 3.67 (s, 2 H), 3.48 (s, 3 H), 3.47 (m, 4 H), 2.82 (m, 4 H).

20

Example 67

25 2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-2-methyl-propan-1-one.



5


The same procedure used in example 54 was followed except that 2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-8-piperazin-1-yl-quinoline **63** was used in the place of 1-{2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **33** and N-t-boc- α -methylalanine was used in the place of N-(tert-butoxycarbonyl)glycine to give the title compound **67**.

- 10 ^1H NMR (CDCl_3) δ 8.62 (s, 1 H), 8.42 (d, J = 8.7 Hz, 1 H), 8.31 (d, J = 8.7 Hz, 1 H), 7.68 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.34 (d, J = 2.5 Hz, 1 H), 7.21 (dd, J = 2.2, 7.5 Hz, 1 H), 7.11 (dd, J = 2.5, 8.7 Hz, 1 H), 4.22 (m, 2 H), 4.17 (m, 4 H), 3.81 (m, 2 H), 3.47 (s, 3 H), 3.39 (m, 4 H), 2.13 (brs, 2 H), 1.49 (s, 6 H).

15

Example 68

(S)-2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-one.




- 20 The same procedure used in example 54 was followed except that 2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-8-piperazin-1-yl-quinoline **63** was used in the place of 1-{2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **33** and N-(tert-butoxycarbonyl)-L-alanine was used in the place of N-(tert-butoxycarbonyl)glycine to give the title compound **68** as a yellow solid.

- 25 ^1H NMR (CDCl_3) δ 8.60 (s, 1 H), 8.40 (d, J = 8.7 Hz, 1 H), 8.30 (d, J = 9.1 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.19 (dd, J = 2.3, 7.5 Hz, 1 H), 7.10 (m, 1 H), 4.20 (m, 2 H), 3.96 (m, 2 H), 3.88 (m, 1 H), 3.79 (m, 4 H), 3.46 (s, 3 H), 3.38 (m, 4 H), 2.11 (brs, 2 H), 1.30 (d, J = 7.0 Hz, 3 H).

5

Example 69

(S)-2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-one.




The same procedure used in example 54 was followed except that 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline **63** was used in the place of 1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **33** and N-(tert-butoxycarbonyl)-D-alanine was used in the place of N-(tert-butoxycarbonyl)glycine to give the title compound **69** as a yellow solid.

¹H NMR (CDCl₃) δ 8.60 (s, 1 H), 8.40 (d, J = 9.1 Hz, 1 H), 8.30 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.33 (m, 1 H), 7.18 (m, 1 H), 7.11 (m, 1 H), 4.20 (m, 2 H), 3.96 (m, 2 H), 3.88 (m, 1 H), 3.79 (m, 4 H), 3.46 (s, 3 H), 3.36 (m, 4 H), 2.10 (brs, 2 H), 1.29 (d, J = 6.2 Hz, 3 H).

20

Example 70

2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone.




The same procedure used in example 54 was followed except that 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline **63** was used in the place of 1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **33** to give the title compound **70**.

¹H NMR (CDCl₃) δ 8.59 (s, 1 H), 8.40 (d, J = 8.7 Hz, 1 H), 8.29 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.51 (dd, J = 2.2, 7.9, 1 H), 7.45 (m, 1 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.18 (dd, J

5 = 2.2, 7.5 Hz, 1 H), 7.09 (d, J = 2.5, 8.7 Hz, 1 H), 4.19 (m, 2 H), 3.96 (m, 2 H), 3.79 (m, 2 H), 3.68 (m, 2 H), 3.54 (m, 2 H), 3.46 (s, 3 H), 3.39 (m, 2 H), 3.33 (m, 2 H), 2.05 (brs, 2 H).

Example 71

(1-Amino-cyclopropyl)-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-methanone.



10

The same procedure used in example 54 was followed except that 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline **63** was used in the place of 1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine **33** and 1-tert-butoxycarbonylamino-cyclopropanecarboxylic acid was used in the place of N-(tert-butoxycarbonyl)glycine to give the title compound **71**.


15

C.I. m/z 487 [M+1]; ¹H NMR (CDCl₃) δ 8.62 (s, 1 H), 8.40 (d, J = 9.1 Hz, 1 H), 8.31 (d, J = 8.7 Hz, 1 H), 7.68 (d, J = 8.7 Hz, 1 H), 7.52 (dd, J = 2.2, 7.9, 1 H), 7.45 (m, 1 H), 7.34 (d, J = 2.1 Hz, 1 H), 7.20 (dd, J = 2.2, 7.5 Hz, 1 H), 7.10 (m, 1 H), 4.20 (m, 2 H), 4.00 (m, 4 H), 3.79 (m, 2 H), 3.46 (s, 3 H), 3.39 (m, 4 H), 2.02 (brs, 2 H), 1.04 (m, 2 H), 0.84 (m, 2 H).

20

Example 72

2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethylamine.



2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline **63** (100 mg, 0.248 mMol), and t-butyl-N-(2-oxoethyl)-carbamate (39.5 mg, 0.248 mMol) were dissolved in 1 mL of methanol under an atmosphere of dry N₂. To this solution was added 200 μL of AcOH followed by NaCNBH₃ (19 mg, 0.297 mMol) and the reaction mixture was then stirred at ambient temperature overnight. The reaction mixture was then concentrated under vacuum and the resulting residue was partitioned between DCM and 1 N aqueous NaOH. The DCM


25

5 layer was then successively washed with 1 N aqueous NaOH and brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellowish-green film. The film was then dissolved in a solution of 1 mL of DCM and 2 mL of TFA under an atmosphere of dry N₂. The reaction mixture was stirred at ambient temperature for 30 minutes after which time it was concentrated under the vacuum. The resulting residue was partitioned between 0.1 N
10 aqueous NaOH. The DCM layer was dried over Na₂SO₄, filtered and concentrated under vacuum to give a green film. The film was chromatographed on flash silica gel eluting with MeOH/DCM (5/95) to remove the less polar impurities and then MeOH/DCM/NH₄OH (5/94.9/0.1) to give 71 mg of the title compound **72**.

C.I. m/z 447 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.45 (d, J = 9.1 Hz, 1 H), 8.27 (d, J = 8.7
15 Hz, 1 H), 7.66 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.22 (m, 1 H), 7.11 (dd, J = 2.5 Hz, 8.7 Hz, 1 H), 4.20 (m, 2 H), 3.80 (m, 2 H), 3.47 (s, 3 H), 3.43 (m, 4 H), 2.89 (t, J = 5.8 Hz, 2 H), 2.80 (m, 4 H), 2.59 (t, J = 5.8 Hz, 2 H), 2.17 (brs, 2 H).

Example 73

20 (R)-2-Amino-3-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-ol.



The same procedure that was used in example 72 was followed except that t-butyl-(S)-(-)-4-formyl-2,2-dimethyl-3-oxazolidinecarboxylate was used in the place of t-butyl-N-(2-oxoethyl)-
25 carbamate to give the title compound **73**.


C.I. m/z 477 [M+1]; ¹H NMR (CDCl₃) δ 8.61 (s, 1 H), 8.43 (d, J = 9.1 Hz, 1 H), 8.27 (d, J = 8.7
Hz, 1 H), 7.64 (d, J = 8.7 Hz, 1 H), 7.45 (m, 2 H), 7.32 (d, J = 2.1 Hz, 1 H), 7.20 (dd, J = 2.6,
7.1 Hz, 1 H), 7.10 (dd, J = 2.5 Hz, 9.1 Hz, 1 H), 4.20 (m, 2 H), 3.80 (m, 2 H), 3.66 (m, 2 H),
3.46 (s, 3 H), 3.45 (m, 4 H), 3.25 (m, 1 H), 2.82 (m, 4 H), 2.65 (m, 1 H), 2.50 (m, 3 H).

30

Example 74

3-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-3-aza-bicyclo[3.1.0]hex-6-
ylamine.

-99-



5


The same procedure that was used in example 33 was followed except that (3-azabicyclo[3.1.0]hex-6-yl)-carbamic acid tert-butyl ester was used in the place of piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound **74** as a yellow solid.

¹H NMR (CDCl₃) δ 8.59 (s, 1 H), 8.22 (d, J = 8.7 Hz, 1 H), 7.99 (d, J = 9.1 Hz, 1 H), 7.59 (d, J = 8.7 Hz, 1 H), 7.37 (m, 2 H), 7.35 (d, J = 2.5 Hz, 1 H), 7.09 (dd, J = 2.5, 9.1 Hz, 1 H), 6.83 (d, J = 7.9 Hz, 1 H), 4.34 (d, J = 9.6 Hz, 2 H), 4.20 (m, 2 H), 3.80 (m, 2 H), 3.46 (s, 3 H), 3.42 (d, J = 9.6 Hz, 2 H), 2.93 (brs, 2 H), 2.65 (s, 1 H), 1.73 (s, 2 H).

10

Example 75

15 (S)-1-{2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-pyrrolidin-3-ylamine.



The same procedure that was used in example 33 was followed except that (3S)-(-)-3-(tert-butoxycarbonyl-amino)pyrrolidine was used in the place of piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound **75** as a tan solid.


20 ¹H NMR (CD₃OD) δ 8.75 (d, J = 1.7 Hz, 1 H), 8.30 (d, J = 8.7 Hz, 1 H), 8.00 (d, J = 9.1 Hz, 1 H), 7.70 (dd, J = 2.1, 8.7 Hz, 1 H), 7.38 (t, J = 7.9 Hz, 1 H), 7.24 (m, 2 H), 7.04 (dd, J = 2.6, 9.1 Hz, 1 H), 6.90 (d, J = 7.9 Hz, 1 H), 4.16 (m, 2 H), 3.90 (m, 1 H), 3.82 (m, 1 H), 3.77 (m, 2 H), 3.66 (m, 1 H), 3.60 (m, 1 H), 3.48 (m, 1 H), 3.43 (s, 3 H), 2.21 (m, 1 H), 1.80 (m, 1 H).

25

Example 76

(R)-1-{2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-pyrrolidin-3-ylamine.

-100-



5


The same procedure that was used in example 33 was followed except that (3R)-(-)-3-(tert-butoxycarbonyl-amino)pyrrolidine was used in the place of piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound **76** as a tan solid.

¹H NMR (CD₃OD) δ 8.80 (s, 1 H), 8.35 (d, J = 8.7 Hz, 1 H), 8.05 (d, J = 8.7 Hz, 1 H), 7.76 (d, J = 8.7 Hz, 1 H), 7.41 (t, J = 7.9 Hz, 1 H), 7.28 (m, 2 H), 7.09 (dd, J = 2.5, 9.1 Hz, 1 H), 6.93 (d, J = 7.5 Hz, 1 H), 4.18 (m, 2 H), 3.91 (m, 1 H), 3.84 (m, 1 H), 3.78 (m, 2 H), 3.66 (m, 1 H), 3.62 (m, 1 H), 3.48 (m, 1 H), 3.44 (s, 3 H), 2.23 (m, 1 H), 1.82 (m, 1 H).

15

Example 77

2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-pyridin-3-yl-quinoline.



Example 77A

8-Benzyloxy-quinolin-2-ol.

20

2,8-Quinolinediol (133.3 g, 0.827 Mol) was dissolved in 800 mL of anhydrous DMF under an atmosphere of dry N₂. To this solution was added potassium carbonate (183 g, 1.32 Mol) followed by benzyl bromide (110 mL, 0.909 Mol) and the solution was then warmed up to 65°C and reacted at this temperature overnight. The reaction mixture was then poured into 9 L of water and the resulting solution was stirred at ambient temperature for 5.5 hours after which time it was filtered. The solid was washed with water, collected and suspended in toluene and finally the solution was concentrated under vacuum to give 142 g of the title compound **77A**.

25

Example 77B

8-Benzyloxy-2-chloro-quinoline.

30

- 5 8-Benzyloxy-quinolin-2-ol **77A** (142 g, 0.565 Mol) was dissolved in 500 mL of DCE under an atmosphere of dry N₂. Oxalyl chloride (99 mL, 1.13 Mol) was added dropwise to this solution followed by 1 mL of DMF. After the addition was complete, the reaction was stirred at ambient temperature for 30 minutes after which time the reaction was warmed to 84°C. The reaction mixture was stirred at this temperature for 10 hours and then concentrated under vacuum.
- 10 The resulting residue was partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was washed again with aqueous saturated NaHCO₃, dried over Na₂SO₄, filtered and concentrated under vacuum to give a brown solid. The solid was recrystallised from toluene to give two crops of (68.3 g and 38.3 g) of the title compound **77B**.

15

Example 77C

(8-Benzyloxy-quinolin-2-yl)-[4-(2-methoxy-ethoxy)-2-nitro-phenyl]-amine.

- 8-Benzyloxy-2-chloro-quinoline **77B** (25.53 g, 94.64 mMol) was added to 350 mL of anhydrous toluene under an atmosphere of dry N₂. To this solution was added 5-(2-methoxy-ethoxy)-2-nitro-phenylamine **42A** (20.08 g, 94.64 mMol), palladium acetate (433 mg, 1.89 mMol), Cs₂CO₃ (43.2 g, 132 mMol), phenyl boronic acid (584 mg, 4.79 mMol) and 1,2-bis(diphenylphosphino)-ethane (2.33 g, 5.68 mMol) and the mixture was then deoxygenated by bubbling argon through for ten minutes. The reaction mixture was then heated to 95°C and reacted at this temperature for two hours. Then the reaction was diluted (while still hot) with
- 20 350 mL of DCE and then filtered while still hot. The filter cake was washed with 100 mL of hot DCE. The filtrate was concentrated to ~175 mL during which time a large amount of precipitate formed. To this mixture was added 400 mL of EtOH and the reaction mixture was stirred overnight. The resulting red precipitate was collected via suction filtration and dried under vacuum to give 36.8g of the title compound as a red solid **77C**.

25

Example 77D

2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-ol.

- (8-Benzyloxy-quinolin-2-yl)-[4-(2-methoxy-ethoxy)-2-nitro-phenyl]-amine **77C** (36.8 g, 85.4 mMol) was suspended in 275 mL of EtOH under an atmosphere of dry N₂. The reaction mixture was then cooled to 0°C after which time 119 mL of triethylamine (NEt₃) followed by 34 mL of formic acid (slowly) were added. The ice bath was removed and the reaction mixture was then heated to ~80°C and reacted at this temperature until the nitro group has been fully reduced to the amine and the benzyl group has been fully removed (as determined by mass
- 35 spectral analysis) which, in this case, was ~2 hours. To this mixture is then added
- 40

5 formamidine acetate (11.56 g, 111 mMol) and the reaction temperature was increased to reflux. After a couple of hours more formamidine acetate (3.20 g) was added and the reaction mixture was stirred at reflux for an additional 4 hours. The reaction mixture was then cooled to ambient temperature and the filtered through Celite™. The Celite™ was washed with
10 copious amounts of a 1:1 MeOH/DCM solution and the combined filtrates were concentrated under vacuum and the resulting solids were triturated with 200 mL of EtOH. The solution was filtered and the precipitate was dried under vacuum to give 23.03 g of the title compound **77D** was a white solid.

Example 77E

15 Trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester.

2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-ol **77D** (23.0 g, 68.6 mMol) was dissolved in a solution of 120 mL of anhydrous DMF and 250 mL of THF under an atmosphere of dry N₂. To this solution were added N-phenyl-bis(trifluoromethanesulfonimide) (26.95 g,
20 75.5 mMol) and NEt₃ (6.75 mL, 137.2 mMol). The reaction mixture was stirred over night at ambient temperature after which time an additional 2.70 g of N-phenyl-bis(trifluoromethanesulfonimide) and an 700 µL of NEt₃ were added. The reaction mixture was stirred an additional 1 hour and then was filtered and the filter cake was washed with ethyl ether. The filter cake was dried under vacuum to give 6.70 g of the title compound **77E**
25 as a white solid. A second crop was obtained by concentrating the filtrate to ~ 75 mL. To this solution was added 200 mL of ethyl ether and the resulting mixture was stirred for 1 hour and then filtered. The filter cake was washed with ethyl ether and then vacuum dried to give an additional 19.91 g of the title compound **77E** for a total of 28.61 g.

30

Example 77F

2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-pyridin-3-yl-quinoline.


Trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** (1.00 g, 2.14 mMol), diethyl(3-pyridyl)-borane (620 mg, 4.36 mMol), lithium chloride (188
35 mg, 4.36 mMol) and tetrakis(triphenylphosphine)palladium (0) (248 mg, 0.215 mMol) were dissolved in a solution of 14 mL of toluene, 4 mL of EtOH and 1.5 mL of 2 N aqueous sodium carbonate (Na₂CO₃) under an atmosphere of dry N₂. The reaction mixture was heated to 90°C and reacted at this temperature overnight. The reaction mixture was then cooled to ambient temperature, concentrated under vacuum and partitioned between DCM and aqueous
40 saturated NaHCO₃. The DCM layer was then washed with brine, dried over Na₂SO₄, filtered

5 and concentrated under vacuum. The resulting residue was chromatographed on flash silica gel eluting with a gradient from MeOH/DCM (1/99) to MeOH/DCM (2/98) to give 460 mg of the title compound 77.

C.I. m/z 397 [M+1]; ¹H NMR (CDCl₃) δ 8.86 (d, J = 1.6 Hz, 1 H), 8.67 (dd, J = 1.3, 4.6 Hz, 1 H), 8.56 (s, 1 H), 8.35 (d, J = 8.7 Hz, 1 H), 8.08 (m, 1 H), 7.88 (m, 1 H), 7.78 (m, 2 H), 7.70 (d, J = 8.7 Hz, 1 H), 7.62 (m, 1 H), 7.42 (m, 1 H), 7.25 (m, 1 H), 6.84 (dd, J = 2.5 Hz, 9.1 Hz, 1 H), 4.15 (m, 2 H), 3.77 (m, 2 H), 3.45 (s, 3 H).

Example 78

2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-8-(6-methoxy-pyridin-3-yl)-quinoline.



15

Trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl ester 77E (850 mg, 1.82 mMol), 2-methoxy-5-pyridineboronic acid (305 mg, 2.00 mMol), potassium phosphate (K₂CO₃) (772 mg, 3.64 mMol) and tetrakis(triphenylphosphine)palladium (0) (208 mg, 0.182 mMol) were dissolved in 5 mL of 1,4-dioxane under an atmosphere of dry N₂. The reaction mixture was heated to 100°C and reacted at this temperature overnight. The reaction mixture was then cooled to ambient temperature, concentrated under vacuum and partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was then washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum. The resulting residue was chromatographed on flash silica gel eluting with a gradient from EtOAc/DCM (20/80) to EtOAc/DCM (75/25) to give 856 mg of the title compound 78.

25


C.I. m/z 427 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.39 (d, J = 2.1 Hz, 1 H), 8.31 (d, J = 8.7 Hz, 1 H), 7.96 (m, 1 H), 7.91 (d, J = 9.1 Hz, 1 H), 7.80 (dd, J = 2.2, 8.3 Hz, 1 H), 7.72 (m, 2 H), 7.56 (m, 1 H), 7.24 (m, 1 H), 6.87 (m, 1 H), 6.83 (d, J = 8.7 Hz, 1 H), 4.14 (m, 2 H), 4.00 (s, 3 H), 3.76 (m, 2 H), 3.44 (s, 3 H).

30

Example 79

4-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-benzoic acid methyl ester.

-104-



5


The same procedure that was used in example 78 was followed except that 4-methoxycarbonylphenylboronic acid was used in the place of 2-methoxy-5-pyridineboronic acid to give the title compound **79** as a white solid.

C.I. m/z 454 [M+1]; ¹H NMR (CDCl₃) δ 8.77 (s, 1 H), 8.37 (d, J = 8.7 Hz, 1 H), 8.13 (d, J = 8.3 Hz, 2 H), 7.91 (d, J = 9.1 Hz, 1 H), 7.88 (dd, J = 2.2, 7.9 Hz, 1 H), 7.77 (m, 4 H), 7.62 (m, 1 H), 7.26 (d, J = 2.1 Hz, 1 H), 6.82 (dd, J = 2.5, 9.1 Hz, 1 H), 4.16 (m, 2 H), 3.99 (s, 3 H), 3.77 (m, 2 H), 3.46 (s, 3 H).

10

Example 80

1-[2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-4-methyl-piperidin-4-ylamine.



15

Example 80A

Piperidine-1,4-dicarboxylic acid tert-butyl ester ethyl ester.

20 Piperidine-4-carboxylic acid ethyl ester (9.80 mL, 63.2 mMol) was dissolved in 60 mL of DCM at ambient temperature under an atmosphere of dry N₂. Di-tert-butyl dicarbonate (13.77 g, 63.16 mMol) was slowly added to the reaction mixture after which time the reaction mixture was stirred overnight. The reaction mixture was then partitioned between DCM and saturated aqueous NaHCO₃. The DCM layer was washed two more times with saturated aqueous
25 NaHCO₃, dried over Na₂SO₄, filtered and concentrated under vacuum to give 16.13 g of the title compound **80A** as an oil.

Example 80B

5 4-Methyl-piperidine-1,4-dicarboxylic acid tert-butyl ester ethyl ester.

Piperidine-1,4-dicarboxylic acid tert-butyl ester ethyl ester **80A** (10.9 g, 42.3 mMol) was dissolved in 44 mL anhydrous THF under an atmosphere of dry N₂. The solution was then cooled to -40°C and 85 mL of a solution of 1 M lithium bis(trimethylsilyl)amide in tetrahydrofuran was slowly added. The reaction mixture was stirred at -40°C for 1 hour after
10 which time iodomethane (5.3 mL, 85 mMol) was added. The solution was allowed to warm up to ambient temperature and after 1 hour was quenched with water. The reaction mixture was partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was washed two more times with aqueous saturated NaHCO₃, dried over Na₂SO₄, filtered and concentrated under vacuum to give 14.78 g of the title compound **80B** as an orange oil. The compound
15 was used without further purification.

Example 80C

4-Methyl-piperidine-1,4-dicarboxylic acid mono-tert-butyl ester.

20 4-Methyl-piperidine-1,4-dicarboxylic acid tert-butyl ester ethyl ester **80B** (12.8 g, 47.2 mMol) was dissolved in a solution of 94 mL of EtOH and 47 ml of aqueous 2 N NaOH. The reaction mixture was heated to 60°C and reacted at this temperature for 6 hours. The reaction mixture was concentrated under vacuum to remove the EtOH. The remaining aqueous solution was washed three times with ethyl ether. The pH was then adjusted to 3 with the careful addition
25 of 1 N HCl. The aqueous layer was then washed 4 times with EtOAc. The EtOAc extracts were combined, dried over Na₂SO₄, filtered and concentrated under vacuum to give 9.54 g of the title compound **80C** as white solid. The compound was used without further purification.

Example 80D

30 4-Carbamoyl-4-methyl-piperidine-1-carboxylic acid tert-butyl ester.

4-Methyl-piperidine-1,4-dicarboxylic acid mono-tert-butyl ester **80C** (9.54 g, 39.2 mMol) was dissolved in 115 mL of anhydrous THF under an atmosphere of dry N₂. The reaction mixture was cooled to 0°C after which time NEt₃ (4.50 mL, 47.1 mMol) followed by ethyl chloroformate
35 (7.10 mL, 60.0 mMol) were added. The mixture was warmed to ambient temperature and stirred for 15 minutes after which time it was cooled back down to 0°C. To this mixture, was added 65 mL of NH₄OH. The reaction mixture was warmed to ambient temperature and the THF was removed under vacuum. The resulting aqueous solution was washed three times with DCM. The DCM extracts were combined, dried over Na₂SO₄, filtered and concentrated

- 5 under vacuum to give 6.19 g of the title compound **80D** as an oil. The compound was used without further purification.

Example 80E

4-Amino-4-methyl-piperidine-1-carboxylic acid tert-butyl ester.

10

- 4-Carbamoyl-4-methyl-piperidine-1-carboxylic acid tert-butyl ester **80D** (6.19 g, 25.6 mMol), [Bis(trifluoroacetoxy)iodo]-benzene (16.5 g, 38.3 mMol) and pyridine (6.20 mL, 76.9 mMol) were dissolved in a solution of 40 mL of ACN and 20 mL of water under an atmosphere of dry N₂. The reaction mixture was stirred for 2 hours at ambient temperature after which time it was diluted with aqueous 1 N NaOH. The resulting solution was washed three times with DCM. The DCM extracts were combined, dried over Na₂SO₄, filtered and concentrated under vacuum to give an oil. The oil was partitioned between DCM and aqueous HCl (pH~3). The aqueous layer was washed 2 more times with DCM, and then was basified with 2 N aqueous NaOH until the pH~10. The basic aqueous layer was washed three times with DCM. The DCM extracts were combined, dried over Na₂SO₄, filtered and concentrated under vacuum to give 4.19 g of the title compound **80E** as a yellow oil. The compound was used without further purification.

Example 80F

25

(4-Methyl-piperidin-4-yl)-carbamic acid benzyl ester.

- 4-Amino-4-methyl-piperidine-1-carboxylic acid tert-butyl ester **80E** (4.19 g, 19.6 mMol) was dissolved in a solution of 50 mL of dioxane and 20 mL of aqueous 2 N NaOH under an atmosphere of dry N₂. To this solution was slowly added benzyl chloroformate (5.6 mL, 39 mMol) and the reaction was stirred for 3 hours at ambient temperature. The reaction mixture was concentrated under vacuum. The reaction mixture was dissolved in DCM and successively washed three with aqueous 1 N NaOH. The DCM layer was dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow oil. The oil was dissolved in 15 mL of TFA under an atmosphere of dry N₂. The reaction mixture was concentrated under vacuum and partitioned between DCM and 0.1 N aqueous NaOH. The DCM layer was dried over Na₂SO₄, filtered and concentrated under vacuum to give 3.06 g of an oil. The oil was chromatographed on flash silica gel eluting with DCM to remove less polar impurities then with MeOH/DCM/NH₄OH (20/79.9/0.1) to give 2.85 g of the title compound **80F** as an oil.

40

Example 80G


5 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-4-methyl-piperidin-4-ylamine.

The same procedure that was used in example 1 was followed except that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** and (4-methyl-piperidin-4-yl)-carbamic acid benzyl ester **80F** was used in the place of piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound **80** as a yellow solid.

C.I. m/z 432 [M+1]; ¹H NMR (CD₃OD) δ 8.80 (s, 1 H), 8.49 (d, J = 9.1 Hz, 1 H), 8.24 (d, J = 9.1 Hz, 1 H), 7.70 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.24 (dd, J = 1.2, 8.7 Hz, 1 H), 7.18 (d, J = 2.5 Hz, 1 H), 7.00 (m, 1 H), 4.12 (m, 2 H), 3.76 (m, 2 H), 3.43 (s, 3 H), 3.26 (m, 2 H), 3.13 (m, 2 H), 1.75 (m, 4 H), 1.17 (s, 3 H).

Example 81

20 1-[2-(6,7-Dihydro-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalen-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.



Example 81A

8-Bromo-2-chloro-quinoline.

25

8-Bromo-quinolin-2-ol **61B** (44.11 g, 196.7 mMol) was dissolved in 450 mL of DCE under an atmosphere of dry N₂. To this solution was added oxalyl chloride (65.3 mL, 749 mMol) dropwise followed by the addition of 400 µL of dimethylformamide. After the addition was complete, the reaction mixture was heated to reflux and reacted for 3 hours. The reaction mixture was concentrated under vacuum. The resulting residue was partitioned between DCM and saturated aqueous NaHCO₃, washed again with saturated aqueous NaHCO₃, dried over Na₂SO₄, filtered and concentrated under vacuum to give 48.40 g of the title compound **81A** as an orange solid.

30

5

Example 81B

6,7-Dihydro-1H-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalene.

5-6-Ethylenedioxy-2-mercaptobenzimidazole (1.27 g, 6.10 mMol) and 3.5 mL of Raney ® nickel (50% slurry in water) were suspended in 30 mL of EtOH under an atmosphere of dry N₂. The reaction mixture was heated to 80°C and reacted at this temperature for 3 hours. The reaction mixture was then filtered through Celite™ and then the Celite™ was washed with ethanol and DCM. The filtrates were combined and concentrated under vacuum to give 750 mg of the title compound **81B** as a foam.

15

Example 81C

1-(8-Bromo-quinolin-2-yl)-6,7-dihydro-1H-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalene.

6,7-Dihydro-1H-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalene **81B** (600 mg, 3.4 mMol) was dissolved in 15 mL of 1-methyl-2-pyrrolidinone under an atmosphere of dry N₂. To this solution was added 60% sodium hydride in oil (136 mg, 3.40 mMol) and the reaction mixture was stirred for 15 minutes. 8-Bromo-2-chloro-quinoline **81A** (750 mg, 3.10 mMol) was then added to the reaction mixture which was subsequently heated to 65°C and reacted at this temperature overnight. The reaction mixture was then cooled to ambient temperature, partitioned between DCM and saturated aqueous NaHCO₃, washed again with saturated aqueous NaHCO₃, dried over Na₂SO₄, filtered and concentrated under vacuum to give 1.60 g of an orange solid. The solid was chromatographed on flash silica gel eluting with a gradient from DCM to MeOH/DCM (0.5/99.5) to give 700 mg of the title compound **81C** as a light yellow solid.

30

Example 81D

{1-[2-(6,7-Dihydro-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalen-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester.


1-(8-Bromo-quinolin-2-yl)-6,7-dihydro-1H-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalene **81C** (700 mg, 1.83 mMol) and piperidin-4-yl-carbamic acid tert-butyl ester (733 mg, 3.66 mMol) were dissolved in 6.0 mL of dioxane under an atmosphere of dry N₂. To this solution was added Cs₂CO₃ (835 mg, 2.56 mMol), racemic-BINAP (68 mg, 0.109 mMol) and tris(dibenzylideneacetone)dipalladium (0) (34 mg, 0.037 mMol) and the reaction mixture was heated to reflux and reacted at this temperature overnight. The mixture was then cooled to ambient temperature, filtered, and concentrated under vacuum to give 1.12 g of an orange

5 film. The film was dissolved in a solution of 5 mL of TFA and 5 mL of DCM under an atmosphere of dry N₂. The reaction mixture was stirred at ambient temperature for 30 minutes after which time it was concentrated under vacuum to give a yellow oil. The oil was partitioned between ethyl ether and 0.1 N aqueous HCl. The aqueous layer was then washed with DCM. The aqueous layer was basified to pH~11 with NaOH. The resulting solution was
10 washed three times with DCM. The DCM extracts were combined, dried over Na₂SO₄, filtered and concentrated to give 500 mg of a yellow solid. The solid was chromatographed on flash silica gel eluting with MeOH/DCM (10/90) to give 127 mg of the title compound **81** as a yellow solid.

C.I. m/z 402 [M+1]; ¹H NMR (CD₃OD) δ 8.63 (s, 1 H), 8.19 (s, 1 H), 8.08 (d, J = 8.7 Hz, 1 H),
15 7.56 (d, J = 9.1 Hz, 1 H), 7.30 (m, 2 H), 7.08 (dd, J = 2.1 Hz, 6.7 Hz, 1 H), 7.06 (s, 1 H), 4.21 (m, 4 H), 3.57 (m, 2 H), 2.63 (m, 3 H), 1.86 (m, 2 H), 1.73 (m, 2 H).

Example 82

20 4-Cyclopropylaminomethyl-1-[2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ol.



The same procedure that was used in example 24 was followed except that 5-(2-Methoxy-ethoxy)-2-nitro-phenylamine **42A** was used in the place of 4-methoxy-2-nitroaniline in example 1B and cyclopropylamine was used in the place of 2.0 M solution of dimethylamine in
25 THF to give the title compound **82** as a yellow foam **82**.


C.I. m/z 488 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.33 (d, J = 9.1 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.65 (d, J = 8.7 Hz, 1 H), 7.44 (m, 2 H), 7.33 (d, J = 2.1 Hz, 1 H), 7.27 (m, 1 H), 7.08 (m, 1 H), 4.21 (m, 2 H), 3.81 (m, 2 H), 3.68 (m, 2 H), 3.47 (s, 3 H), 3.22 (m, 2 H), 2.80 (brs, 2 H), 2.28 (m, 1 H), 1.99 (m, 2 H), 1.73 (m, 2 H), 0.51 (m, 2 H), 0.40 (m, 2 H).

30

Example 83

2-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethanol.

-110-



5


The procedure that was used in example 2 was followed except that 1-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine **33** was used in the place of 1-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** to give the title compound **83** as a yellow solid.

- 10 C.I. m/z 404 [M+1]; ¹H NMR (CD₃OD) δ 8.96 (s, 1 H), 8.69 (d, J = 8.7 Hz, 1 H), 8.42 (d, J = 8.7 Hz, 1 H), 7.93 (d, J = 8.7 Hz, 1 H), 7.56 (d, J = 6.5 Hz, 1 H), 7.48 (t, J = 7.9 Hz, 1 H), 7.33 (d, J = 7.5 Hz, 1 H), 7.29 (d, J = 2.5 Hz, 1 H), 7.22 (m, 1 H), 4.14 (m, 2 H), 3.92 (m, 2 H), 3.85 (m, 2 H), 2.83 (m, 3 H), 2.00 (m, 2 H), 1.83 (m, 2 H).

15

Example 84

1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazole-5-sulfonic acid dimethylamide.



Example 84A

20

4-Amino-N,N-dimethyl-3-nitro-benzenesulfonamide.

- 6.00 g (25.0 mmol) sodium 2-nitroaniline-4-sulfonate was taken into 25.0 mL carbon tetrachloride (CCl₄) with 13.0 g (62.4 mmol) phosphorus pentachloride (PCl₅) and allowed to heat to reflux for five hours. The reaction was allowed to cool to room temperature and was poured into ice water. Chloroform (CHCl₃) was added and normal aqueous work-up yielded a bright yellow solid. Half of the isolated material was taken into 10.0 mL H₂O and cooled to 0 °C. To this solution was slowly added 44.0 mL (87.5 mmol) 2.0 M dimethyl amine in methanol. After the initial exotherm and gas discharge had subsided, reaction was heated to 80°C for four hours. The reaction was allowed to cool to room temperature and the pH was

-111-

- 5 adjusted to 3.0 using dilute HCl. The quenched reaction was allowed to slowly stir at room temperature overnight. 1.13 g (4.62 mmol, 37%) of the desired product **84A** was isolated as a bright yellow solid through filtration.

Example 84B

- 10 {1-[2-(5-Dimethylsulfamoyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester.

The same procedure that used for Example 1 was followed with the exception that 4-amino-N,N-dimethyl-3-nitro-benzenesulfonamide **84A** was used in the place of that 4-methoxy-2-nitroaniline in example 1B to give the title compound **84B** solid.

15


Example 84C

1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazole-5-sulfonic acid dimethylamide.

- 20 {1-[2-(5-Dimethylsulfamoyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester **84B** (129 mg, 0.234 mMol) was dissolved into 1.0 mL of 1,4-dioxane. To this solution, was slowly added 234 μ L (0.936 mmol) of a solution of 4.0 M HCl in 1,4-dioxane. Solid formation was noted and the slurry was allowed to stir at room temperature for two hours. Diethyl ether was added to the reaction mixture, and the crude reaction product was
- 25 isolated through filtration (141 mg yellow solid). This yellow solid was taken into DCM and washed once with 10% K_2CO_3 . Organic layer was dried over $MgSO_4$ and evaporated under reduced pressure. The resulting yellow solid was purified over silica gel (94 $CHCl_3$: 6 CH_3OH : 0.6 NH_4OH) to give 10 mg of the title compound **84** a brownish solid.
- 30 C.I. m/z 451 $[M+1]$; 1H NMR ($CDCl_3$): δ 8.81 (s, 1 H), 8.64 (m, 1 H), 8.38 (m, 1 H), 8.34 (s, 1H), 7.88 (m, 1 H), 7.71 (m, 1H), 7.52 (m, 2 H), 7.29 (m, 1 H), 3.89 (m, 1 H), 2.98 (m, 2 H), 2.74 (s, 6 H), 2.09 (m, 2 H), 1.80 (m, 2 H), 1.78 (m, 2 H).

Example 85

1-[2-(6-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.




- 5 The same procedure that was used in example 84 was followed except that 5-methoxy-2-nitroaniline was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **85** as an off-white solid.

C.I. m/z 374.1 [M+1]; ¹H NMR (CDCl₃): δ 8.60 (s, 1 H), 8.31 (d, J = 8.7 Hz, 1 H), 7.90 (s, 1 H), 7.77 (d, J = 8.7 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.23 (m, 1 H), 7.02 (m, 1 H),
10 3.92 (s, 3 H), 3.91 (m, 2 H), 2.90 (m, 3 H), 2.10 (m, 2 H), 1.89 (m, 2 H), 1.58 (brs, 2 H).

Example 86

1-[2-(5,6-Dimethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.



15

Example 86A

4,5-Dimethoxy-2-nitroaniline.

- 4.50 g (20.0 mmol) 4,5-dimethoxy-1,2-dinitrobenzene was taken into 50.0 mL EtOH/ 8.4 mL
20 AcOH mixture. To this was added 4.00 g powdered iron (0). The reaction was heated to an oil bath temperature of 70°C overnight. The reaction was allowed to cool to room temperature and was poured into 400 mL H₂O. Aqueous layer was extracted several times with diethyl ether. Organic layers were dried over magnesium sulfate and evaporated under reduced pressure to give 4.73 g orange solid as a mixture of starting material and product.
25 This solid was purified over silica gel (30 % DCM in hexanes) to give the title compound **86A** as an orange solid (1.15 g, 5.80 mmol).

Example 86B

1-[2-(5,6-Dimethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.


30

The same procedure that was used in example 84 was followed except that 4,5-dimethoxy-2-nitroaniline **86A** was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the title compound **86** as an off-white solid.

- 5 C.I. m/z 404 [M+1]; ¹H NMR (CDCl₃): δ 8.54 (s, 1H), 8.30 (d, J = 8.7 Hz, 1H), 7.86 (s, 1 H), 7.64 (d, J = 8.7 Hz, 1H), 7.45 (m, 1 H), 7.33 (s, 1 H), 7.30 (m, 1 H), 7.20 (m, 1 H), 4.00 (s, 3H), 3.97 (s, 3H), 3.90 (m, 2 H), 2.90 (m, 3 H), 2.02 (m, 2H), 1.89 (m, 2 H).

Example 87

- 10 2-Dimethylamino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone.



Example 87A

- 15 2-Chloro-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone.

2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline **63** (215 mg, 0.53 mMol) was dissolved in 2.5 mL of anhydrous DCM under an atmosphere of dry N₂. To this solution, was sequentially added 2,6-lutidine (120 μL, 1.06 mMol) and chloroacetyl chloride (630 μL, 0.800 mMol) and then reaction was then stirred overnight at ambient temperature. The reaction mixture was filtered and the solid was collected and dried under vacuum to give 50 mg of the title compound **87A**.

Example 87B


- 25 2-Dimethylamino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone.

2-Chloro-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone **87A** (50 mg, 0.10 mMol) was dissolved in 2.0 mL of a 2.0 M solution of

- 5 dimethylamine in methanol and the reaction mixture was subsequently stirred for 2 hours at ambient temperature. The reaction mixture was concentrated under vacuum and the residue was chromatographed on flash silica gel eluting with MeOH/DCM (2/98) then MeOH/DCM/NH₄OH (4/95.9/ 0.1) and finally MeOH/DCM/NH₄OH (6/93.9/ 0.1) to give 32.5 mg of the title compound **87**.
- 10 C.I. m/z 489 [M+1]; ¹H NMR (CD₃OD): δ 8.79 (s, 1H), 8.49 (d, J = 9.1 Hz, 1H), 8.24 (d, J = 8.7 Hz, 1H), 7.69 (d, J = 9.1 Hz, 1H), 7.48 (m, 1 H), 7.39 (m, 1 H), 7.17 (m, 2 H), 6.98 (dd, J = 2.5, 9.1 Hz, 1 H), 4.11 (m, 2 H), 3.76 (m, 6 H), 3.42 (s, 3 H), 3.14-3.21 (m, 6 H), 2.28 (s, 6 H).

Example 88

- 15 1-[2-(5-Benzyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-4-methyl-piperidin-4-ol.



Example 88A

- 1-[2-(5-Benzyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one.

20

- 2-(5-benzyloxy-benzoimidazol-1-yl)-8-bromo-quinoline (3.46 g, 8.05 mMol), which was prepared using the procedure outlined in example **61** except that 4-benzyloxy-2-nitro-phenylamine was used in the place of 5-(2-methoxy-ethoxy)-2-nitro-phenylamine **42A** in example **61D**, and 4-piperidone hydrochloride hydrate (2.47 g, 16.1 mMol) were dissolved in
- 25 ~40 mL of 1,4-dioxane under an atmosphere of dry N₂. To this solution was added Cs₂CO₃ (8.91 g, 27.4 mMol), racemic-BINAP (300 mg, 0.482 mMol) and tris(dibenzylideneacetone)dipalladium (0) (147 mg, 0.160 mMol) and the reaction mixture was heated to 100°C and reacted at this temperature overnight. The mixture was then cooled to ambient temperature, filtered, and the precipitate was washed several times with DCM/MeOH.

- 5 The combined filtrates were concentrated under vacuum and the resulting residue was purified via flash silica gel chromatography eluting with MeOH/DCM/NH₄OH (1.5/98.3/0.2) to give 1.91 g of the title compound **88A** as an orange solid.

Example 88B

- 10 1-[2-(5-Benzyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-4-methyl-piperidin-4-ol.


1-[2-(5-Benzyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one **88A** (832 mg, 1.85 mMol) was dissolved in 10 mL of anhydrous THF under an atmosphere of dry N₂. The solution was then cooled to -78°C after which time 1.2 mL of a 3 M solution methylmagnesium bromide in
15 THF was added. The reaction mixture warmed up to ambient temperature overnight after which time it was quenched with water. The mixture was then concentrated under vacuum and subsequently partitioned between DCM and saturated aqueous NaHCO₃. The organic layer was then saved, washed again with saturated aqueous NaHCO₃, then with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow foam. The foam was
20 chromatographed over flash silica gel eluting with MeOH/DCM/NH₄OH (1.5/98.3/0.2) to give 585 mg of the title compound **88** as yellow solid.

C.I. m/z 465 [M+1]; ¹H NMR (CDCl₃): δ 8.63 (s, 1H), 8.35 (d, J = 8.7 Hz, 1H), 8.23 (d, J = 8.7 Hz, 1H), 7.60 (d, J = 8.7 Hz, 1H), 7.24-7.48 (m, 9 H), 7.10 (dd, J = 2.5, 8.7 Hz, 1 H), 5.12 (s, 2 H), 3.58 (m, 2 H), 3.26 (m, 2 H), 2.08 (m, 2 H), 1.98 (brs, 1 H), 1.84 (m, 2 H), 1.37 (s, 3 H).

25

Example 89

(4-[2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-benzyl)-dimethyl-amine.




The same procedure that used for Example 6 was followed with the exception that trifluoromethanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E**
30

5 was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F to give the title compound **89**.

C.I. m/z 453 [M+1]; ¹H NMR (CDCl₃) δ 8.57 (s, 1 H), 8.31 (d, J = 9.1 Hz, 1 H), 7.99 (d, J = 9.1 Hz, 1 H), 7.80 (m, 1 H), 7.77 (m, 1 H), 7.65 (m, 3 H), 7.57 (m, 1 H), 7.44 (d, J = 8.1 Hz, 2 H), 7.26 (d, J = 2.5 Hz, 1 H), 6.81 (dd, J = 2.5, 9.1, 1 H), 4.15 (m, 2 H), 3.77 (m, 2 H), 3.56 (s, 2 H), 3.45 (s, 3 H), 2.33 (s, 6 H).

Example 90

(4-{2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-benzyl)-methyl-amine.




15 The same procedure that used for Example 5 was followed with the exception trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F to give the title compound **90**.

C.I. m/z 439 [M+1]; ¹H NMR (CDCl₃) δ 8.56 (s, 1 H), 8.31 (d, J = 9.1 Hz, 1 H), 7.94 (d, J = 9.1 Hz, 1 H), 7.80 (m, 1 H), 7.75 (m, 1 H), 7.63 (m, 3 H), 7.56 (m, 1 H), 7.43 (d, J = 7.9 Hz, 2 H), 7.23 (d, J = 2.5 Hz, 1 H), 6.80 (m, 1 H), 4.14 (m, 2 H), 3.87 (s, 2 H), 3.76 (m, 2 H), 3.43 (s, 3 H), 2.53 (s, 3 H)

Example 91

2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-8-(4-morpholin-4-ylmethyl-phenyl)-quinoline.




- 5 The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and morpholine was used in the place of methylamine in example 5B to give the title compound **91**.
- 10 C.I. m/z 495 [M+1]; ¹H NMR (CDCl₃) δ 8.56 (s, 1 H), 8.31 (d, J = 8.7 Hz, 1 H), 7.98 (d, J = 8.7 Hz, 1 H), 7.81 (m, 1 H), 7.76 (m, 1 H), 7.65 (m, 3 H), 7.57 (m, 1 H), 7.43 (d, J = 7.9 Hz, 2 H), 7.25 (d, J = 2.5 Hz, 1 H), 6.80 (dd, J = 2.5, 9.1, 1 H), 4.15 (m, 2 H), 3.78 (m, 6 H), 3.62 (s, 2 H), 3.46 (s, 3 H), 2.85 (m, 4 H)

15

Example 92


2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamino)-ethanol.



- The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and ethanolamine was used in the place of methylamine in example 5B to give the title compound **92**.
- 20 C.I. m/z 469 [M+1]; ¹H NMR (CD₃OD) δ 8.63 (s, 1 H), 8.14 (d, J = 9.1 Hz, 1 H), 7.71 (d, J = 9.1 Hz, 1 H), 7.68 (dd, J = 2.2, 7.9 Hz, 1 H), 7.59 (d, J = 9.1 Hz, 1 H), 7.54 (dd, J = 2.2, 7.0 Hz, 1 H), 7.43 (m, 1 H), 7.35 (d, J = 8.3 Hz, 2 H), 7.28 (d, J = 8.3 Hz, 2 H), 6.99 (d, J = 2.5 Hz, 1 H), 6.50 (m, 1 H), 4.00 (m, 2 H), 3.83 (s, 2 H), 3.70 (m, 4 H), 3.41 (s, 3 H), 3.21 (m, 2 H)
- 25

Example 93

4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamine.



5


The same procedure that used for Example 78 was followed with the exception that 4-aminomethylphenylboronic acid hydrochloride was used in the place 2-methoxy-5-pyridineboronic acid and the number of equivalents of potassium phosphate was doubled to give the title compound **93**.

- 10 C.I. m/z 425 [M+1]; ¹H NMR (CD₃OD) δ 8.69 (s, 1 H), 8.21 (d, J = 9.1 Hz, 1 H), 7.77 (d, J = 8.7 Hz, 1 H), 7.73 (m, 1 H), 7.66 (d, J = 9.1 Hz, 1 H), 7.59 (dd, J = 1.7, 7.0 Hz, 1 H), 7.48 (m, 1 H), 7.41 (d, J = 8.3 Hz, 2 H), 7.32 (d, J = 8.3 Hz, 2 H), 7.01 (d, J = 2.5 Hz, 1 H), 6.54 (d, J = 2.5, 9.1 Hz, 1 H), 4.02 (m, 2 H), 3.86 (s, 2 H), 3.71 (m, 2 H), 3.41 (s, 3 H).

15

Example 94

2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline.




- 20 The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and pyrrolidine was used in the place of methylamine in example 5B to give the title compound **94**.

- 25 C.I. m/z 479 [M+1]; ¹H NMR (CD₃OD) δ 8.72 (s, 1 H), 8.22 (d, J = 9.1 Hz, 1 H), 7.79 (d, J = 9.1 Hz, 1 H), 7.74 (m, 1 H), 7.69 (d, J = 8.7 Hz, 1 H), 7.57 (dd, J = 1.7, 7.0 Hz, 1 H), 7.48 (m, 1 H), 7.36 (d, J = 8.3 Hz, 2 H), 7.27 (d, J = 8.3 Hz, 2 H), 7.03 (d, J = 2.5 Hz, 1 H), 6.50 (m, 1 H), 4.02 (m, 2 H), 3.70 (m, 2 H), 3.65 (s, 2 H), 3.41 (s, 3 H), 2.55 (m, 4 H), 1.80 (m, 4 H).

5

Example 95

2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline.




The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and piperidine was used in the place of methylamine in example 5B to give the title compound **95**.

C.I. m/z 493 [M+1]; ¹H NMR (CDCl₃) δ 8.56 (s, 1 H), 8.30 (d, J = 8.7 Hz, 1 H), 8.00 (d, J = 9.1 Hz, 1 H), 7.78 (dd, J = 1.2, 7.9 Hz, 1 H), 7.76 (m, 1 H), 7.64 (m, 3 H), 7.56 (m, 1 H), 7.45 (d, J = 7.9 Hz, 2 H), 7.25 (d, J = 2.5 Hz, 1 H), 6.81 (d, J = 2.5, 9.1 Hz, 1 H), 4.15 (m, 2 H), 3.77 (m, 2 H), 3.60 (s, 2 H), 3.46 (s, 3 H), 2.48 (m, 4 H), 1.63 (m, 4 H), 1.47 (m, 2 H).

Example 96

2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline.



20

The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and azetidine was used in the place of methylamine in example 5B to give the title compound **96**.


25

- 5 C.I. m/z 465 [M+1]; ¹H NMR (CD₃OD) δ 8.68 (s, 1 H), 8.17 (d, J = 9.1 Hz, 1 H), 7.72 (d, J = 9.1 Hz, 1 H), 7.71 (m, 1 H), 7.62 (d, J = 9.1 Hz, 1 H), 7.54 (dd, J = 1.7, 7.0 Hz, 1 H), 7.44 (m, 1 H), 7.32 (d, J = 7.9 Hz, 2 H), 7.19 (d, J = 7.9 Hz, 2 H), 7.02 (d, J = 2.5 Hz, 1 H), 6.49 (dd, J = 2.5, 9.1 Hz, 1 H), 4.01 (m, 2 H), 3.70 (m, 2 H), 3.60 (s, 2 H), 3.41 (s, 3 H), 3.26 (m, 4 H), 2.09 (m, 2 H).

10

Example 97

1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-cis-pyrrolidine-3,4-diol.




- 15 The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and cis-pyrrolidine-3,4-diol was used in the place of methylamine in example 5B to give the title compound **97**.
- 20 C.I. m/z 511 [M+1]; ¹H NMR (CD₃OD) δ 8.68 (s, 1 H), 8.18 (d, J = 9.1 Hz, 1 H), 7.74 (d, J = 9.1 Hz, 1 H), 7.71 (dd, J = 1.3, 8.3 Hz, 1 H), 7.63 (d, J = 9.1 Hz, 1 H), 7.54 (dd, J = 1.3, 7.0 Hz, 1 H), 7.44 (m, 1 H), 7.33 (d, J = 8.0 Hz, 2 H), 7.25 (d, J = 8.0 Hz, 2 H), 7.00 (d, J = 2.5 Hz, 1 H), 6.50 (m, 1 H), 4.14 (m, 2 H), 4.01 (m, 2 H), 3.71 (m, 2 H), 3.66 (s, 2 H), 3.42 (s, 3 H), 2.96 (m, 2 H), 2.55 (m, 2 H).

25

Example 98

R,R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol).

-121-



5

The same procedure that used for Example 5 was followed with the exception that trifluoromethanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoromethanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in step 1F and R,R-trans-pyrrolidine-3,4-diol was used in the place of methylamine in step 5B to give the title compound **98**.


10

C.I. m/z 511 [M+1]; ¹H NMR (CD₃OD) δ 8.87 (s, 1 H), 8.40 (d, J = 8.7 Hz, 1 H), 7.96 (d, J = 9.1 Hz, 1 H), 7.88 (m, 2 H), 7.71 (dd, J = 1.2, 7.0 Hz, 1 H), 7.58 (m, 1 H), 7.54 (d, J = 8.3 Hz, 2 H), 7.44 (d, J = 8.3 Hz, 2 H), 7.12 (d, J = 2.5 Hz, 1 H), 6.67 (dd, J = 2.5, 9.1 Hz, 1 H), 4.10 (m, 4 H), 3.77 (m, 4 H), 3.44 (s, 3 H), 3.04 (m, 2 H), 2.60 (m, 2 H).

15

Example 99

1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol.



The same procedure that used for Example 5 was followed with the exception that trifluoromethanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoromethanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and racemic 3-pyrrolidinol was used in the place of methylamine in example 5B to give the title compound **99**.

20


C.I. m/z 495 [M+1]; ¹H NMR (CD₃OD) δ 8.60 (s, 1 H), 8.08 (d, J = 8.7 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.63 (m, 1 H), 7.53 (d, J = 9.1 Hz, 1 H), 7.49 (dd, J = 1.2, 7.1 Hz, 1 H), 7.39 (m, 1 H), 7.39 (d, J = 8.3 Hz, 2 H), 7.26 (d, J = 8.3 Hz, 2 H), 6.97 (d, J = 2.5 Hz, 1 H), 6.44 (dd, J = 2.5,

25

- 5 9.1 Hz, 1 H), 4.35 (m, 1 H), 3.98 (m, 2 H), 3.69 (m, 2 H), 3.61 (d, J = 12.5 Hz, 1 H), 3.57 (d, J = 12.5 Hz, 1 H), 3.41 (s, 3 H), 2.82 (m, 1 H), 2.67 (m, 1 H), 2.48 (m, 2 H), 2.12 (m, 1 H), 1.70 (m, 1 H).

Example 100

- 10 R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol).




- The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and R-3-pyrrolidinol was used in the place of methylamine in example 5B to give the title compound **100**.

- C.I. m/z 495 [M+1]; ¹H NMR (CD₃OD) δ 8.69 (s, 1 H), 8.19 (d, J = 9.1 Hz, 1 H), 7.76 (d, J = 9.1 Hz, 1 H), 7.71 (m, 1 H), 7.64 (d, J = 8.7 Hz, 1 H), 7.55 (dd, J = 1.3, 7.0 Hz, 1 H), 7.46 (m, 1 H), 7.34 (d, J = 8.3 Hz, 2 H), 7.27 (d, J = 8.3 Hz, 2 H), 7.01 (d, J = 2.5 Hz, 1 H), 6.49 (dd, J = 2.5, 9.1 Hz, 1 H), 4.36 (m, 1 H), 4.01 (m, 2 H), 3.70 (m, 2 H), 3.67 (d, J = 12.5 Hz, 1 H), 3.62 (d, J = 12.5 Hz, 1 H), 3.42 (s, 3 H), 2.84 (m, 1 H), 2.70 (m, 1 H), 2.55 (m, 1 H), 2.50 (m, 1 H), 2.15 (m, 1 H), 1.72 (m, 1 H).

Example 101

- 25 S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol).

-123-



5

The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and S-3-pyrrolidinol was used in the place of methylamine in example 5B to give the title compound **101**.


10

C.I. m/z 495 $[M+1]$; 1H NMR (CD_3OD) δ 8.84 (s, 1 H), 8.37 (d, $J = 8.7$ Hz, 1 H), 7.93 (d, $J = 9.1$ Hz, 1 H), 7.86 (m, 2 H), 7.68 (dd, $J = 1.7, 7.1$ Hz, 1 H), 7.56 (m, 1 H), 7.53 (d, $J = 7.9$ Hz, 2 H), 7.40 (d, $J = 7.9$ Hz, 2 H), 7.11 (d, $J = 2.5$ Hz, 1 H), 6.63 (dd, $J = 2.5, 9.1$ Hz, 1 H), 4.39 (m, 1 H), 4.09 (m, 2 H), 3.74 (m, 4 H), 3.43 (s, 3 H), 2.90 (m, 1 H), 2.76 (m, 1 H), 2.62 (m, 1 H), 2.52 (m, 1 H), 2.19 (m, 1 H), 1.75 (m, 1 H).

15

Example 102

1-(4-[2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-benzyl)-azetidin-3-ol.



The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and azetidin-3-ol was used in the place of methylamine in example 5B to give the title compound **102**.

C.I. m/z 481 $[M+1]$; 1H NMR (CD_3OD) δ 8.52 (s, 1 H), 8.00 (d, $J = 9.1$ Hz, 1 H), 7.63 (d, $J = 8.7$ Hz, 1 H), 7.56 (dd, $J = 2.3, 7.9$ Hz, 1 H), 7.45 (m, 2 H), 7.33 (m, 1 H), 7.26 (d, $J = 8.3$ Hz, 2 H), 7.15 (d, $J = 8.3$ Hz, 2 H), 6.96 (d, $J = 2.5$ Hz, 1 H), 6.46 (dd, $J = 2.5, 8.7$ Hz, 1 H), 4.86 (brs, 1


25

-124-

5 H), 4.37 (m, 1 H), 3.96 (m, 2 H), 3.68 (m, 2 H), 3.63 (s, 2 H), 3.59 (m, 2 H), 3.40 (s, 3 H), 2.98 (m, 2 H).

Example 103

10 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-quinoline.




The same procedure that used for Example 5 was followed with the exception that trifluoromethanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoromethanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and 1-methylpiperazine was used in the place of methylamine in example 5B to give the title compound **103**.

15 C.I. m/z 508 [M+1]; ¹H NMR (CDCl₃) δ 8.56 (s, 1 H), 8.30 (d, J = 9.1 Hz, 1 H), 7.98 (d, J = 8.7 Hz, 1 H), 7.80 (m, 1 H), 7.75 (m, 1 H), 7.62 (m, 3 H), 7.57 (m, 1 H), 7.44 (d, J = 8.3 Hz, 2 H), 7.24 (d, J = 2.1 Hz, 1 H), 6.80 (dd, J = 2.5, 8.7 Hz, 1 H), 4.15 (m, 2 H), 3.77 (m, 2 H), 3.62 (s, 20 2 H), 3.45 (s, 3 H), 2.70 (m, 8 H), 2.30 (s, 3 H).

Example 104

4-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperazine-1-carboxylic acid tert-butyl ester.




25

- 5 The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and 1-tert-butoxycarbonyl-piperazine was used in the place of methylamine in example 5B to give the title compound **104**.
- 10 C.I. m/z 594 [M+1]; ¹H NMR (CDCl₃) δ 8.58 (s, 1 H), 8.35 (d, J = 9.1 Hz, 1 H), 8.00 (d, J = 9.1 Hz, 1 H), 7.82 (m, 1 H), 7.78 (m, 1 H), 7.67 (m, 3 H), 7.60 (m, 1 H), 7.47 (d, J = 7.9 Hz, 2 H), 7.26 (d, J = 2.1 Hz, 1 H), 6.83 (dd, J = 2.5, 8.7 Hz, 1 H), 4.17 (m, 2 H), 3.79 (m, 2 H), 3.64 (s, 2 H), 3.49 (m, 4 H), 3.46 (s, 3 H), 2.50 (m, 4 H), 1.45 (s, 9 H).

15

Example 105

[1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-yl]-carbamic acid tert-butyl ester.




- The same procedure that used for Example 5 was followed with the exception that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** in example 1F and 4-N-tert-butoxycarbonyl-aminopiperidine was used in the place of methylamine in example 5B to give the title compound **105**.
- 20 C.I. m/z 608 [M+1]; ¹H NMR (CD₃OD) δ 8.92 (s, 1 H), 8.48 (d, J = 9.1 Hz, 1 H), 8.04 (d, J = 9.1 Hz, 1 H), 7.96 (d, J = 9.1 Hz, 1 H), 7.92 (m, 1 H), 7.75 (m, 1 H), 7.61 (m, 3 H), 7.45 (d, J = 7.9 Hz, 2 H), 7.16 (d, J = 2.5 Hz, 1 H), 6.65 (dd, J = 2.5, 8.7 Hz, 1 H), 4.12 (m, 2 H), 3.77 (m, 2 H), 3.65 (s, 2 H), 3.44 (s, 3 H), 3.41 (m, 1 H), 3.00 (m, 2 H), 2.22 (m, 2 H), 1.88 (m, 2 H), 1.75 (m, 2 H), 1.42 (s, 9 H).

30

Example 106

1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-ylamine.



5


[1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-yl]-carbamic acid tert-butyl ester **105** (110 mg, 0181 mMol) was dissolved in a solution of 3 mL of TFA and 3 mL of DCM under an atmosphere of dry N_2 . The reaction mixture was stirred for 40 minutes after which time it was concentrated under vacuum and the resulting residue was partitioned between DCM and 1 N aqueous NaOH. The DCM layer was then washed with again with 1 N aqueous NaOH, dried over Na_2SO_4 , filtered and concentrated under vacuum to give 45 mg of the title compound **106**.

C.I. m/z 508 $[M+1]$; 1H NMR ($CDCl_3$) δ 8.55 (s, 1 H), 8.29 (d, J = 8.7 Hz, 1 H), 7.98 (d, J = 9.1 Hz, 1 H), 7.78 (d, J = 8.3 Hz, 1 H), 7.76 (m, 1 H), 7.60 (m, 4 H), 7.42 (d, J = 7.9 Hz, 2 H), 7.23 (d, J = 2.5 Hz, 1 H), 6.80 (m, 1 H), 4.14 (m, 2 H), 3.76 (m, 2 H), 3.59 (s, 2 H), 3.44 (s, 3 H), 2.93 (m, 2 H), 2.68 (m, 1 H), 2.11 (m, 2 H), 1.94 (brs, 2 H), 1.80 (m, 2 H), 1.45 (m, 2 H).

15

Example 107

(1-(2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-yl)-methanol.



20

The same procedure that was used in example 1 was followed except that trifluoromethanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** was used in the place of trifluoromethanesulfonic acid 2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl ester **1E** and 4-(hydroxymethyl)-piperidine was used in the place of piperidin-4-yl-carbamic acid tert-butyl ester in example 1F to give the title compound **107** as a yellow solid.

25


C.I. m/z 433 $[M+1]$; 1H NMR ($CDCl_3$) δ 8.66 (s, 1 H), 8.38 (d, J = 9.1 Hz, 1 H), 8.29 (d, J = 9.1 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.47 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.28 (m, 1 H), 7.10

- 5 (dd, $J = 2.5, 9.1$ Hz, 1 H), 4.21 (m, 2 H), 3.95 (m, 2 H), 3.80 (m, 2 H), 3.63 (m, 2 H), 3.47 (s, 3 H), 2.83 (m, 2 H), 1.95 (m, 2 H), 1.75 (m, 3 H).

Example 108

(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-methyl-amine.

10



Example 108A

1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidine-4-carbaldehyde.

- 15 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methanol **107**
 (1.47 g, 3.40 mMol) was dissolved in 15 mL of anhydrous DCM under an atmosphere of dry N_2 . To this mixture was added ~2 gm of 4A molecular sieves, 4-methylmorpholine N-oxide (1.19 g, 10.2 mMol) and finally tetrapropylammonium perruthenate (119 mg, 0.340 mMol). The mixture was stirred at ambient temperature over night after which time it was filtered
 20 through Celite™. The Celite™ was washed with DCM and the resulting combined filtrate was concentrated under vacuum. The resulting residue was chromatographed on flash silica gel eluting with a gradient from MeOH/DCM (1/99) to MeOH/DCM (2/98) to give 446 mg of the title compound **108A**.

25

Example 108B

(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-methyl-amine.

- 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidine-4-carbaldehyde **108A**
 30 (100 mg, 0.232 mMol) was dissolved in 1 mL of MeOH under an atmosphere of dry N_2 . To this solution was added 580 μ L of a 2.0 M solution of methylamine in MeOH. AcOH (~100 μ L) was added to this solution until the pH ~ 5 after which time $NaCNBH_3$ (30 mg, 0.46 mMol) was added to the mixture. The reaction mixture was stirred overnight at ambient temperature. The reaction mixture was then concentrated under vacuum and partitioned between DCM and


- 5 1 N aqueous NaOH. The DCM layer was saved and washed again with 1 N aqueous NaOH, dried over Na₂SO₄, filtered and concentrated under vacuum. The resulting residue was chromatographed on flash silica gel eluting with a gradient from MeOH/DCM (3/97) to MeOH/DCM (5/95) and then a gradient from MeOH/DCM/NH₄OH (7/92.9/0.1) to MeOH/DCM/NH₄OH (9/90.9/0.1) to give 60 mg of the title compound **108** as a yellow solid.
- 10 C.I. m/z 446 [M+1]; ¹H NMR (CD₃OD) δ 8.96 (s, 1 H), 8.68 (d, J = 9.1 Hz, 1 H), 8.44 (d, J = 9.1 Hz, 1 H), 7.93 (d, J = 8.7 Hz, 1 H), 7.57 (d, J = 7.9 Hz, 1 H), 7.49 (m, 1 H), 7.34 (d, J = 7.9 Hz, 1 H), 7.27 (d, J = 2.5 Hz, 1 H), 7.10 (dd, J = 2.5, 9.1 Hz, 1 H), 4.19 (m, 2 H), 3.86 (m, 2 H), 3.79 (m, 2 H), 3.45 (s, 2 H), 2.78 (m, 2 H), 2.63 (m, 2 H), 2.45 (s, 3 H), 1.92 (m, 2 H), 1.68 (m, 3 H).

15

Example 109

2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-piperidin-1-yl]-quinoline.

20



The same procedure that was used in example 108 was followed except that 1-methylpiperazine was used in the place of methylamine in example 108B to give the title compound **109**.


- 25 C.I. m/z 515 [M+1]; ¹H NMR (CD₃OD) δ 8.79 (s, 1 H), 8.48 (d, J = 9.1 Hz, 1 H), 8.19 (d, J = 8.7 Hz, 1 H), 7.67 (d, J = 9.1 Hz, 1 H), 7.38 (m, 2 H), 7.16 (d, J = 2.1 Hz, 1 H), 7.12 (d, J = 7.4 Hz, 1 H), 6.90 (m, 1 H), 4.11 (m, 2 H), 3.75 (m, 2 H), 3.63 (m, 2 H), 3.43 (s, 3 H), 2.20-2.60 (m, 15 H), 1.76 (m, 2 H), 1.55 (m, 1 H), 1.45 (m, 2 H).

30

Example 110

(1-(2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylmethyl)-dimethylamine.

5




The same procedure that was used in example 108 was followed except that dimethylamine was used in the place of methylamine in step 108B to give the title compound 110.

C.I. m/z 460 [M+1]; ¹H NMR (CD₃OD) δ 8.85 (s, 1 H), 8.55 (d, J = 9.1 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.76 (d, J = 8.7 Hz, 1 H), 7.45 (d, J = 7.9 Hz, 1 H), 7.40 (m, 1 H), 7.21 (m, 2 H), 6.98 (dd, J = 2.1, 9.1 Hz, 1 H), 4.14 (m, 2 H), 3.75 (m, 4 H), 3.43 (s, 3 H), 2.64 (m, 2 H), 2.27 (m, 2 H), 2.24 (s, 6 H), 1.83 (m, 2 H), 1.63 (m, 1 H), 1.54 (m, 2 H).

Example 111

15 1-(1-(2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylmethyl)-pyrrolidin-3-ol.




20 The same procedure that was used in example 108 was followed except that racemic 3-pyrrolidinol was used in the place of methylamine in example 108B to give the title compound 111.

C.I. m/z 502 [M+1]; ¹H NMR (CD₃OD) δ 8.93 (s, 1 H), 8.65 (d, J = 8.7 Hz, 1 H), 8.39 (d, J = 8.7 Hz, 1 H), 7.88 (d, J = 8.7 Hz, 1 H), 7.55 (m, 1 H), 7.46 (m, 1 H), 7.30 (dd, J = 1.3, 7.5 Hz, 1 H), 7.25 (d, J = 2.1 Hz, 1 H), 7.06 (m, 1 H), 4.35 (m, 1 H), 4.18 (m, 2 H), 3.78 (m, 4 H), 3.45 (s, 3 H), 2.86 (m, 1 H), 2.74 (m, 3 H), 2.58 (m, 1 H), 2.50 (m, 3 H), 2.12 (m, 1 H), 1.93 (m, 2 H), 1.60-1.75 (m, 4 H).

5

Example 112

C-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methylamine.




The same procedure that was used in example 108 was followed except that ammonium acetate was used in the place of methylamine in example 108B to give the title compound **112**.

C.I. m/z 432 $[M+1]$; 1H NMR (CD_3OD) δ 8.91 (s, 1 H), 8.61 (d, $J = 8.7$ Hz, 1 H), 8.37 (d, $J = 8.7$ Hz, 1 H), 7.85 (d, $J = 8.7$ Hz, 1 H), 7.52 (d, $J = 7.1$ Hz, 1 H), 7.45 (m, 1 H), 7.29 (d, $J = 7.5$ Hz, 1 H), 7.23 (d, $J = 2.1$ Hz, 1 H), 7.04 (dd, $J = 2.5, 9.1$ Hz, 1 H), 4.17 (m, 2 H), 3.79 (m, 4 H), 3.44 (s, 3 H), 2.65-2.80 (m, 4 H), 1.89 (m, 2 H), 1.60 (m, 3 H).

15

Example 113

1-[2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-4-methyl-piperidin-4-ol.



20

Example 113A

1-[8-(4-Hydroxy-4-methyl-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-ol.

1-[2-(5-Benzoyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-4-methyl-piperidin-4-ol **88** (535 mg, 1.15 mMol) was dissolved in 5 mL of anhydrous DCM under an atmosphere of dry N_2 . The solution

25

-131-

5 was then cooled to 0°C after which time 32 μ L of NEt_3 and 1.50 mL of a 1 M solution of dimethylbromoborane in DCM were added. The reaction mixture was allowed to warm up to ambient temperature and reacted at this temperature for ~ 2 hours. The mixture was then partitioned between iso-propanol/DCM (18/82) and saturated aqueous NaHCO_3 . The aqueous layer was extracted two more times with iso-propanol/DCM. The organic extracts
10 were combined, dried over Na_2SO_4 , filtered and concentrated under vacuum to give a yellow solid. The solid was chromatographed over flash silica gel eluting with $\text{MeOH/DCM/NH}_4\text{OH}$ (4/95.8/0.2) to give 189 mg of the title compound **113A** as yellow solid.

Example 113B

15 1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-4-methyl-piperidin-4-ol.


1-[8-(4-Hydroxy-4-methyl-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-ol **113A** (189 mg, 0.505 mMol), Cs_2CO_3 (592 mg, 1.81 mMol), sodium iodide (76 mg, 0.505 mMol) and (2-dimethylamino)ethyl chloride hydrochloride (87 mg, 0.61 mMol) were added to 2 mL of
20 anhydrous DMF and the resulting mixture was heated to 80°C under an atmosphere of dry N_2 . The reaction was run for 48 hours after which time it was cooled to ambient temperature and concentrated under vacuum. The residue was chromatographed over flash silica gel eluting with a gradient starting from MeOH/DCM (3/97) to MeOH/DCM (8/92) then switched to $\text{MeOH/DCM/NH}_4\text{OH}$ (8/91.8/0.2) to give 67mg of the title compound **113** as a yellow oil.

25 C.I. m/z 446 $[\text{M}+1]$; ^1H NMR (CDCl_3): δ 8.64 (s, 1H), 8.33 (d, J = 8.7 Hz, 1H), 8.29 (d, J = 8.7 Hz, 1H), 7.66 (d, J = 8.7 Hz, 1H), 7.46 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.28 (dd, J = 2.9, 5.8 Hz, 1 H), 7.08 (dd, J = 2.5, 9.1 Hz, 1 H), 4.20 (m, 2 H), 3.60 (m, 2 H), 3.26 (m, 2 H), 2.86 (m, 2 H), 2.41 (s, 6 H), 2.10 (m, 2 H), 1.85 (m, 2 H), 1.38 (s, 3 H).

30

Example 114

1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol.



5

Example 114A

1-[2-(5-Benzoyloxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol.

1-[2-(5-Benzoyloxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one **88A** (790 mg, 1.76 mMol)
10 was dissolved in 6 mL of MeOH under an atmosphere of dry N₂. To this solution was added sodium borohydride (66 mg, 1.76 mMol) and the reaction mixture was stirred at ambient temperature overnight. The mixture was then concentrated under vacuum and the resulting residue was partitioned between DCM and 0.1 N aqueous NaOH. The DCM layer was saved and washed again with 0.1 N aqueous NaOH and then two times with brine. The DCM layer
15 was dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow solid. The solid was purified via flash silica gel chromatography eluting with MeOH/DCM/NH₄OH (1.5/98.3/0.2) to give 690 mg of the title compound **114A**.

Example 114B

20 1-[8-(4-Hydroxy-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazol-5-ol.

1-[2-(5-Benzoyloxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol **114A** (440 mg, 0.976 mMol) was suspended in 10 mL of DCE under an atmosphere of dry N₂. To this heterogeneous solution was added 5 mL of a 1 M solution of boron tribromide in DCM. The
25 reaction mixture was stirred for 1 hour at ambient temperature and then heated to reflux and reacted at that temperature for 2 hours. The reaction mixture was then cooled to ambient temperature and was added to ice water. The pH of the resulting reaction mixture was adjusted to 8 with the addition of Na₂CO₃ and the solution was extracted 5 times with isopropanol/DCM (18/82). The organic extracts were combined and dried over Na₂SO₄,
30 filtered and concentrated under vacuum to give a yellow solid. The solid was purified via flash silica gel chromatography eluting with MeOH/DCM/NH₄OH (3/96.8/0.2) to give 201 mg of the title compound **114B**.

Example 114C


35 1-[2-[5-(2-Dimethylamino-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ol.

5 1-[8-(4-Hydroxy-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazol-5-ol **114B** (201 mg, 0.558 mMol), Cs_2CO_3 (400 mg, 1.22 mMol), sodium iodide (84 mg, 0.559 mMol) and (2-dimethylamino)ethyl chloride hydrochloride (88 mg, 0.62 mMol) were added to 5 mL of anhydrous DMF and the resulting mixture was heated to 65°C under an atmosphere of dry N_2 . After one hour, the reaction temperature was increased to 100°C. After reacting at 100°C for
10 1 hour, an additional 44 mg of (2-dimethylamino)ethyl chloride hydrochloride and 200 mg of Cs_2CO_3 were added. After reacting overnight at 100°C, the reaction mixture was cooled to ambient temperature and concentrated under vacuum. The resulting residue was partitioned between DCM and aqueous 0.1 N NaOH. The aqueous layer was extracted 2 more times with DCM and the organic extracts were combined, dried over Na_2SO_4 , filtered and
15 concentrated under vacuum to give a yellow oil. The residue was purified via High Pressure Liquid Chromatography (C-8 reverse phase; linear gradient from 0.1% TFA in water to 0.1% TFA in water/acetonitrile (50/50) over 8 minutes) to give 43 mg of the title compound **114** as a yellow oil.

C.I. m/z 432 [M+1]; ^1H NMR (CD_3OD) δ 8.92 (s, 1H), 8.68 (d, J = 9.1 Hz, 1H), 8.38 (d, J = 9.1 Hz, 1H), 7.87 (d, J = 9.1 Hz, 1H), 7.52 (dd, J = 2.3, 7.9 Hz, 1 H), 7.45 (m, 1 H), 7.30 (dd, J = 1.2, 7.5 Hz, 1 H), 7.25 (d, J = 2.1 Hz, 1 H), 7.13 (m, 1 H), 4.17 (m, 2 H), 3.80 (m, 1 H), 3.70 (m, 2 H), 2.91 (m, 2 H), 2.81 (m, 2 H), 2.37 (s, 6 H), 2.05 (m, 2 H), 1.92 (m, 2 H).

Example 115

25 S,S-(1-(4-(2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl)-benzyl)-trans-pyrrolidine-3,4-diol).



The same procedure that used for Example **98** was followed with the exception S,S-trans-pyrrolidine-3,4-diol was used in the place of R,R-trans-pyrrolidine-3,4-diol to give the title
30 compound **115**.


C.I. m/z 511 [M+1]; ^1H NMR (CD_3OD) δ 8.95 (s, 1 H), 8.49 (d, J = 9.1 Hz, 1 H), 8.04 (d, J = 9.1 Hz, 1 H), 7.98 (d, J = 9.1 Hz, 1 H), 7.94 (m, 1 H), 7.761 (m, 1 H), 7.65 (m, 1 H), 7.62 (d, J =

-134-

- 5 8.3 Hz, 2 H), 7.51 (d, J = 8.3 Hz, 2 H), 7.17 (d, J = 2.5 Hz, 1 H), 6.74 (dd, J = 2.5, 9.1 Hz, 1 H), 4.13 (m, 4 H), 3.88 (d J = 12.5 Hz, 1 H), 3.82 (d, J = 12.5 Hz, 1 H), 3.77 (m, 2 H), 3.44 (s, 3 H), 3.11 (m, 2 H), 2.67 (m, 2 H).

Example 116

- 10 4-{2-[5-(3-Amino-propoxy)-benzimidazol-1-yl]-quinolin-8-yl}-phenol.



Example 116A

Trifluoro-methanesulfonic acid 2-(5-hydroxy-benzimidazol-1-yl)-quinolin-8-yl ester.

- 15 Trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl ester **77E** (2.00 g, 4.28 mMol) was suspended in 20 mL of anhydrous DCM under an atmosphere of dry N₂. To this heterogeneous solution was added 12.9 mL of a solution of 1 M borontribromide in DCM and the resulting mixture was stirred at ambient temperature for 24
- 20 hours. The reaction mixture was then poured into ice water. The resulting heterogeneous mixture was neutralised with the addition of NaHCO₃ and then partitioned between isopropanol/DCM (18/82) and aqueous saturated NaHCO₃. The organic layer was washed again aqueous saturated NaHCO₃, dried over Na₂SO₄, filtered, concentrated under vacuum to give 890 mg of a brownish yellow solid as the title compound **116A**.

25

Example 116B

Trifluoro-methanesulfonic acid 2-[5-(3-tert-butoxycarbonylamino-propoxy)-benzimidazol-1-yl]-quinolin-8-yl ester.

- 30 (3-Hydroxy-propyl)-carbamic acid tert-butyl ester (180 mg, 1.03 mMol) was dissolved in 7 mL of anhydrous THF was under an atmosphere of dry N₂. Trifluoro-methanesulfonic acid 2-(5-hydroxy-benzimidazol-1-yl)-quinolin-8-yl ester **116A** (420 mg, 1.03 mMol) and triphenyl phosphine (538 mg, 2.05 mMol) were added to the solution resulting in a brown heterogeneous solution. A solution of 320 µL of diethyl azodicarboxylate dissolved in 3 mL of
- 35 anhydrous THF was then added. After ~ 30 minutes, the reaction mixture was concentrated

- 5 under vacuum and the residue was chromatographed on flash silica gel eluting with MeOH/DCM/NH₄OH (1/98.8/0.2) to give 184 mg of the title compound **116B**.

Example 116C

- 10 [3-(1-{8-[4-(Tetrahydro-pyran-2-yloxy)-phenyl]-quinolin-2-yl}-1H-benzoimidazol-5-yloxy)-propyl]-carbamic acid tert-butyl ester.

- Trifluoro-methanesulfonic acid 2-[5-(3-tert-butoxycarbonylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **116B** (144 mg, 0.254 mMol), was dissolved in 1.0 mL of 1,4-dioxane under an atmosphere of dry N₂. To this solution were added 4-hydroxyphenyl boronic acid
15 tetrahydropyranyl ether (67 mg, 0.31 mMol), potassium phosphate (108 mg, 0.508 mMol) and tetrakis(triphenylphosphine)palladium (0) (15 mg, 0.013 mMol). The reaction mixture was heated to 100°C and reacted overnight at this temperature in a seal tube. The reaction mixture was then cooled to ambient temperature, concentrated under vacuum and the residue was taken up in DCM. The resulting heterogeneous solution was filtered and the filter cake
20 was washed several times with DCM/MeOH (~1:1). The filtrates were combined and concentrated under vacuum and the resulting residue was chromatographed on flash silica gel eluting with MeOH/DCM/NH₄OH (1.5/98.3/0.2) to give 128 mg of the title compound **116C**.

Example 116D

- 25 4-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol.

- [3-(1-{8-[4-(Tetrahydro-pyran-2-yloxy)-phenyl]-quinolin-2-yl}-1H-benzoimidazol-5-yloxy)-propyl]-carbamic acid tert-butyl ester **116C** (128 mg, 0.215 mMol) was dissolved in 1 mL of TFA under an atmosphere of dry N₂ and was then stirred for 10 minutes. The reaction mixture
30 was concentrated under vacuum and the resulting residue was partitioned between isopropanol/DCM (18/82) and aqueous saturated NaHCO₃. The organic layer was dried over Na₂SO₄, filtered and concentrated under vacuum to give a brown oil. The oil was purified via High Pressure Liquid Chromatography (C-8 reverse phase; linear gradient from 0.01% HCl in water to 0.01% HCl /acetonitrile (over 15 minutes) to give 19 mg of the title compound **116** as
35 a white solid which was further purified by recrystallizing from EtOH to give 9 mg of the title compound as the bis-HCl salt.


C.I. m/z 411 [M+1]; ¹H NMR (CD₃OD) δ 9.75 (s, 1 H), 8.61 (d, J = 8.7 Hz, 1 H), 8.19 (d, J = 9.1 Hz, 1 H), 8.03 (d, J = 8.7 Hz, 1 H), 7.95 (m, 1 H), 7.80 (d, J = 6.2 Hz, 1 H), 7.70 (m, 1 H), 7.45

- 5 (d, J = 8.3 Hz, 2 H), 7.30 (m, 1 H), 6.95 (m, 1 H), 6.94 (d, J = 8.3 Hz, 2 H), 4.20 (m, 2 H), 3.29 (m, 2 H), 2.20 (m, 2 H).

10

Example 117

4-{2-[5-(3-Dimethylamino-propoxy)-benzimidazol-1-yl]-quinolin-8-yl}-phenol.




- The same procedure that was used in example 4 was followed except that 4-{2-[5-(3-amino-propoxy)-benzimidazol-1-yl]-quinolin-8-yl}-phenol **116** was used in the place of 1-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine **1** to give the title compound **117**.

- C.I. m/z 439 [M+1]; ¹H NMR (CD₃OD) δ 8.82 (s, 1 H), 8.33 (d, J = 8.7 Hz, 1 H), 7.99 (d, J = 9.1 Hz, 1 H), 7.80 (m, 2 H), 7.66 (m, 1 H), 7.52 (m, 1 H), 7.39 (d, J = 8.3 Hz, 2 H), 7.07 (d, J = 2.5 Hz, 1 H), 6.90 (d, J = 8.3 Hz, 2 H), 6.65 (dd, J = 2.5, 9.1 Hz, 1 H), 3.99 (m, 2 H), 2.75 (m, 2 H), 2.46 (s, 6 H), 2.00 (m, 2 H).

Example 118

1-[2-(5-Phenyl-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.



25

5

Example 118A

Trifluoro-methanesulfonic acid 1-[8-(4-tert-butoxycarbonylamino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl ester.

10 {1-[2-(5-Hydroxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester **3A** (2.37 g, 5.17 mMol) was dissolved in 35 mL of anhydrous THF under an atmosphere of dry N₂. To this solution were added triethylamine (790 µL, 5.68 mMol) followed by N-phenyl-bis(trifluoromethanesulfonamide) (2.02 g, 5.68 mMol). The reaction mixture was reacted at this temperature for 2 days. The yellow precipitate was collected via suction filtration, washed with isopropylether and dried under vacuum to give 1.48 g of the title
15 compound as a yellow solid.

Example 118B

{1-[2-(5-Phenyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester.

20 Trifluoro-methanesulfonic acid 1-[8-(4-tert-butoxycarbonylamino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl ester **118A** (150 mg, 0.253 mMol), was dissolved in 2.0 mL of 1,4-dioxane under an atmosphere of dry N₂. To this solution were added phenylboronic acid (46 mg, 0.38 mMol), potassium phosphate (161 mg, 0.759 mMol) and tetrakis(triphenylphosphine)palladium (0) (15 mg, 0.013 mMol). The reaction mixture was
25 heated to 100°C and reacted overnight at this temperature. The reaction mixture was then cooled to ambient temperature, concentrated under vacuum and the residue was partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was washed again with aqueous saturated NaHCO₃, dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow oil. The resulting oil was chromatographed on flash silica gel eluting with
30 EtOAc/Hexanes (70:30) to give 121 mg of the title compound **118B**.

Example 118C

1-[2-(5-Phenyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.


35 {1-[2-(5-Phenyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester **118B** (121 mg, 0.233 mMol) was dissolved in 0.50 mL of TFA under an atmosphere of dry N₂ and was then stirred for 45 minutes. The reaction mixture was concentrated under vacuum and the resulting residue was partitioned between DCM and aqueous 1 N NaOH. The DCM

- 5 layer was washed again with concentrated under vacuum and the resulting residue was partitioned between DCM and aqueous, dried over Na_2SO_4 , filtered and concentrated under vacuum to give a yellow oil as the title compound **118**. Compound **118** was dissolved in DCM and then 3 mL of 1 N HCl in ethyl ether was added to the mixture. The resulting precipitate was collected via suction filtration and dried under vacuum to give 91 mg of the title
10 compound **118** as the bis-HCl salt.

C.I. m/z 420 [M+1]; ^1H NMR (CD_3OD) δ 10.49 (s, 1 H), 8.84 (d, J = 8.7 Hz, 1 H), 8.80 (d, J = 9.1 Hz, 1 H), 8.27 (d, J = 8.7 Hz, 1 H), 8.16 (m, 2 H), 7.91 (m, 1 H), 7.80 (m, 4 H), 7.53 (m, 2 H), 7.44 (m, 1 H), 4.12 (m, 2 H), 3.50 (m, 1 H), 3.38 (m, 2 H), 2.36 (m, 4 H).

Example 119

- 15 1-[2-(5-Pyridin-4-yl-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.




The same procedure that was used in example 118 was followed except that 4-pyridylboronic acid was used in the place of phenylboronic acid in step 118B to give the title compound **119** as the free base.

- 20 C.I. m/z 421 [M+1]; ^1H NMR (CD_3OD) δ 9.00 (s, 1 H), 8.80 (d, J = 8.5 Hz, 1 H), 8.52 (d, J = 6.0 Hz, 2 H), 8.33 (d, J = 8.7 Hz, 1 H), 8.03 (s, 1 H), 7.84 (m, 2 H), 7.72 (d, J = 6.0 Hz, 2 H), 7.44 (m, 2 H), 7.27 (d, J = 8.5 Hz, 1 H), 3.78 (m, 2 H), 2.84 (m, 1 H), 2.76 (m, 2 H), 1.99 (m, 2 H), 1.85 (m, 2 H).

25

Example 120

- 1-[2-[5-(3-Methoxy-phenyl)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine.




- 5 The same procedure that was used in example 118 was followed except that (3-methoxyphenyl)boronic acid was used in the place of phenylboronic acid in example 118B to give the title compound 120 as the free base.

C.I. m/z 450 [M+1]; ¹H NMR (CDCl₃) δ 8.74 (s, 1 H), 8.46 (d, J = 8.5 Hz, 1 H), 8.30 (d, J = 8.7 Hz, 1 H), 8.08 (d, J = 1.3 Hz, 1 H), 7.69 (d, J = 8.7 Hz, 1 H), 7.66 (dd, J = 1.7, 8.5 Hz, 1 H),
10 7.46 (m, 2 H), 7.38 (m, 1 H), 7.22-28 (m, 3 H), 6.90 (m, 1 H), 3.89 (m, 2 H), 3.87 (s, 3 H), 2.90 (m, 3 H), 2.05 (m, 2 H), 1.83 (m, 2 H).

Example 121

1-[2-(5-Pyridin-3-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine.



15


- The same procedure that was used in example 77 was followed except that {1-[2-(5-phenyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester **118B** was used in the place of trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl ester **77E** in example 77F. The resulting compound was deprotected using the
20 procedure outlined in example 118C to give the title compound **121** as the free base.

C.I. m/z 421 [M+1]; ¹H NMR (CD₃OD) δ 9.32 (s, 1 H), 9.27 (s, 1 H), 8.93 (m, 1 H), 8.87 (d, J = 8.7 Hz, 1 H), 8.82 (d, J = 5.7 Hz, 1 H), 8.55 (d, J = 8.9 Hz, 1 H), 8.24 (s, 1 H), 8.13 (m, 1 H), 8.03 (m, 2 H), 7.70 (d, J = 8.1 Hz, 1 H), 7.57 (m, 1 H), 7.47 (d, J = 7.7 Hz, 1 H), 4.01 (m, 2 H), 3.38 (m, 1 H), 3.05 (m, 2 H), 2.13-2.24 (m, 4 H).

25

Example 122

1-[2-[5-(6-Methoxy-pyridin-3-yl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine.




- 5 The same procedure that was used in example 118 was followed except that 2-methoxy-5-pyridineboronic acid was used in the place of phenylboronic acid in example 118B to give the title compound 122 as the HCl salt.

C.I. m/z 451 [M+1]; ¹H NMR (DMSO) δ 9.27 (s, 1 H), 9.02 (d, J = 8.7 Hz, 1 H), 8.61 (d, J = 1.5 Hz, 1 H), 8.57 (d, J = 9.1 Hz, 1 H), 8.19 (d, J = 8.7 Hz, 1 H), 8.15 (m, 1 H), 8.07 (s, 1 H), 8.02 (d, J = 8.7 Hz, 1 H), 7.63 (d, J = 7.9 Hz, 1 H), 7.50 (m, 1 H), 7.31 (d, J = 7.5 Hz, 1 H), 6.92 (d, J = 8.3 Hz, 1 H), 3.89 (s, 3 H), 3.87 (m, 2 H), 3.22 (m, 1 H), 2.79 (m, 2 H), 2.07 (m, 4 H).

Example 123

1-{2-[5-(4-Aminomethyl-phenyl)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.



15


The same procedure that was used in example 118 was followed except that the HCl salt of 4-aminomethylphenylboronic acid was used in the place of phenylboronic acid in example 118B and 2.5 equivalents of N,N-diisopropylethylamine was added to the reaction mixture to give the title compound 123 as the HCl salt.

- 20 C.I. m/z 449 [M+1]; ¹H NMR (CD₃OD) δ 9.10 (s, 1 H), 8.71 (d, J = 8.7 Hz, 1 H), 8.55 (d, J = 8.7 Hz, 1 H), 8.04 (m, 2 H), 7.87 (dd, J = 1.7, 8.7 Hz, 1 H), 7.83 (m, 2 H), 7.66 (d, J = 7.1 Hz, 1 H), 7.58 (m, 3 H), 7.40 (m, 1 H), 4.18 (s, 2 H), 4.03 (m, 2 H), 3.34 (m, 1 H), 2.95 (m, 2 H), 2.20 (m, 2 H), 2.10 (m, 2 H).

25

Example 124

4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazol-5-yl}-benzoic acid methyl ester.




- 5 The same procedure that was used in example 118 was followed except that 4-methoxycarbonylphenylboronic acid was used in the place of phenylboronic acid in example 118B to give the title compound **124** as the HCl salt.

C.I. m/z 478 [M+1]; ¹H NMR (CD₃OD) δ 9.08 (s, 1 H), 8.79 (d, J = 8.7 Hz, 1 H), 8.52 (d, J = 8.7 Hz, 1 H), 8.11 (m, 2 H), 8.07 (d, J = 1.3 Hz, 1 H), 8.02 (d, J = 9.1 Hz, 1 H), 7.87 (m, 1 H), 7.83 (m, 2 H), 7.65 (m, 1 H), 7.55 (m, 1 H), 7.39 (m, 1 H), 4.03 (m, 2 H), 3.93 (s, 3 H), 3.33 (m, 1 H), 2.93 (m, 2 H), 2.19 (m, 2 H), 2.11 (m, 2 H).

Example 125

4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazol-5-yl}-phenol.



15


The same procedure that was used in example 118 was followed except that 4-hydroxyphenylboronic acid tetrahydropyranyl ether was used in the place of phenylboronic acid in example 118B to give the title compound **125** as the HCl salt.

C.I. m/z 436 [M+1]; ¹H NMR (CD₃OD) δ 9.20 (s, 1 H), 8.72 (d, J = 8.3 Hz, 1 H), 8.49 (d, J = 8.7 Hz, 1 H), 7.98 (d, J = 8.7 Hz, 1 H), 7.90 (s, 1 H), 7.79 (m, 1 H), 7.64 (d, J = 7.5 Hz, 1 H), 7.53 (m, 3 H), 7.39 (d, J = 7.9 Hz, 1 H), 6.88 (m, 2 H), 4.03 (m, 2 H), 3.34 (m, 1 H), 2.93 (m, 2 H), 2.19 (m, 2 H), 2.10 (m, 2 H).

20

Example 126

25 2-(5-Methoxy-benzimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester.



Trifluoro-methanesulfonic acid 2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl ester **1E** (1.0 g, 2.3 mMol), palladium acetate (16 mg, 0.07 mMol), 1,3-bis(diphenylphosphine)propane (30 mg, 0.07 mMol) and triethylamine (730 μL, 5.20 mMol) were dissolved in a solution of 8 mL of DMF and 4 mL of MeOH in a Paar bottle. The reaction vessel was evacuated under vacuum

30


5 and then filled with carbon monoxide (50 psi). The reaction mixture was shaken for 24 hours after which time it was concentrated under vacuum. The resulting residue was partitioned between DCM and aqueous saturated NaHCO_3 . The DCM layer was then washed again with aqueous saturated NaHCO_3 and then with brine, dried over Na_2SO_4 , filtered and concentrated under vacuum to give a light tan foam. The foam was chromatographed on flash silica gel
10 eluting with MeOH/DCM (1:99) to give 770 mg of the title compound **126**.

C.I. m/z 334 $[\text{M}+1]$; ^1H NMR (CDCl_3) δ 8.63 (d, $J = 8.9$ Hz, 1 H), 8.56 (s, 1 H), 8.25 (d, $J = 8.9$ Hz, 1 H), 8.10 (dd, $J = 1.5, 7.2$ Hz, 1 H), 7.90 (dd, $J = 1.5, 8.1$ Hz, 1 H), 7.64 (d, $J = 8.9$ Hz, 1 H), 7.51 (m, 1 H), 7.27 (d, $J = 2.5$ Hz, 1 H), 7.07 (dd, $J = 2.5, 9.1$ Hz, 1 H), 4.04 (s, 3 H), 3.87 (s, 3 H).

15

Example 127

2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinoline-8-carboxylic acid methyl ester.




The same procedure that was used in example 126 was followed except that trifluoro-methanesulfonic acid 2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl ester **77E**
20 was used in the place of trifluoro-methanesulfonic acid 2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl ester **1E** to give the title compound **127** as a white solid.

C.I. m/z 378 $[\text{M}+1]$; ^1H NMR (DMSO) δ 9.17 (s, 1 H), 8.73 (d, $J = 9.1$ Hz, 1 H), 8.64 (d, $J = 9.1$ Hz, 1 H), 8.23 (d, $J = 8.7$ Hz, 1 H), 8.18 (m, 1 H), 8.05 (m, 1 H), 7.64 (m, 1 H), 7.29 (d, $J = 2.5$ Hz, 1 H), 7.07 (dd, $J = 2.5, 9.1$ Hz, 1 H), 4.14 (m, 2 H), 4.00 (s, 3 H), 3.67 (m, 2 H), 3.31 (s, 3 H).
25

Example 128

2-(5-Methoxy-benzimidazol-1-yl)-quinoline-8-carboxylic acid (2-dimethylamino-ethyl)-amide.

-143-



5

2-(5-Methoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester **126** (166.7 mg, 0.500 mMol) was dissolved in 2 mL of anhydrous DCM under an atmosphere of dry N₂. In a separate flask, N,N-dimethylethylenediamine (2.20 mL, 20 mMol) was dissolved in 20 mL of anhydrous DCM under an atmosphere of dry N₂. To this solution was slowly added 10 mL of a solution of 2.0 M trimethylaluminum in toluene. The solution was stirred for 30 minutes after which time 1 mL of this solution was added to the solution containing compound **126**. The temperature of the reaction mixture was raised to 40°C and reacted over night after which time 1 mL of water was slowly added to quench the reaction. The resulting mixture was partitioned between DCM and aqueous 0.1 N NaOH. The DCM layer was washed again with aqueous 0.1 N NaOH and then with brine. The DCM layer was then dried over Na₂SO₄, filtered and concentrated under vacuum. The resulting residue was dissolved in 2 mL of DCM. To this solution was added 1.5 mL of a solution of HCl in ethyl ether which caused an immediate precipitate to form. The precipitate was collected via suction filtration, washed with ethyl ether, and dried under vacuum to give 165 mg of the bis-HCl salt of the title compound **128** as a white solid.


15
20

C.I. m/z 390 [M+1]; ¹H NMR (CD₃OD) δ 10.22 (s, 1 H), 8.85 (d, J = 8.9 Hz, 1 H), 8.54 (d, J = 9.0 Hz, 1 H), 8.49 (dd, J = 1.5, 7.2 Hz, 1 H), 8.30 (dd, J = 1.4, 8.1 Hz, 1 H), 8.18 (d, J = 8.9 Hz, 1 H), 7.86 (m, 1 H), 7.42 (m, 2 H), 3.97 (s, 3 H), 3.90 (m, 2 H), 3.44 (m, 2 H), 2.96 (s, 6 H).

25

Example 129

2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester.




- 5 The same procedure that was used in example 126 was followed except that 4-cyclopropylmethoxy-2-nitro-phenylamine **52A** was used in the place of 4-methoxy-2-nitroaniline in example 1B to give the titled compound **129**.

C.I. m/z 374 [M+1]; ¹H NMR (CDCl₃) δ 8.65 (d, J = 9.1 Hz, 1 H), 8.60 (s, 1 H), 8.34 (d, J = 8.7 Hz, 1 H), 8.14 (m, 1 H), 7.97 (m, 1 H), 7.72 (d, J = 9.1 Hz, 1 H), 7.57 (m, 1 H), 7.29 (d, J = 2.5 Hz, 1 H), 7.10 (m, 1 H), 4.09 (s, 3 H), 3.89 (d, J = 7.1 Hz, 2 H), 1.32 (m, 1 H), 0.66 (m, 2 H), 0.38 (m, 2 H).

Example 130

[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-pyrrolidin-1-yl-methanone.



15

- The same procedure that was used in example 128 was followed except that 2-(5-cyclopropylmethoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester **129** was used in the place of 2-(5-methoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester **126** and pyrrolidine was used in the place of N,N-dimethylethylenediamine to give the title compound **130** as a white solid.

20


C.I. m/z 413 [M+1]; ¹H NMR (CDCl₃) δ 8.58 (s, 1 H), 8.51 (d, J = 9.1 Hz, 1 H), 8.26 (d, J = 9.1 Hz, 1 H), 7.80 (d, J = 7.9 Hz, 1 H), 7.76 (d, J = 7.1 Hz, 1 H), 7.67 (d, J = 8.7 Hz, 1 H), 7.52 (m, 1 H), 7.25 (d, J = 2.1 Hz, 1 H), 7.07 (m, 1 H), 3.92 (m, 1 H), 3.88 (d, J = 7.1 Hz, 2 H), 3.85 (m, 1 H), 3.19 (m, 1 H), 3.10 (m, 1 H), 1.95 (m, 2 H), 1.75 (m, 2 H), 1.32 (m, 1 H), 0.67 (m, 2 H), 0.39 (m, 2 H).

25

Example 131

[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-morpholin-4-yl-methanone.

-145-



5


The same procedure that was used in example 128 was followed except that 2-(5-cyclopropylmethoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester **129** was used in the place of 2-(5-methoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester **126** and morpholine was used in the place of N,N-dimethylethylenediamine to give the

10

C.I. m/z 429 [M+1]

Example 132

[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-1-yl-methanone.



15 The same procedure that was used in example 128 was followed except that 2-(5-cyclopropylmethoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester **129** was used in the place of 2-(5-methoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester **126** and piperidine was used in the place of N,N-dimethylethylenediamine to give the title compound **132** as a yellow solid.


20 C.I. m/z 427 [M+1]; ¹H NMR (CDCl₃) δ 8.61 (s, 1 H), 8.49 (d, J = 9.1 Hz, 1 H), 8.31 (d, J = 9.1 Hz, 1 H), 7.85 (d, J = 7.9 Hz, 1 H), 7.72 (m, 2 H), 7.56 (m, 1 H), 7.29 (d, J = 2.5 Hz, 1 H), 7.09 (m, 1 H), 4.09 (m, 1 H), 3.88 (d, J = 6.7 Hz, 2 H), 3.78 (m, 1 H), 3.13 (m, 2 H), 1.20-1.80 (m, 7 H), 0.66 (m, 2 H), 0.39 (m, 2 H).

25

Example 133

(3-Amino-pyrrolidin-1-yl)-[2-(5-cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-methanone.

-146-



5

Example 133A

4-Cyclopropylmethoxy-2-nitro-phenylamine.

- 10 4-Amino-3-nitrophenol (26.00 g, 165.5 mMol) was dissolved in 200 mL of anhydrous DMF under an atmosphere of dry N₂. The reaction mixture was then cooled to 0°C and to this mixture was added Cs₂CO₃ (64.7 g, 199 mMol), and cyclopropyl methane bromide (17.7 mL, 182 mMol). After stirring for 15 minutes, the reaction mixture was then warmed up to ambient temperature and then stirred overnight. The reaction mixture is then poured into 800 mL of
- 15 water. The precipitate was collected via suction filtration and partitioned between DCM and aqueous 0.1 N NaOH. The DCM layer was saved and washed again aqueous 0.1 N NaOH, then with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give 31.52 g of 4-cyclopropylmethoxy-2-nitro-phenylamine **133A** as an orange solid.

20

Example 133B

{1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinoline-8-carbonyl]-pyrrolidin-3-yl}-
carbamic acid tert-butyl ester.

- Trifluoro-methanesulfonic acid 2-(5-cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl ester (602 mg, 1.30 mMol), (which was prepared by following example 77E except that 4-cyclopropylmethoxy-2-nitro-phenylamine **133A** was used in the place of 5-(2-methoxy-ethoxy)-2-nitro-phenylamine **42A** in example 77C) palladium acetate (9.0 mg, 0.04 mMol), 1,3-bis(diphenylphosphine)propane (16 mg, 0.04 mMol), (+/-)-3-(tert-butoxycarbonylamino)pyrrolidine (484 mg, 2.60 mMol) and triethylamine (400 µL, 2.86 mMol) were dissolved in 6 mL of DMF and 4 mL of MeOH in a Paar bottle. The reaction vessel was
- 25 evacuated under vacuum and then filled with carbon monoxide (50 psi). The reaction mixture was shaken for 24 hours after which time it was concentrated under vacuum. The resulting residue was partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was then washed again with aqueous saturated NaHCO₃ and then with brine, dried over Na₂SO₄,
- 30

-147-

- 5 filtered and concentrated under vacuum. The resulting residue was chromatographed on flash silica gel eluting with MeOH/DCM (1:99) to give 139 mg of the title compound **133B**.

Example 133C


- 10 (3-Amino-pyrrolidin-1-yl)-[2-(5-cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-methanone.

- {1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinoline-8-carbonyl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester **133B** (130 mg, 0.246 mMol) was dissolved in a solution of 1 mL of TFA and 1 mL of DCM under an atmosphere of dry N₂. The reaction mixture was stirred for 15 30 minutes. The reaction mixture was concentrated under vacuum and the resulting residue was partitioned between DCM and aqueous 1 N NaOH. The DCM layer was washed again with concentrated under vacuum and the resulting residue was partitioned between DCM and aqueous, dried over Na₂SO₄, filtered and concentrated under vacuum to give the title compound **133**.

- 20 C.I. m/z 428 [M+1]

Example 134

8-Allyloxy-2-(5-methoxy-benzoimidazol-1-yl)-quinoline.




- 25 2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-ol **1D** (292 mg, 1.00 mMol) was dissolved in 5 mL of anhydrous THF under an atmosphere of dry N₂ and the reaction mixture was cooled to 0°C. To this solution was slowly added 60% sodium hydride in oil (44 mg, 1.1 mMol. After reacting for 30 minutes, allyl bromide (100 μL, 1.1 mMol) was added and the reaction mixture was then warmed to ambient temperature and 1 mL of DMF was added. After stirring 30 overnight at ambient temperature, the reaction mixture was partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was washed again with aqueous saturated NaHCO₃, then with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give a red solid. The red solid was chromatographed over flash silica gel eluting with EtOAc to give 245 mg of the title compound **134** as a tan solid.

- 5 C.I. m/z 332 [M+1]; ¹H NMR (CDCl₃) δ 8.60 (d, J = 8.9 Hz, 1 H), 8.57 (s, 1 H), 8.19 (d, J = 8.7 Hz, 1 H), 7.59 (d, J = 8.7 Hz, 1 H), 7.37 (m, 2 H), 7.30 (d, J = 2.5 Hz, 1 H), 7.07 (dd, J = 1.9, 7.0 Hz, 1 H), 7.02 (dd, J = 2.5, 9.1 Hz, 1 H), 6.21 (m, 1 H), 5.62 (m, 1 H), 5.36 (m, 1 H), 4.76 (m, 2 H), 3.87 (s, 3 H).

10

Example 135

{2-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yloxy]-ethyl}-methyl-amine.

Example 135A

15

[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yloxy]-acetaldehyde.

- 8-Allyloxy-2-(5-methoxy-benzoimidazol-1-yl)-quinoline **134** (245 mg, 0.74 mMol), and trimethylamine-N-oxide dihydrate (101 mg, 0.88 mMol) were dissolved in 3 mL of anhydrous DCM under an atmosphere of dry N₂. To this solution was added osmium tetroxide (245 mg, 0.74 mMol) and the reaction mixture was stirred for ~ 1 hour at ambient temperature. The reaction mixture was then concentrated under vacuum and the resulting residue was dissolved in 2 mL of THF. To this solution was added 2 mL of water followed by sodium periodate (238 mg, 1.11 mMol) and the reaction mixture was stirred overnight at ambient temperature. The reaction mixture was partitioned between DCM and aqueous saturated NaHCO₃ causing a precipitate to form. The brown solid was collected by suction filtration and dried under vacuum to give 214 mg of the title compound **135A**.

Example 135B

{2-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yloxy]-ethyl}-methyl-amine.

30


[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yloxy]-acetaldehyde **135A** (214 mg, 0.64 mMol), 400 μL of AcOH, 1.6 mL of a solution of 2 M methylamine in MeOH and Na(OAc)₃BH (204,

- 5 0.96 mMol) were added to 3 mL of DCE under an atmosphere of dry N₂. The reaction mixture was stirred overnight at ambient temperature after which time it was partitioned between DCM and aqueous 1 N NaOH. The DCM layer was washed 2 more times with aqueous 1 N NaOH, then with brine, dried over Na₂SO₄, filtered and concentrated under vacuum. The resulting residue was chromatographed on flash silica gel initially eluting with MeOH/EtOAc (4:96),
10 then MeOH/EtOAc/NH₄OH (8:91.9:0.1), then MeOH/EtOAc/NH₄OH (10:89.9:0.1) and finally MeOH/EtOAc/NH₄OH (15:84.9:0.1) to give 28 mg of the title compound **135** as a white solid.

C.I. m/z 349 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (d, J = 9.1 Hz, 1 H), 8.60 (s, 1 H), 8.27 (d, J = 8.7 Hz, 1 H), 7.66 (d, J = 8.9 Hz, 1 H), 7.43 (m, 2 H), 7.33 (d, J = 2.5 Hz, 1 H), 7.14 (dd, J = 2.1, 6.9 Hz, 1 H), 7.05 (dd, J = 2.5, 8.9 Hz, 1 H), 4.35 (m, 2 H), 3.89 (s, 3 H), 3.21 (m, 2 H), 2.61
15 (s, 3 H), 2.02 (brs, 1 H).

Example 136

{2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethyl}-dimethyl-amine.



- 20 {2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethyl}-methyl-amine **135** (23 mg, 0.066 mMol), 20 μL of 37% formaldehyde in water and 20 μL of formic acid were added to 500 μL of chloroform under an atmosphere of dry N₂. The reaction mixture was then heated to reflux for ~ 2 hours after which time it was concentrated under vacuum and partitioned between DCM and aqueous 1 N NaOH. The DCM layer was washed again with aqueous 1 N NaOH, then
25 with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to give 20 mg of the title compound **136**.


C.I. m/z 363 [M+1]; ¹H NMR (CDCl₃) δ 8.67 (d, J = 9.1 Hz, 1 H), 8.61 (s, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.67 (d, J = 8.9 Hz, 1 H), 7.43 (m, 2 H), 7.32 (d, J = 2.3 Hz, 1 H), 7.13 (d, J = 7.3 Hz, 1 H), 7.06 (dd, J = 2.5, 8.9 Hz, 1 H), 4.34 (m, 2 H), 3.90 (s, 3 H), 3.00 (m, 2 H), 2.46 (s, 6 H).

30

Example 137

2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethylamine.

-150-




5
2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-ol **1D** (290 mg, 0.996 mMol) was dissolved in 5 mL of anhydrous DMF under an atmosphere of dry N₂. To this solution was slowly added 60% sodium hydride in oil (44.1 mg, 1.10 mMol). After reacting for 30 minutes, N-(2-bromoethyl)-phthalimide (280 mg, 1.10 mMol) was added and the reaction mixture was then
10 heated to 80°C. After stirring for 2 hours at this temperature, K₂CO₃ (360 mg, 2.61 mMol) was added and the reaction mixture was stirred overnight at 80°C. The reaction mixture was then cooled to ambient temperature and partitioned between EtOAc and water. The EtOAc layer was then washed 4 more times with water, then with brine, dried over MgSO₄, filtered and concentrated under vacuum. The resulting residue was chromatographed over flash silica gel
15 eluting with a gradient from DCM/Hex (50:50) to DCM and then to MeOH/DCM (2:98) to give a white solid. The white solid was dissolved in 5 mL of hot (reflux) EtOH to which was added 500 µL of anhydrous hydrazine. After reacting at reflux for two hours, the reaction mixture was cooled to ambient temperature, filtered and the filter cake was washed with EtOH. The combined filtrates were concentrated under vacuum and the resulting residue was
20 chromatographed over flash silica gel eluting with a gradient of DCM to MeOH/DCM (8:92) then switched to MeOH/NH₄OH (8:91.9: 0.1) to give 110 mg of the title compound **137** as a white solid.

¹H NMR (CDCl₃) δ 8.60 (d, J = 8.9 Hz, 1 H), 8.58 (d, J = 8.9 Hz, 1 H), 8.57 (s, 1 H), 8.21 (d, J = 8.7 Hz, 1 H), 7.60 (d, J = 8.7 Hz, 1 H), 7.39 (m, 2 H), 7.29 (d, J = 2.5 Hz, 1 H), 7.08 (dd, J =
25 1.9, 7.1 Hz, 1 H), 7.03 (dd, J = 2.5, 8.9 Hz, 1 H), 4.21 (m, 2 H), 3.86 (s, 3 H), 3.27 (m, 2 H), 2.26 (brs, 2 H).

Example 138

1-[[2-[5-(3-Morpholinoethoxy)-1H-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride.
30



5

Example 138A,B

8-Chloro-2-(5-methoxy-1*H*-benzimidazol-1-yl)-quinoline and 8-Chloro-2-(6-methoxy-1*H*-benzimidazol-1-yl)-quinoline.

- 10 2,8-Dichloroquinoline (11.89 g, 60 mMol), sodium hydride (60% dispersion in mineral oil, 2.88 g, 72 mMol) and 50 mL of toluene were combined under nitrogen, and a solution containing 5-methoxy-benzimidazole (10.67 g, 72 mMol) dissolved in 100 mL of anhydrous 1-methyl-2-pyrrolidinone was carefully added by syringe. The resultant mixture was warmed slowly to 110° C and stirred at this temperature for 16 hrs. After cooling, the reaction mixture was quenched
- 15 with 650 mL of water and then extracted multiple times with EtOAc. The combined EtOAc extracts were dried over Na₂SO₄ and evaporated to leave a semi-solid residue containing the title product isomers **138A,B**. An ¹H NMR spectrum of this product mixture revealed that formation of the 6-methoxybenzimidazol-1-yl isomer was favored over the 5-methoxybenzimidazol-1-yl isomer by about a 3:2 ratio. Fractional crystallization from EtOH
- 20 provided three early crystalline crops that were combined to give 7.34 g, comprised of about 90% pure 6-methoxybenzimidazol-1-yl isomer. Recrystallization from EtOH gave needle-like crystals of the 6-methoxybenzimidazol-1-yl isomer **138A** melting at 175-176° C.

C.I. m/z 310 [M+1]; ¹H NMR (DMSO) δ 9.18 (s, 1 H), 8.77 (d, J = 2.5 Hz, 1 H), 8.65 (d, J = 9.0 Hz, 1 H), 8.30 (d, J = 9.0 Hz, 1 H), 8.01 (m, 2 H), 7.68 (d, 8.7 Hz, 1 H), 7.58 (m, 1 H), 7.00 (dd, J = 2.5, 8.7 Hz, 1 H), 3.92 (s, 3 H).

25

Additional fractional crystallization efforts ultimately yielded two crops, 2.157 g combined weight, of the 5-methoxybenzimidazol-1-yl isomer **138B**, melting at 184-185° C.

¹H NMR (DMSO) δ 9.27 (s, 1 H), 8.98 (d, J = 9.0 Hz, 1 H), 8.67 (d, J = 9.0 Hz, 1 H), 8.31 (d, J = 9.0 Hz, 1 H), 8.02 (m, 2 H), 7.59 (m, 1 H), 7.35 (d, J = 2.2 Hz, 1 H), 7.12 (dd, J = 2.2, 9.0 Hz, 1 H), 3.87 (s, 3 H).

30

5

Example 138C

8-Chloro-2-(5-hydroxy-1*H*-benzimidazol-1-yl)-quinoline.

To a pre-cooled (-78°C) solution containing 8-chloro-2-(5-methoxy-1*H*-benzimidazol-1-yl)-quinoline **138B** (2.16 g, 7.0 mMol) in anhydrous DCM (28 mL) was added a 1.0 M solution of borontribromide in DCM (20.9 mL, 20.9 mMol). The cooling bath was removed and the
10 reaction mixture stirred at ambient temperature under nitrogen for 22 hours. The reaction was quenched by the addition of aqueous 1 N NaOH to a pH of about 8.8. After stirring for about 15 minutes, the suspended solids were removed by filtration. The solids were subsequently stirred briefly with 1 N NaOH (21 mL, 21 mMol) to hydrolyse remaining borate esters, and then neutralized by adding 1 N HCl (21 mL, 21 mMol). The solid precipitate was filtered off,
15 stirred briefly with a small amount of warm EtOAc to remove traces of starting material, and re-filtered to give 1.91 g, mp >250° C, of the title product **138C**.

C.I. m/z 296 [M+1]; ¹H NMR (DMSO) δ 9.41 (br. s, 1 H), 9.20 (s, 1 H), 8.86 (d, J = 9.1 Hz, 1 H), 8.65 (d, J = 9.1 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 8.00 (d, J = 7.9 Hz, 2 H), 7.57 (t, J = 7.9 Hz, 1 H), 7.08 (d, J = 2.5 Hz, 1 H), 6.93 (dd, J = 2.5, 8.7 Hz, 1 H).

20

Example 138D

8-Chloro-2-[5-(3-morpholinopropoxy)-1*H*-benzimidazol-1-yl]-quinoline.

8-Chloro-2-(5-hydroxy-1*H*-benzimidazol-1-yl)-quinoline **138C** (0.30 g, 1 mMol) and NaH (0.047 g of 60% dispersion in mineral oil, 1.1 mMol) were combined with 3 mL of anhydrous
25 DMF under argon, and the mixture allowed to react for 1 hr at ambient temperature. To this was added 4-(3-chloropropyl)morpholine (0.199 g, 1.22 mMol) and the reaction mixture stirred at ambient temperature for 4 days. Remaining sodium hydride, if any, was quenched by water addition and the solvents evaporated under vacuum. Water (3 mL) was added to the residue,
30 and the mixture extracted (5 x 20 mL) with chloroform. The combined chloroform extracts were dried over Na₂SO₄ and evaporated to give an off-white solid. The solids were stirred with a small amount of EtOH and filtered to give 0.157 g of title product **138D**.

C.I. m/z 423 [M+1]; ¹H NMR (DMSO) δ 9.24 (s, 1 H), 8.95 (d, J = 9.1 Hz, 1 H), 8.66 (d, J = 9.1 Hz, 1 H), 8.30 (d, J = 9.1 Hz, 1 H), 8.01 (d, J = 7.9 Hz, 2 H), 7.57 (t, J = 7.9 Hz, 1 H), 7.30 (d, J = 2.1 Hz, 1 H), 7.08 (dd, J = 2.5, 9.1 Hz, 1 H), 4.08 (t, J = 6.4 Hz, 2 H), 3.56 (m, 4 H), 2.43 (t, J = 7.3 Hz, 2 H), 2.36 (m, 4 H), 1.89 (m, 2H).

Example 138E

- 5 {1-[[2-[5-(3-Morpholinopropoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-yl}-carbamic acid tert-butyl ester.

8-Chloro-2-[5-(3-morpholinopropoxy)-1*H*-benzimidazol-1-yl]-quinoline **138D** (226 mg, 0.53 mMol), racemic-BINAP (50 mg, 0.08 mMol), Cs₂CO₃ (243 mg, 0.75 mMol), piperidin-4-yl-carbamic acid tert-butyl ester (214 mg, 1.07 mMol), xylenes (1 mL), and Pd(OAc)₂ (12 mg, 0.053 mMol) were combined under argon and heated at reflux for 2 days. After allowing the mixture to cool, hexanes were added and decanted off, several times, to remove hexane solubles. The residue was stirred with EtOAc and filtered, and the solids washed with additional EtOAc. The combined EtOAc solutions were evaporated, and the residue
10 chromatographed on silica gel with EtOAc/MeOH elution. Fractions containing the title product **138E** were combined and concentrated to provide 130 mg.
C.I. m/z 587 [M+1]; ¹H NMR (DMSO) δ 9.15 (s, 1 H), 8.88 (d, J = 9.6 Hz, 1 H), 8.51 (d, J = 8.7 Hz, 1 H), 8.14 (d, J = 9.1 Hz, 1 H), 7.59 (d, J = 7.0 Hz, 1 H), 7.46 (t, J = 7.9 Hz, 1 H), 7.30 (m, 3 H), 7.13 (d, J = 8.3 Hz, 1 H), 4.08 (t, J = 6.2 Hz, 2 H), 3.75 (m, 2 H), 3.56 (m, 4 H), 3.42 (m, 1 H), 2.74 (t, J = 10.8
20 Hz, 2 H), 2.44 (m, 2 H), 2.36 (m, 4 H), 1.8-1.9 (m, 6 H), 1.39 (s, 9 H).

Example 138F

1-[[2-[5-(3-Morpholinopropoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride.

25

{1-[[2-[5-(3-Morpholinopropoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-yl}-carbamic acid tert-butyl ester **138E** (98 mg, 0.17 mMol) was stirred at ambient temperature with 4N HCl in dioxane (0.5 mL, 2 mMol). After 4 hr the mixture was rotoevaporated to provide the title product **138** as a pale yellow solid.


30 C.I. m/z 487 [M+1]; ¹H NMR (DMSO) δ 9.55 (brs, 1 H), 8.94 (d, J = 8.7 Hz, 1 H), 8.65 (d, J = 9.0 Hz, 1 H), 8.4 (brs, 3 H), 8.24 (d, J = 9.0 Hz, 1 H), 7.72 (d, J = 8.1 Hz, 1 H), 7.58 (m, 2 H), 7.42 (m, 2 H), 4.24 (t, J = 5.6 Hz, 2 H), 4.0 (m, 2 H), 3.9 (m, 4 H), 3.5 (m, 2 H), 3.3 (m, 3 H), 3.1 (m, 2 H), 2.88 (m, 2 H), 2.29 (m, 2 H), 2.12 (m, 4 H).

35

Example 139

1-[[2-[5-(3-Morpholinoethoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride.

-154-



5

Example 139A**8-Chloro-2-[5-(3-morpholinoethoxy)-1H-benzimidazol-1-yl]-quinoline.**

- 10 8-Chloro-2-(5-hydroxy-1H-benzimidazol-1-yl)-quinoline **138B** (296 mg, 1 mMol) and Cs₂CO₃ (717 mg, 2.2 mMol) and 3 ml anhydrous dioxane were combined and heated at 80° C for 1.5 hr under nitrogen. Morpholinoethylchloride hydrochloride (223 mg, 1.2 mMol) was added and heating continued at 80° C overnight. Additional morpholinoethylchloride hydrochloride (112 mg, 0.6 mMol) and Cs₂CO₃ (358 mg, 1.1 mMol) were added, and the mixture heated for an
- 15 additional 2 days. After evaporation of the solvent, water (2 mL) was added to the residue, and the mixture extracted with EtOAc. The EtOAc extract was dried over Na₂SO₄ and evaporated to give 200 mg of title product **139A** as an off-white solid.

Example 139B

- 20 {1-[[2-[5-(3-Morpholinoethoxy)-1H-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-yl}-carbamic acid tert-butyl ester.

- 8-Chloro-2-[5-(3-morpholinoethoxy)-1H-benzimidazol-1-yl]-quinoline **139A** (182 mg, 0.32 mMol), racemic-BINAP (30 mg, 0.05 mMol), Cs₂CO₃ (145 mg, 0.45 mMol), piperidin-4-yl-carbamic acid tert-butyl ester (127 mg, 0.64 mMol), xylenes (1 mL), and Pd(OAc)₂ (7.1 mg, 0.032 mMol) were combined under argon and heated at reflux for 2 days. This reaction was
- 25 worked in the same way as in example 138E, including chromatography, to provide 120 mg of title product **139B**.

30

Example 139C


1-[[2-[5-(3-Morpholinoethoxy)-1H-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride.

- 5 {1-[[2-[5-(3-Morpholinoethoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl]-carbamic acid tert-butyl ester **139B** (121 mg, 0.21 mMol) was stirred at ambient temperature with 4*N* HCl in dioxane (0.63 mL, 2.5 mMol). After 4 hr the mixture was rotoevaporated, and the residue stirred briefly with fresh anhydrous dioxane. Filtration provided the title product **139** as a pale yellow solid.
- 10 C.I. *m/z* 473 [M+1]; ¹H NMR (DMSO) δ 9.67 (brs, 1 H), 8.92 (d, *J* = 9.1 Hz, 1 H), 8.62 (d, *J* = 8.7 Hz, 1 H), 8.50 (brs, 3 H), 8.21 (d, *J* = 8.7 Hz, 1 H), 7.76 (m, 1 H), 7.69 (d, *J* = 8.3 Hz, 1 H), 7.55 (t, *J* = 7.9 Hz, 1 H), 7.44 (d, *J* = 2.5 Hz, 1 H), 7.40 (m, 1H), 4.59 (m, 2 H), 3.97 (d, *J* = 11.2 Hz, 2 H), 3.84 (m, 4 H), 3.6-3.5 (m, 4 H), 3.21-3.24 (m, 3 H), 2.86 (m, 2 H), 2.09 (m, 4 H).

15

Example 140

5-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-[1,3,4]oxadiazol-2-ylamine.

Example 140A

20 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinoline-8-carboxylic acid hydrazide.

- 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinoline-8-carboxylic acid methyl ester **127** (1.00 g, 2.66 mMol) was dissolved in 10 mL of anhydrous THF under an atmosphere of dry N₂. To this solution was added 400 μL of anhydrous hydrazine and the solution was then
- 25 stirred for 18 hours at ambient temperature. The solution was concentrated under vacuum and the resulting residue was partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was washed again with aqueous saturated NaHCO₃, then with brine, dried over Na₂SO₄, filtered and concentrated under vacuum. The resulting residue was chromatographed on flash silica gel eluting with MeOH/DCM (5/95) to give 553 mg of the title
- 30 compound **140A** as a light yellow solid.

Example 140B

5-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-[1,3,4]oxadiazol-2-ylamine.

5


- 2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinoline-8-carboxylic acid hydrazide **140A** (403 mg, 1.07 mMol) and NaHCO_3 (275 mg, 3.28 mMol) were dissolved in a solution of 5 mL of 1,4-dioxane and 5 mL of water. To this solution was added 500 μL of a 3 M solution of cyanogen bromide in DCM. The reaction mixture was heated to reflux and reacted at this temperature for 48 hours. The reaction mixture was concentrated under vacuum and then partitioned between EtOAc and aqueous saturated NaHCO_3 . The DCM layer was then washed with brine, dried over Na_2SO_4 and filtered. A precipitate formed in the filtrate that was collected via suction filtration and dried under vacuum to give 83.7 mg of the title compound **140** as a yellow solid.
- 15 C.I. m/z 403 $[\text{M}+1]$; ^1H NMR (DMSO) δ 9.18 (s, 1 H), 8.75 (d, $J = 8.7$ Hz, 1 H), 8.68 (d, $J = 9.1$ Hz, 1 H), 8.27 (d, $J = 9.1$ Hz, 1 H), 8.20 (dd, $J = 1.2, 8.3$ Hz, 1 H), 8.12 (m, 1 H), 7.70 (m, 1 H), 7.30 (m, 1 H), 7.28 (d, $J = 2.5$ Hz, 3 H), 6.97 (dd, $J = 2.5, 8.7$ Hz, 1 H), 4.14 (m, 2 H), 3.67 (m, 2 H), 3.31 (s, 3 H).

20

Example 141

Ethyl 1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate.

25

Example 141A,B

- Ethyl 1-(8-chloroquinolin-2-yl)-benzimidazole-5-carboxylate and Ethyl 1-(8-chloroquinolin-2-yl)-benzimidazole-6-carboxylate.

30

- 2,8-Dichloroquinoline (13.3 g, 67.2 mMol), sodium hydride (60% dispersion in mineral oil, 2.75 g, 68.8 mMol), 55 mL toluene and 109 mL of 1-methyl-2-pyrrolidinone were combined under nitrogen. Ethyl benzimidazole-5-carboxylate (12.81 g, 67.4 mMol) was added portionwise over a 10 minute period, and the reaction mixture heated at reflux for 20 hr. The cooled mixture was diluted with EtOAc (~450 mL), and water (~600 mL) was added slowly to facilitate product crystallization. Filtration gave 25 g of light gray crystals, which were shown by ^1H NMR analysis to contain both title regioisomers **141A,B** in a ratio of about 3 to 2 in favor of the

35

- 5 5-carboxylate regioisomer **141A**. Fractional crystallization from chloroform/ethanol allowed separation of the regioisomers.

5'-Ester regioisomer **141A**: C.I. m/z 352 [M+1]; ¹H NMR (CDCl₃) δ 8.91 (d, J = 8.7 Hz, 1 H), 8.75 (s, 1H), 8.58 (s, 1 H), 8.40 (d, J = 8.7 Hz, 1 H), 8.22 (d, J = 8.7 Hz, 1 H), 7.92 (d, J = 7.9 Hz, 1 H), 7.80 (d, J = 8.3 Hz, 1 H), 7.78 (d, J = 8.7 Hz, 1 H), 7.51 (m, 1 H), 4.43 (q, J = - 7 Hz, 10 2 H), 1.44 (t, J = 7 Hz, 3 H).

6'-Ester regioisome **141Br**: C.I. m/z 352 [M+1]; ¹H NMR (CDCl₃) δ 9.70 (s, 1H), 8.82 (s, 1 H), 8.40 (d, J = 8.7 Hz, 1 H), 8.15 (d, J = 8.7 Hz, 1 H), 7.93 (d, J = 7.5 Hz, 1 H), 7.89 (d, J = 8.3 Hz, 1 H), 7.80 (m, 2 H), 7.51 (t, J = 7.9 Hz, 1 H), 4.44 (q, J = - 7.2 Hz, 2 H), 1.46 (t, J = 7.1 Hz, 15 3 H).

Example 141C

Ethyl-1-[8-(4-tert-butoxycarbonylaminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate.

- 20 Ethyl 1-(8-chloroquinolin-2-yl)-benzimidazole-5-carboxylate **141A** (3.00 g, 8.53 mMol), racemic-BINAP (806 mg, 1.29 mMol), Cs₂CO₃ (4.00 g, 12.3 mMol), piperidin-4-yl-carbamic acid tert-butyl ester (3.49 g, 17.4 mMol), xylenes (26 mL), and Pd(OAc)₂ (198 mg, 0.88 mMol) were combined under argon and heated at reflux for 3 days. After cooling, hexanes (25 mL) was added and the mixture stirred overnight. Solids were removed by filtration, and repeatedly
25 washed with hexanes (5 X 30 mL). The solids were then extracted with EtOAc (3 X 500 mL), and the EtOAc solutions combined and evaporated, leaving 4.18 g of greenish-black foam. The title product **141C** (2.06 g) was isolated as pale-yellow crystals (mp = 170 – 175 °C) after silica gel chromatography (hexanes/EtOAc) followed by brief ethyl ether trituration.

30

Example 141D

Ethyl 1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate.

- Ethyl-1-[8-(4-tert-butoxycarbonylaminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate **141C** (125 mg, 0.24 mMol) was stirred at ambient temperature with 4N HCl in dioxane (0.62 mL, 2.5 mMol). After 4 hr the mixture was evaporated under reduced pressure.
35 The solid residue was slurried briefly in 0.5 mL ethyl ether, filtered and vacuum dried to give the title product **141** as a hydrochloride salt.

-158-


- 5 C.I. m/z 416 [M+1]; ¹H NMR (DMSO) δ 9.38 (s, 1 H), 8.93 (d, J = 8.7 Hz, 2 H), 8.62 (d, J = 9.1 Hz, 1 H), 8.37 (s, 1H), 8.30 (d, J = 8.7 Hz, 1H), 8.21 (d, J = 8.7 Hz, 1 H), 8.16 (brs, 3 H), 7.68 (d, J = 8.3 Hz, 1 H), 7.54 (m, 1 H), 7.37 (m, 1 H), 4.36 (q, J = 7 Hz, 2 H), 3.88 (m, 2 H), 3.22 (m, 1 H), 2.84 (m, 2 H), 2.08 (m, 2 H), 1.97-2.00 (m, 2 H), 1.36 (t, J = 7 Hz, 3 H).

10

Example 142

1-[8-(4-Aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylic acid.

15

Example 142A1-[8-(4-*tert*-Butoxycarbonylaminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylic acid.

- 20 Ethyl-1-[8-(4-*tert*-butoxycarbonylaminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate **141C** (1.300 g, 2.52 mMol), 1,4-dioxane (5.2 mL), MeOH (1.3 mL) and aqueous 1N NaOH (3.9 mL, 3.9 mMol) were combined and heated at 70 °C for 1 hr. After allowing the mixture to cool, aqueous 1N HCl (3.9 mL, 3.9 mMol) was added slowly with stirring, followed by 120 mL of EtOAc. Precipitated solids were removed by filtration to give 0.916 mg of title
- 25 product **142A**. An additional 264 mg of compound **142A** was recovered upon evaporation of the filtrate EtOAc layer.

Example 142B

1-[8-(4-Aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylic acid.

30

1-[8-(4-*tert*-Butoxycarbonylaminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylic acid **142A** (51 mg, 0.105 mMol) was stirred at ambient temperature with 4N HCl in 1,4-dioxane (0.26 mL, 1.04 mMol). After 4 hr the mixture was evaporated under reduced pressure. The solid residue was slurried briefly in 1 mL Et₂O, filtered and vacuum dried to give the title

35 product **142** as a hydrochloride salt.


C.I. m/z 388 [M+1]; ¹H NMR (DMSO) δ 9.36 (s, 1 H), 8.95 (d, J = 8.3 Hz, 1 H), 8.60 (d, J = 9.1 Hz, 1 H), 8.34 (s, 1H), 8.28 (d, J = 9.1 Hz, 1H), 8.21 (d, J = 8.7 Hz, 1 H), 8.08 (br, 3 H), 7.66

-159-

- 5 (d, J = 8.3 Hz, 1 H), 7.52 (m, 1 H), 7.34 (d, J = 7.1 Hz, 1 H), 3.85-3.88 (m, 2 H), 3.20 (m, 1 H), 2.81 (m, 2 H), 2.06 (m, 2 H), 1.95-1.98 (m, 2 H).

Example 143

10 N-(4-Morpholino)ethyl-1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxamide.



15

Example 143A

- 20 N-(4-Morpholino)ethyl-1-[8-(4-*tert*-butoxycarbonylaminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxamide.
- 1-[8-(4-*tert*-Butoxycarbonylaminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylic acid **142A** (146 mg, 0.30 mMol), DMF (1 mL) and 1,1'-carbonyldiimidazole were combined and stirred under nitrogen at ambient temperature for 2 hr. 4-(2-Aminoethyl)morpholine (0.050 mL, 0.38 mMol) was added and the mixture stirred at ambient temperature for 16 hr. EtOAc (10 mL) and saturated aqueous NaHCO₃ (5 mL) were added and stirred for 30 min. The white precipitate was removed by filtration and vacuum dried to give 158 mg of title compound **143A**.

25

Example 143B

- 30 N-(4-Morpholino)ethyl-1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxamide.

- N-(4-Morpholino)ethyl-1-[8-(4-*tert*-butoxycarbonylaminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxamide **143A** (133 mg, 0.22 mMol) was stirred at ambient temperature with 4N HCl in dioxane (2.0 mL, 8 mMol). After 4 hr the mixture was evaporated under reduced pressure, 2 mL 1,4-dioxane was added and evaporated, twice, leaving a solid residue, which was slurried briefly in 3 mL Et₂O, filtered and vacuum dried to give the title product **143** as its bis-HCl salt.
- 35


-160-

- 5 C.I. m/z 500 [M+1]; ¹H NMR (DMSO) δ 9.43 (s, 1 H), 9.07-9.12 (m, 2 H), 8.61 (m, 2 H), 8.46 (br, 3 H), 8.38 (s, 1 H), 8.23 (d, J = 8.7 Hz, 1 H), 7.68 (m, 1 H), 7.54 (m, 1 H), 7.39 (m, 1H), 3.95-3.98 (m, 2 H), 3.78-3.84 (m, 4 H), 3.70-3.74 (m, 2 H), 3.57-3.68 (m, 2 H), 3.36-3.38 (m, 2 H), 3.11-3.23 (m, 3 H), 2.85 (m, 2 H), 2.11 (m, 4 H).

10

Example 144

4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazol-5-yl}-benzaldehyde.




- The same procedure that was used in example 118 was followed except that 4-
15 formylbenzeneboronic acid was used in the place of phenylboronic acid in example 118B to give the title compound **144** as the bis-HCl salt.

- C.I. m/z 448 [M+1]; ¹H NMR (DMSO) δ 10.08 (s, 1 H), 9.56 (s, 1 H), 9.08 (d, J = 8.7 Hz, 1 H), 8.64 (d, J = 8.7 Hz, 1 H), 8.40 (brs, 4 H), 8.33 (d, J = 8.7 Hz, 1 H), 8.25 (s, 1 H), 8.13 (d, J =
20 8.3 Hz, 2 H), 8.02 (d, J = 8.3 Hz, 1 H), 7.73 (d, J = 7.5 Hz, 1 H), 7.58 (m, 2 H), 7.44 (m, 1 H), 3.91 (m, 2 H), 3.26 (m, 1 H), 2.92 (m, 2 H), 2.14 (m, 4 H).

Example 145

1-{2-[5-(4-Methylaminomethyl-phenyl)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.



25

4-{1-[8-(4-amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazol-5-yl}-benzaldehyde **144** (205 mg, 0.374 mMol) was dissolved in a solution of 3 mL of MeOH and 2 mL of DCE under an atmosphere of dry N₂. To this solution was added 935 μL of a solution of 2.0 M methylamine


-161-

5 in MeOH and then AcOH was added dropwise until the pH of the solution was ~5. To this solution was added (47 mg, 0.75 mMol) NaCNBH₃ and the reaction mixture was stirred at ambient temperature for 4 hours. The reaction mixture was then concentrated under vacuum and subsequently partitioned between a solution of 2-propanol/DCM (18:82) and 0.1 N aqueous NaOH. The water layer was washed 2 more times with a solution of 2-propanol/DCM (18:82). The organic extracts were combined and concentrated under vacuum to give a yellow foam. The foam was chromatographed on flash silica gel eluting with a gradient DCM/MeOH/NH₄OH (2/97.8/0.2) to DCM/MeOH/NH₄OH (10/89.8/0.2) to give 83 mg of a yellow foam. The foam was dissolved in 2 mL of EtOH in a pressure vial. To this solution was added 50 µL of concentrated HCl. The reaction mixture was heated to ~90°C and reacted at this temperature for 2 hours. The reaction mixture was then cooled to ambient temperature and the resulting precipitate was collected via suction filtration. The solid was dried under vacuum to give 82 mg of the tri-HCl salt of the title compound **145** as a white solid.

C.I. m/z 463 [M+1]; ¹H NMR (CD₃OD) δ 10.57 (s, 1 H), 8.90 (d, J = 8.7 Hz, 1 H), 8.75 (d, J = 9.5 Hz, 1 H), 8.32 (d, J = 9.1 Hz, 1 H), 8.23 (m, 2 H), 8.16 (d, J = 8.3 Hz, 1 H), 8.03 (d, J = 8.3 Hz, 1 H), 7.94 (d, J = 8.3 Hz, 2 H), 7.86 (m, 1 H), 7.67 (d, J = 8.3 Hz, 2 H), 4.28 (s, 2 H), 4.18 (m, 2 H), 3.73 (m, 2 H), 3.63 (m, 1 H), 2.77 (s, 3 H), 2.48 (m, 2 H), 2.41 (m, 2 H).

Example 146

1-{2-[5-(4-Dimethylaminomethyl-phenyl)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-amine.



The same procedure that was used in example 145 was followed except that a solution of 2.0 M dimethylamine in MeOH was used in the place of a solution of 2.0 M methylamine in MeOH to give the tri-HCl salt of the title compound **146** as a white solid.


C.I. m/z 477 [M+1]; ¹H NMR (CD₃OD) δ 10.74 (s, 1 H), 8.97 (d, J = 9.1 Hz, 1 H), 8.70 (d, J = 8.7 Hz, 1 H), 8.39 (d, J = 9.1 Hz, 1 H), 8.28 (m, 4 H), 7.97 (d, J = 8.3 Hz, 2 H), 7.93 (m, 1 H), 7.72 (d, J = 8.3 Hz, 2 H), 4.43 (s, 2 H), 4.22 (m, 2 H), 3.97 (m, 2 H), 3.73 (m, 1 H), 2.91 (s, 6 H), 2.61 (m, 2 H), 2.49 (m, 2 H).

-162-

5

Example 147

1-(2-{5-[2-(2-Methyl-imidazol-1-yl)-ethoxy]-benzoimidazol-1-yl}-quinolin-8-yl)-piperidin-4-ylamine.



10

Example 147A

Methanesulfonic acid 2-{1-[8-(4-tert-butoxycarbonylamino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethyl ester.

15

2-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethanol **83** (1.89 g, 4.68 mMol) and di-tert-butylidicarbonate (1.02 g, 4.68 mMol) were added to a solution of 40 mL of anhydrous DCM and 20 mL of anhydrous THF under an atmosphere of dry N₂. The heterogeneous reaction mixture was stirred at ambient temperature for 3 hours after which time 1.2 mL of NEt₃ followed by 2.10 mL of methanesulphonyl chloride were added. The reaction mixture was stirred at ambient temperature for 48 hours and then partitioned between DCM and aqueous saturated NaHCO₃. The DCM layer was saved and washed 2 more times with aqueous saturated NaHCO₃, dried over Na₂SO₄, filtered and concentrated under vacuum to give a yellow foam as the title compound **147A** and was used without further purification.

25

Example 147B

1-(2-{5-[2-(2-Methyl-imidazol-1-yl)-ethoxy]-benzoimidazol-1-yl}-quinolin-8-yl)-piperidin-4-ylamine.


30 Methanesulfonic acid 2-{1-[8-(4-tert-butoxycarbonylamino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethyl ester **147A** (344 mg, 0.591 mMol) was slurried in 2.0 mL of anhydrous DMF under an atmosphere of dry N₂. To this heterogeneous solution was added 2-methylimidazole (53 mg, 0.65 mMol) followed by sodium hydride (60% in oil) (16 mg, 0.65 mMol). The reaction mixture was then heated to 60°C. After 1 hour, the reaction mixture was cooled to ambient temperature and 26 mg more of sodium hydride (60% in oil) was added.

35

- 5 The reaction mixture was reheated to 60°C and reacted at this temperature overnight. The reaction mixture was cooled to ambient temperature and then quenched with water. The reaction mixture was concentrated under vacuum and the resulting brown oil was partitioned between DCM and aqueous 1.0 M NaOH. The DCM layer was saved and washed 2 more times with aqueous 1.0 M NaOH, dried over Na₂SO₄, filtered and concentrated under vacuum.
- 10 The resulting residue was chromatographed on flash silica gel eluting with a gradient from DCM/MeOH/NH₄OH (2/97.8/0.2) to DCM/MeOH/NH₄OH (10/89.8/0.2) to give 158 mg of a yellow foam. The foam was dissolved in 1 mL of TFA under an atmosphere of dry N₂ and stirred at ambient temperature for 1 hour. The reaction mixture was concentrated under vacuum, and the resulting residue was partitioned between DCM and aqueous 0.1 M NaOH.
- 15 The water layer was washed 2 more times with DCM. The DCM extracts were combined, dried over Na₂SO₄, filtered and concentrated under vacuum to give 128 mg of the title compound **147**.
- C.I. m/z 468 [M+1]; ¹H NMR (CDCl₃) δ 8.63 (s, 1 H), 8.37 (d, J = 9.1 Hz, 1 H), 8.28 (d, J = 8.7 Hz, 1 H), 7.64 (d, J = 8.7 Hz, 1 H), 7.46 (m, 2 H), 7.26 (d, J = 2.5 Hz, 1 H), 7.22 (m, 1 H), 6.97 (m, 2 H), 6.91 (d, J = 1.3 Hz, 1 H), 4.28 (s, 4 H), 3.88 (m, 2 H), 2.88 (m, 3 H), 2.46 (s, 3 H), 2.05 (m, 2 H), 1.86 (brs, 2 H), 1.80 (m, 2 H).
- 20

Example 148

- 25 1-{2-[5-(2-[1,2,4]Triazol-1-yl-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine.



- Methanesulfonic acid 2-{1-[8-(4-tert-butoxycarbonylamino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethyl ester **147A** (500 mg, 0.860 mMol) was slurried in 2.0 mL of anhydrous DMF under an atmosphere of dry N₂. To this heterogeneous solution was added 1,2,4-triazole (65 mg, 0.95 mMol) followed by sodium hydride (60% in oil) (23 mg, 0.95 mMol). The reaction mixture was then heated to 60°C and reacted at this temperature for 3 hours. The reaction mixture was cooled to ambient temperature and then quenched with water. The reaction mixture was concentrated under vacuum and the resulting brown oil was partitioned
- 35 between a solution of 2-propanol/DCM (18:82) and aqueous 0.1 M NaOH. The organic layer

- 5 was saved and washed 2 more times with aqueous 0.1 M NaOH, dried over Na₂SO₄, filtered and concentrated under vacuum to a yellow oil. The yellow oil was chromatographed on flash silica gel eluting with DCM/MeOH/NH₄OH (1.5/98.4/0.1) to give 126 mg of a yellow foam. The foam was dissolved in 2 mL of EtOH in a pressure vial. To this solution was added 75 μ L of concentrated HCl. The reaction mixture was heated to -90°C and reacted at this temperature
- 10 for 2 hours. The reaction mixture was then cooled to ambient temperature and the resulting precipitate was collected via suction filtration. The solid was dried under vacuum to give 86 mg of the bis-HCl salt of the title compound **148** as a yellow solid.


C.I. m/z 455 [M+1]; ¹H NMR (CD₃OD) δ 10.46 (s, 1 H), 9.71 (s, 1 H), 8.84 (d, J = 9.7 Hz, 1 H), 8.73 (s, 1 H), 8.60 (d, J = 9.1 Hz, 1 H), 8.23 (d, J = 8.7 Hz, 1 H), 8.10 (d, J = 8.3 Hz, 1 H), 8.00

15 (d, J = 7.5 Hz, 1 H), 7.83 (m, 1 H), 7.54 (dd, J = 2.5, 9.1 Hz, 1 H), 7.50 (d, J = 2.1 Hz, 1 H), 4.93 (m, 2 H), 4.67 (m, 2 H), 4.12 (m, 2 H), 3.60 (m, 3 H), 2.37 (m, 4 H).

5

CLAIMS

1. A compound of the formula



or a pharmaceutically acceptable salt, prodrug or solvate thereof, wherein:

wherein X is CH or N;

- 10 R^1 is selected from $-(CR^4R^5)_tC(O)OR^3$, $-(CR^4R^5)_tC(O)NR^3R^4$, $-(CR^4R^5)_tOR^3$, $-(CR^4R^5)_tC(O)(C_3-C_{10} \text{ cycloalkyl})$, $-(CR^4R^5)_tC(O)(C_6-C_{10} \text{ aryl})$, $-(CR^4R^5)_tC(O)(4 \text{ to } 10 \text{ membered heterocyclic})$, $-(CR^4R^5)_t(C_3-C_{10} \text{ cycloalkyl})$, $-(CR^4R^5)_t(C_6-C_{10} \text{ aryl})$, and $-(CR^4R^5)_t(4 \text{ to } 10 \text{ membered heterocyclic})$, wherein each t is independently an integer from 0 to 5; said cycloalkyl, aryl and heterocyclic R^1 moieties are optionally fused to a benzene ring, a C_5-C_8
- 15 cycloalkyl group, or a 4 to 10 membered heterocyclic group; the $-(CR^4R^5)_t$ moieties of the foregoing R^1 groups optionally include a carbon-carbon double or triple bond where t is an integer between 2 and 5; the foregoing R^1 groups are each optionally substituted by 1 or 2 groups independently selected from $-NR^3R^4$, $-OR^3$, C_1-C_{10} alkyl, C_2-C_{10} alkenyl, and C_2-C_{10} alkynyl, wherein said alkyl, alkenyl and alkynyl groups are substituted by 1 or 2 groups
- 20 independently selected from $-NR^3R^4$ and $-OR^3$; and the foregoing R^1 groups are optionally substituted by 1 to 3 R^2 groups;

- each R^2 is independently selected from H, C_1-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_3-C_{10} cycloalkyl, oxo, halo, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, $-OR^3$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)OR^6$, $-OC(O)R^3$, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$,
- 25 $-NR^4C(O)R^3$, $-C(O)NR^3R^4$, $-NR^5C(O)NR^3R^4$, $-NR^3R^4$, $-S(O)_j(CR^4R^5)_m(C_6-C_{10} \text{ aryl})$, $-S(O)_j(C_1-C_6 \text{ alkyl})$, wherein j is an integer from 0 to 2, $-(CR^4R^5)_m(C_6-C_{10} \text{ aryl})$, $-O(CR^4R^5)_m(C_6-C_{10} \text{ aryl})$, $-NR^4(CR^4R^5)_m(C_6-C_{10} \text{ aryl})$, $-O(CR^4R^5)_m(4 \text{ to } 10 \text{ membered heterocycle})$, $-NR^4(CR^4R^5)_m(4 \text{ to } 10 \text{ membered heterocycle})$, $-(CR^4R^5)_m(4 \text{ to } 10 \text{ membered heterocyclic})$, and $-(CR^4R^5)_m(C_3-C_{10} \text{ cycloalkyl})$ wherein each m is independently an integer from 0 to 4; said alkyl, alkenyl and
- 30 alkynyl groups optionally contain 1 or 2 hetero moieties selected from O, $-S(O)_j$ wherein j is an integer from 0 to 2, and $-N(R^3)-$ with the proviso that two O atoms, two S atoms, or an O and S atom are not attached directly to each other, and the proviso that an O atom, a S atom or a N atom are not attached directly to a triple bond or a non-aromatic double bond; said cycloalkyl, aryl and heterocyclic R^2 groups are optionally fused to a C_6-C_{10} aryl group, a C_5-C_8
- 35 cycloalkyl group, or a 4 to 10 membered heterocyclic group; and said alkyl, cycloalkyl, aryl

- 5 and heterocyclic R^2 groups are optionally substituted by 1 to 5 substituents independently selected from oxo ($=O$), halo, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-C(O)R^3$, $-C(O)OR^3$, $-OC(O)R^3$, $-NR^4C(O)OR^6$, $-NR^4C(O)R^3$, $-C(O)NR^3R^4$, $-NR^3R^4$, $-OR^3$, C_1-C_{10} alkyl, $-(CR^4R^5)_m(C_6-C_{10}$ aryl), and $-(CR^4R^5)_m$ (4 to 10 membered heterocyclic), wherein each m is independently an integer ranging from 0 to 4;
- 10 each R^3 is independently selected from H, C_1-C_{10} alkyl, $-(CR^4R^5)_m(C_6-C_{10}$ aryl), and $-(CR^4R^5)_m$ (4 to 10 membered heterocyclic), wherein each m is independently an integer from 0 to 4; said alkyl group optionally includes 1 or 2 hetero moieties selected from O, $-S(O)_j-$ wherein j is an integer ranging from 0 to 2, and $-N(R^4)-$ with the proviso that two O atoms, two S atoms, or an O and S atom are not attached directly to each other; said cycloalkyl, aryl and
- 15 heterocyclic R^3 groups are optionally fused to a C_6-C_{10} aryl group, a C_5-C_8 cycloalkyl group, or a 4 to 10 membered heterocyclic group; and the foregoing R^3 substituents, except H, are optionally substituted by 1 to 5 substituents independently selected from oxo, halo, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, $-C(O)R^4$, $-C(O)OR^4$, $-OC(O)R^4$, $-NR^4C(O)R^5$, $-C(O)NR^4R^5$, $-NR^4R^5$, hydroxy, C_1-C_6 alkyl, and C_1-C_6 alkoxy;
- 20 each R^4 and R^5 is independently H or C_1-C_6 alkyl;
- or, where R^4 and R^5 are attached to the same carbon or nitrogen atom, R^4 and R^5 together with said carbon or nitrogen may be taken together to form a 4 to 10 membered ring which may be carbocyclic or heterocyclic;
- each R^6 is selected from the substituents provided in the definition of R^3 except R^6 is
- 25 not H;
- each R^7 , R^8 , R^9 , R^{10} and R^{11} is independently selected from the group of substituents provided in the definition of R^2 .
2. A compound according to claim 1 wherein R^1 is C_6-C_{10} aryl or 4 to 10 membered heterocyclic wherein the foregoing R^1 groups are each substituted by 1 or 2 groups
- 30 independently selected from $-NR^3R^4$, $-OR^3$ and C_1-C_3 alkyl, wherein said alkyl groups are substituted by 1 or 2 groups independently selected from $-NR^3R^4$ and $-OR^3$; and the foregoing R^1 groups are optionally substituted by 1 to 3 R^2 groups;
- each R^2 is independently selected from H, C_1-C_{10} alkyl, C_3-C_{10} cycloalkyl, oxo ($=O$), $-OR^3$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)R^3$, $-C(O)NR^3R^4$, $-NR^3R^4$, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$,
- 35 $-(CR^4R^5)_m$ (4 to 10 membered heterocyclic), and $(CR^4R^5)_m(C_3-C_{10}$ cycloalkyl); and said alkyl groups optionally contain 1 or 2 hetero moieties selected from O, $-S(O)_j-$ wherein j is an integer from 0 to 2, and $-N(R^3)-$ with the proviso that two O atoms, two S atoms, or an O and S atom are not attached directly to each other; and said alkyl and cycloalkyl R^2 groups are
- 40 optionally substituted by 1 to 5 substituents independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, $-NR^4SO_2R^6$, $-SO_2NR^3R^4$, $-C(O)R^3$, $-C(O)OR^3$, $-NR^4C(O)OR^6$,

5 -NR⁴C(O)R³, -C(O)NR³R⁴, -NR³R⁴, -OR³, and C₁-C₁₀ alkyl, wherein each m is independently an integer ranging from 0 to 4.

3. A compound according to claim 1 wherein R¹ is piperidinyl, piperziny, or phenyl, wherein said R¹ groups are substituted by -NR³R⁴, oxo (=O), -OR³ and C₁-C₃ alkyl, wherein said alkyl group is optionally substituted by 1 or 2 groups independently selected from -NR³R⁴ and -OR³; and the foregoing R¹ groups are optionally substituted by 1 to 3 R² groups.

4. A compound according to claim 2 wherein said R¹ groups are substituted by -NR³R⁴, oxo (=O), OR³, or C₁-C₃ alkyl, wherein said alkyl group is optionally substituted by -NR³R⁴.

5. A compound according to claim 1 wherein R¹ is phenyl substituted by pyrrolidin-1-yl which pyrrolidin-1-yl is optionally substituted by 1 to 3 substituents independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, -NR⁴SO₂R⁶, -SO₂NR³R⁴, -C(O)R³, -C(O)OR³, -NR⁴C(O)OR⁶, -NR⁴C(O)R³, -C(O)NR³R⁴, -NR³R⁴, -OR³, and C₁-C₁₀ alkyl; and R¹¹ is -OR³.

6. A compound according to claim 5 wherein R¹ is 4-pyrrolidin-1-ylmethyl-phenyl optionally substituted by 1 to 3 substituents independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, -NR⁴SO₂R⁶, -SO₂NR³R⁴, -C(O)R³, -C(O)OR³, -NR⁴C(O)OR⁶, -NR⁴C(O)R³, -C(O)NR³R⁴, -NR³R⁴, -OR³, and C₁-C₁₀ alkyl; and R¹¹ is -OR³.

7. A compound according to claim 6 wherein R¹¹ is connected at the 5 position of the benzimidazole moiety of the compound of formula 1 and is -OR³.

8. A compound according to claim 7 wherein R¹¹ is connected at the 5 position of the benzimidazole moiety of the compound of formula 1 and is 2-methoxyethoxy.

9. A compound according to claim 1 wherein R¹ is pyrrolidin-1-yl or piperidin-1-yl, said R¹ being optionally substituted by 1 to 3 substituents independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, -NR⁴SO₂R⁶, -SO₂NR³R⁴, -C(O)R³, -C(O)OR³, -NR⁴C(O)OR⁶, -NR⁴C(O)R³, -C(O)NR³R⁴, -NR³R⁴, -OR³, and C₁-C₁₀ alkyl.

10. A compound according to claim 9 wherein R¹ is pyrrolidin-1-yl or piperidin-1-yl substituted by -NR³R⁴ and optionally substituted by 1 or 2 substituents independently selected from oxo, cyano, trifluoromethyl, trifluoromethoxy, -NR⁴SO₂R⁶, -SO₂NR³R⁴, -C(O)R³, -C(O)OR³, -NR⁴C(O)OR⁶, -NR⁴C(O)R³, -C(O)NR³R⁴, -NR³R⁴, -OR³, and C₁-C₁₀ alkyl; and R¹¹ is -OR³.

11. A compound according to claim 10 or claim 9 wherein R¹¹ is connected at the 5 position of the benzimidazole moiety of the compound of formula 1 and is -OR³ and R⁹ and R¹⁰ are both H.

- 5 12. A compound according to claim 10 or claim 9 wherein R¹¹ is connected at the 5 position of the benzimidazole moiety of the compound of formula 1 and is 2-methoxyethoxy and R⁹ and R¹⁰ are both H.
13. A compound according to claim 1 wherein R⁹ is -C(O)R³ wherein R³ is pyrrolidin-1-yl or azetidin-1-yl wherein said R³ groups are optionally substituted by 1 to 3 substituents
10 independently selected from oxo, halo, cyano, nitro, trifluoromethyl, difluoromethoxy, trifluoromethoxy, azido, -C(O)R⁴, -C(O)OR⁴, -OC(O)R⁴, -NR⁴C(O)R⁵, -C(O)NR⁴R⁵, -NR⁴R⁵, hydroxy, C₁-C₈ alkyl, and C₁-C₈ alkoxy.
14. A compound according to claim 1 selected from the group consisting of:
[1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
15 1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzimidazol-5-ol;
1-[2-[5-(Pyridin-2-ylmethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
{1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethyl-amine;
{4-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine;
{4-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine;
20 Cyclopropyl-{4-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine;
tert-Butyl-{4-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine;
4-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzylamine;
1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
{1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethylamine;
25 1-[2-(5-Trifluoromethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
{4-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine;
Cyclopropyl-{4-[2-(5-ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine;
tert-Butyl-{4-[2-(5-ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-amine;
{4-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine;
30 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one;
1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-one;
1-[2-(5-Ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
tert-Butyl-{1-[2-(5-ethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-amine;
{1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine;
35 2-(5-Methoxy-benzimidazol-1-yl)-8-(1-oxa-6-aza-spiro[2.5]oct-6-yl)-quinoline;
4-Dimethylaminomethyl-1-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-4-methylaminomethyl-piperidin-4-ol;
40 4-Aminomethyl-1-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;

- 5 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-pyrrolidin-3-ylamine;
 1-(2-Benzimidazol-1-yl-quinolin-8-yl)-piperidin-4-ylamine;
 1-(2-Imidazo[4,5-b]pyridin-3-yl-quinolin-8-yl)-piperidin-4-ylamine;
 1-(2-[5-(4-Methoxy-phenyl)-benzimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine;
 1-[2-(5-Cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
 10 1-(2-[5-(3-Dimethylamino-propoxy)-benzimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine;
 1-(2-[5-(3-Amino-propoxy)-benzimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine;
 1-(2-[5-(2-Dimethylamino-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine;
 15 1-(2-[5-(Pyridin-4-ylmethoxy)-benzimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine;
 1-(2-[5-(Benzyloxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
 1-(2-[5-(Pyridin-3-ylmethoxy)-benzimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine;
 1-(2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine;
 1-(2-[5-Methoxy-benzimidazol-1-yl]-quinolin-8-yl)-piperidine-4-carboxylic acid ethyl
 20 ester;
 1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidine-4-carboxylic acid;
 4-Dimethylaminomethyl-1-[2-(5-methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
 N-(1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl)-acetamide;
 25 N-(1-[2-(5-Cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl)-acetamide;
 1-(2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ol;
 {1-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-urea;
 4-Aminomethyl-1-[2-[5-(pyridin-2-ylmethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ol;
 30 Cyclopropyl-(1-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-amine;
 (1-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-dimethyl-amine;
 35 1-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-methyl-amine;
 (1-[2-[5-(3-Dimethylamino-propoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-dimethyl-amine;
 {1-[2-(5-Cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl)-methyl-amine;
 40

- 5 {1-[2-(5-Cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-
dimethyl-amine;
2-Amino-N-(1-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-
yl)- acetamide;
- (S)-2-Amino-N-(1-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-
10 piperidin-4-yl)-propionamide;
- (R)-2-Amino-N-(1-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-
piperidin-4-yl)-propionamide;
2-Amino-N-(1-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-
yl)-isobutyramide;
15 1-(1-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamino)-
2-methyl-propan-2-ol;
(1-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-pyridin-
2-ylmethyl-amine;
(1-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperidin-4-yl)-pyridin-
20 3-ylmethyl-amine;
4-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-phenol;
[2-(4-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-phenoxy)-ethyl]-
dimethyl-amine;
2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-8-piperazin-1-yl-quinoline;
25 [2-(4-[2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperazin-1-yl)-
ethyl]-dimethyl-amine;
2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-8-(4-pyridin-2-ylmethyl-piperazin-1-yl)-
quinoline;
2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-8-(4-pyridin-3-ylmethyl-piperazin-1-yl)-
30 quinoline;
2-Amino-1-(4-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperazin-1-
yl)-2-methyl-propan-1-one;
(S)-2-Amino-1-(4-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-
piperazin-1-yl)-propan-1-one;
35 (S)-2-Amino-1-(4-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-
piperazin-1-yl)-propan-1-one;
2-Amino-1-(4-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-piperazin-1-
yl)-ethanone;
(1-Amino-cyclopropyl)-(4-[2-[5-(2-methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl]-
40 piperazin-1-yl)-methanone;

-171-

- 5 2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethylamine;
(R)-2-Amino-3-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-propan-1-ol;
3-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-3-aza-bicyclo[3.1.0]hex-
10 6-ylamine;
(S)-1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-pyrrolidin-3-ylamine;
(R)-1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-pyrrolidin-3-ylamine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-pyridin-3-yl-quinoline;
15 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(6-methoxy-pyridin-3-yl)-quinoline;
4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzoic acid methyl ester;
1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-4-methyl-piperidin-4-ylamine;
20 1-[2-(6,7-Dihydro-5,8-dioxa-1,3-diaza-cyclopenta[b]naphthalen-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
2-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethanol;
4-Cyclopropylaminomethyl-1-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol;
25 1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazole-5-sulfonic acid dimethylamide;
1-[2-(6-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-[2-(5,6-Dimethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
2-Dimethylamino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-ethanone;
30 1-[2-(5-Benzyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-4-methyl-piperidin-4-ol;
(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-dimethylamine;
(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-methylamine;
35 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-morpholin-4-ylmethyl-phenyl)-quinoline;
2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamino)-ethanol;
4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamine;

- 5 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
- 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
- 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
- 10 1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-cis-pyrrolidine-3,4-diol;
- R,R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol);
- 15 1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol;
- R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol);
- S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol);
- 20 1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-azetidin-3-ol;
- 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-quinoline;
- 4-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperazine-1-carboxylic acid tert-butyl ester;
- 25 [1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-yl]-carbamic acid tert-butyl ester;
- 1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-ylamine;
- 30 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methanol;
- (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-methyl-amine;
- 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-piperidin-1-yl]-quinoline;
- 35 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-dimethyl-amine;
- 1-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-pyrrolidin-3-ol;

- 5 C-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methylaniline;
1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-4-methyl-piperidin-4-ol;
1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ol;
- 10 S,S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol);
4-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
4-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
1-[2-(5-Phenyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
- 15 1-[2-(5-Pyridin-4-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-[2-[5-(3-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
1-[2-(5-Pyridin-3-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-[2-[5-(6-Methoxy-pyridin-3-yl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
1-[2-[5-(4-Aminomethyl-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-
- 20 yllamine;
4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl}-benzoic acid methyl ester;
4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl}-phenol;
2-(5-Methoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester;
- 25 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinoline-8-carboxylic acid methyl ester;
2-(5-Methoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid (2-dimethylamino-ethyl)-amide;
2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinoline-8-carboxylic acid methyl ester;
- 30 [2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-pyrrolidin-1-yl-methanone;
[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-morpholin-4-yl-methanone;

- 5 [2-(5-Cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-piperidin-1-yl-methanone;
(3-Amino-pyrrolidin-1-yl)-[2-(5-cyclopropylmethoxy-benzimidazol-1-yl)-quinolin-8-yl]-methanone;
8-Allyloxy-2-(5-methoxy-benzimidazol-1-yl)-quinoline;
- 10 {2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethyl}-methyl-amine;
{2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethyl}-dimethyl-amine;
2-[2-(5-Methoxy-benzimidazol-1-yl)-quinolin-8-yloxy]-ethylamine;
1-[[2-[5-(3-Morpholinoethoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride;
- 15 1-[[2-[5-(3-Morpholinoethoxy)-1*H*-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride;
5-{2-[5-(2-Methoxy-ethoxy)-benzimidazol-1-yl]-quinolin-8-yl}-[1,3,4]oxadiazol-2-ylamine;
Ethyl 1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate;
- 20 1-[8-(4-Aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylic acid;
N-(4-Morpholino)ethyl-1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxamide;
4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1*H*-benzimidazol-5-yl}-benzaldehyde;
- 25 1-{2-[5-(4-Methylaminomethyl-phenyl)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-{2-[5-(4-Dimethylaminomethyl-phenyl)-benzimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-(2-{5-[2-(2-Methyl-imidazol-1-yl)-ethoxy]-benzimidazol-1-yl}-quinolin-8-yl)-piperidin-4-ylamine and
- 30 1-{2-[5-(2-[1,2,4]Triazol-1-yl)-ethoxy]-benzimidazol-1-yl}-quinolin-8-yl}-piperidin-4-ylamine,
and the pharmaceutically acceptable salts, prodrugs and solvates of the foregoing compounds.

15. A compound according to claim 1 selected from the group consisting of:

- 5 1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-ol;
1-{2-[5-(Pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
{1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethyl-amine;
{4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine;
10 {4-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine;
1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
{1-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-dimethylamine;
{4-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-methyl-amine;
{4-[2-(5-Ethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-benzyl}-dimethyl-amine;
15 {1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-amine;
4-Dimethylaminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
1-[2-(5-Methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-4-methylaminomethyl-piperidin-4-ol;
20 4-Aminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
1-{2-[5-(4-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
25 1-{2-[5-(3-Amino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-{2-[5-(2-Dimethylamino-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-{2-[5-(Pyridin-4-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-[2-(5-Benzyloxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
30 1-{2-[5-(Pyridin-3-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
4-Dimethylaminomethyl-1-[2-(5-methoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ol;
4-Aminomethyl-1-[2-[5-(pyridin-2-ylmethoxy)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ol;
35 1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl}-dimethyl-amine;
1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl}-methyl-amine;

- 5 (1-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-
dimethyl-amine;
{1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-methyl-
amine;
{1-[2-(5-Cyclopropylmethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-yl}-
10 dimethyl-amine;
1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-pyridin-2-
ylmethyl-amine;
(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-pyridin-
3-ylmethyl-amine;
15 4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
[2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenoxy)-ethyl]-
dimethyl-amine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-piperazin-1-yl-quinoline;
[2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-
20 ethyl]-dimethyl-amine;
2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-
yl)-2-methyl-propan-1-one;
(S)-2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-
piperazin-1-yl)-propan-1-one;
25 (S)-2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-
piperazin-1-yl)-propan-1-one;
2-Amino-1-(4-{2-[5-(2-methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-
yl)-ethanone;
2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperazin-1-yl)-
30 ethylamine;
3-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-3-aza-bicyclo[3.1.0]hex-
6-ylamine;
1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-4-methyl-piperidin-4-
ylamine;
35 2-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yloxy}-ethanol;
1-[2-(5,6-Dimethoxy-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
1-[2-(6,7-Dihydro-5,8-dioxo-1,3-diaza-cyclopenta[b]naphthalen-1-yl)-quinolin-8-yl]-
piperidin-4-ylamine;

- 5 (4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-dimethyl-amine;
(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-methyl-amine;
2-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamino)-ethanol;
- 10 4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzylamine;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
- 15 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-(4-pyrrolidin-1-ylmethyl-phenyl)-quinoline;
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-cis-pyrrolidine-3,4-diol;
R,R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol);
- 20 1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol;
R-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol);
- 25 S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-pyrrolidin-3-ol);
1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-azetidin-3-ol
2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-quinoline;
- 30 1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-piperidin-4-ylamine;
(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methanol;
(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-methyl-amine;
- 35 2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-8-[4-(4-methyl-piperazin-1-ylmethyl)-piperidin-1-yl]-quinoline;

- 5 (1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylmethyl)-dimethyl-amine;
- C-(1-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-yl)-methylamine;
- 10 S,S-(1-(4-{2-[5-(2-Methoxy-ethoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-benzyl)-trans-pyrrolidine-3,4-diol);
- 4-{2-[5-(3-Dimethylamino-propoxy)-benzoimidazol-1-yl]-quinolin-8-yl}-phenol;
- 1-[2-(5-Phenyl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
- 1-[2-(5-Pyridin-4-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
- 1-{2-[5-(3-Methoxy-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 15 1-[2-(5-Pyridin-3-yl-benzoimidazol-1-yl)-quinolin-8-yl]-piperidin-4-ylamine;
- 1-[2-[5-(6-Methoxy-pyridin-3-yl)-benzoimidazol-1-yl]-quinolin-8-yl]-piperidin-4-ylamine;
- 1-{2-[5-(4-Aminomethyl-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 20 4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl}-benzoic acid methyl ester;
- 4-{1-[8-(4-Amino-piperidin-1-yl)-quinolin-2-yl]-1H-benzoimidazol-5-yl}-phenol;
- 1-[[2-[5-(3-Morpholinoethoxy)-1H-benzimidazol-1-yl]-quinolin-8-yl]]-piperidin-4-ylamine trihydrochloride;
- Ethyl 1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxylate;
- 25 N-(4-Morpholino)ethyl-1-[8-(4-aminopiperidin-1-yl)-quinolin-2-yl]-benzimidazole-5-carboxamide;
- 1-{2-[5-(4-Methylaminomethyl-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 1-{2-[5-(4-Dimethylaminomethyl-phenyl)-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine;
- 30 1-(2-[5-[2-(2-Methyl-imidazol-1-yl)-ethoxy]-benzoimidazol-1-yl]-quinolin-8-yl)-piperidin-4-ylamine and
- 1-{2-[5-(2-[1,2,4]Triazol-1-yl)-ethoxy]-benzoimidazol-1-yl]-quinolin-8-yl}-piperidin-4-ylamine,

35 and the pharmaceutically acceptable salts, prodrugs and solvates of the foregoing compounds.

16. A method for the treatment of abnormal cell growth in a mammal comprising administering to said mammal an amount of a compound of claim 1 that is effective in treating abnormal cell growth.

17. A method according to claim 16 wherein said abnormal cell growth is cancer.

5 18. A method according to claim 17 wherein said cancer is selected from lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of
10 the vulva, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute leukemia, lymphocytic lymphomas, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasms of the central nervous
15 system (CNS), primary CNS lymphoma, spinal axis tumors, brain stem glioma, pituitary adenoma, or a combination of one or more of the foregoing cancers.

 19. A pharmaceutical composition for the treatment of abnormal cell growth in a mammal comprising an amount of a compound of claim 1 that is effective in treating abnormal cell growth, and a pharmaceutically acceptable carrier.

20 20. The pharmaceutical composition of claim 19 wherein said abnormal cell growth is cancer.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/IB 00/01636

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/14 A61K31/47 A61P35/00 C07D401/04 C07D491/10
C07D471/04 C07D491/04 C07D413/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 16755 A (HUNGATE RANDALL W ;KOESTER TIMOTHY J (US); BILODEAU MARK T (US); M) 8 April 1999 (1999-04-08) claim 1	1-19
Y	WO 92 20642 A (RHONE POULENC RORER INT) 26 November 1992 (1992-11-26) cited in the application claim 5	1-19
Y	PALMER ET AL.: "Structure-Activity relationship for 5-substituted 1-phenylbenzimidazoles ..." J.MED.CHEM., vol. 42, 1999, pages 2373-2382, XP002157335 table 1	1-19

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

15 January 2001

Date of mailing of the international search report

22/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Steendijk, M

INTERNATIONAL SEARCH REPORT

In International Application No

PCT/IB 00/01636

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>PALMER ET AL.: "Structure-Activity relationships for 1-phenylbenzimidazoles as selective ATP site inhibitors..." J.MED.CHEM., vol. 41, 1998, pages 5457-5465, XP002157336 table 2</p> <p>-----</p>	1-19

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 00/01636

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9916755 A	08-04-1999	AU 9500398 A EP 1017682 A US 6162804 A	23-04-1999 12-07-2000 19-12-2000
WO 9220642 A	26-11-1992	AT 159009 T AU 658646 B AU 1993492 A CA 2102780 A DE 69222637 D DE 69222637 T DK 584222 T EP 0584222 A ES 2108120 T GR 3024955 T JP 6507643 T MX 9202181 A SG 64322 A US 5409930 A US 5480883 A US 5710158 A US 5795889 A US 5646153 A US 5721237 A US 5714493 A US 6057320 A US RE36256 E	15-10-1997 27-04-1995 30-12-1992 11-11-1992 13-11-1997 26-02-1998 23-02-1998 02-03-1994 16-12-1997 30-01-1998 01-09-1994 01-05-1993 27-04-1999 25-04-1995 02-01-1996 20-01-1998 18-08-1998 08-07-1997 24-02-1998 03-02-1998 02-05-2000 20-07-1999